

졸업논문청구논문

Orion A Cloud의 쌍극 방출류의 성질

Properties of bipolar outflows of the Orion A Cloud

이 선 재 (李 善 在 Lee, Seon Jae)

16072

과학영재학교 경기과학고등학교

2019

Orion A Cloud의 쌍극 방출류의 성질

Properties of bipolar outflows of the Orion A Cloud

[논문제출 전 체크리스트]

1. 이 논문은 내가 직접 연구하고 작성한 것이다. ☒
2. 인용한 모든 자료(책, 논문, 인터넷자료 등)의 인용표시를 바르게 하였다. ☒
3. 인용한 자료의 표현이나 내용을 왜곡하지 않았다. ☒
4. 정확한 출처제시 없이 다른 사람의 글이나 아이디어를 가져오지 않았다. ☒
5. 논문 작성 중 도표나 데이터를 조작(위조 혹은 변조)하지 않았다. ☒
6. 다른 친구와 같은 내용의 논문을 제출하지 않았다. ☒

Properties of bipolar outflows of the Orion A Cloud

Advisor : Teacher Park, Kie Hyun

by

16072 Lee, Seon Jae

Gyeonggi Science High School for the gifted

A thesis submitted to the Gyeonggi Science High School in partial fulfillment of the requirements for the graduation. The study was conducted in accordance with Code of Research Ethics.*

2018. 7. 21.

**Approved by
Teacher Park, Kie Hyun
[Thesis Advisor]**

*Declaration of Ethical Conduct in Research: I, as a graduate student of GSHS, hereby declare that I have not committed any acts that may damage the credibility of my research. These include, but are not limited to: falsification, thesis written by someone else, distortion of research findings or plagiarism. I affirm that my thesis contains honest conclusions based on my own careful research under the guidance of my thesis advisor.

Orion A Cloud의 쌍극 방출류의 성질

이 선 재

위 논문은 과학영재학교 경기과학고등학교 졸업논문으로
졸업논문심사위원회에서 심사 통과하였음.

2018년 7월 21일

심사위원장 박 용 선 (인)

심사위원 강 동 일 (인)

심사위원 박 기 현 (인)

Properties of bipolar outflows of the Orion A Cloud

Abstract

Stars are born when matter from interstellar molecular clouds fall to its center to increase the mass of the protostar. Bipolar outflows are formed to remove the excess angular momentum of falling matter. Intensities of outflows are known as to be in a close relationship with their bolometric luminosity and evolutionary stages. In this paper, data from Institute for Radio Astronomy in the Millimeter Range (IRAM) 30m Telescope and Taeduk Radio Astronomy Observatory (TRAO) are used. IRAM data were used to map $^{12}\text{CO J} = 2 - 1$ over Orion A molecular cloud. TRAO data were used to map $^{13}\text{CO J} = 1 - 0$ over the same region. Outflows were observed and measured by drawing contour maps and line profiles of red/blue shifted components. The correlation between a protostar's luminosity and outflow momentum flux have been confirmed. Also, outflows could be detected better if the energy level of the emission

line is higher.

Orion A Cloud의 쌍극 방출류의 성질

초 록

별은 성간분자운의 물질이 중심으로 떨어져 원시성의 질량을 증가시켜야만 탄생된다. 이 과정에서 중심으로 떨어지는 물질의 각운동량을 제거하기 위해 방출류가 발생한다. 여기서 방출류의 세기는 원시성의 진화 단계와 광도와 관련이 있다고 알려져 있다. 이를 새로 관측된 데이터를 사용하여 보다 좋은 방출류 측정 방법과 기존의 연구를 검증해 보려고 한다. 이 연구에서는 Institute for Radio Astronomy in the Millimeter Range (IRAM) 30m 망원경으로 관측한 $^{12}\text{CO } J=2-1$ 관측 자료와 대덕 전파 망원경(Taeduk Radio Astronomy Observatory, TRA0)으로 관측한 $^{13}\text{CO } J=1-0$ 천이 선 자료를 이용하였다. 두 자료 모두 Orion A Cloud 영역을 담고 있다. 빠른 속도 성분을 가진 적색/청색편이된 성분의 contour map을 그려 방출류를 관찰하고 방출류의 세기를 구하였다. 방출류의 세기와 원시성의 광도가 대체적으로 비례한다는 것을 알 수 있었다. 그리고 천이 선의 에너지 준위가 높을수록 방출류를 더 잘 검출할 수 있음을 확인할 수 있었다.

Contents

Abstract	i
Contents	iv
List of Figures	v
List of Tables	vi
1 Introduction	1
2 Observations and Data Reduction	2
2.1 Observation Region	2
2.2 Observation Data	2
2.3 Identification of Outflows	3
3 Results	5
3.1 Outflow Identification	5
3.1.1 $^{12}\text{CO J} = 2 - 1$ Observations	5
3.1.2 $^{12}\text{CO J} = 1 - 0$ Observations	9
3.2 Momentum Flux	10
3.3 Momentum flux vs. Bolometric luminosity	11
3.4 Momentum flux by emission line energy level	12
4 Conclusion	14
References	15
감사의 글	17
연구활동	18

List of Figures

Figure 1.	Orion A ^{12}CO ($J = 2 - 1$) integrated intensity map(left) ^{13}CO integrated intensity map(right).	3
Figure 2.	The contour map(left) and the line profile(right) of FIR2.	5
Figure 3.	The contour map and the line profile of FIR3.	6
Figure 4.	The contour map and the line profile of FIR6b.	6
Figure 5.	The contour map and the line profile of MMS2.	7
Figure 6.	The contour map and the line profile of MMS5.	7
Figure 7.	The contour map and the line profile of MMS9.	8
Figure 8.	The contour map of FIR2(left), FIR3(middle), and MMS9(right).	9
Figure 9.	Momentum flux difference by emission line energy.	11
Figure 10.	Momentum flux difference by emission line energy.	12

List of Tables

Table 1.	Protostars with observed outflows.	5
Table 2.	CO outflow parameters.	10

I. Introduction

Stars are formed in molecular clouds by gravitational accretion. In the early stages of star formation, young stellar objects (YSOs) are still embedded in the molecular clouds, increasing its mass and temperature by accretion of interstellar medium around it. Since the angular momentum is conserved while matter is accreted, matter near the surface of the protostar spins quickly, which stops more accretion. Since angular momentum is removed by jets called bipolar outflows, outflows are observed with size proportional to the mass accreted to the protostar [1]. It is already known that the accretion rate and the luminosity correlates to each other [2]. The outflow force decreases as protostars evolve from Class 0 to Class I, which means the strength that the protostar pulls interstellar matter decreases as time passes. In this study, I observed the protostars and their outflows of Orion A Cloud. First, I selected the protostars that outflows can be detected from the Spitzer and the Herchel catalogues [3, 4]. By using different data sets observed by different observatories and different wavelengths, I identified the outflows. I rechecked the correlation between the outflow force and its bolometric luminosity. Also, I compared the outflow momentum flux calculated using different emission lines.

II. Observations and Data Reduction

II.1 Observation Region

The Orion region consists of two giant molecular clouds, the Orion A and B clouds. This research covers the Orion A Cloud. The Orion A Cloud covers about 29 deg^2 of the sky and its distance is about 450 pc [5]. The total mass is estimated to be about $10^5 M_{\odot}$. It contains several hot molecular cores, such as the BN-KL nebula. It is known that the Orion Cloud was formed by a collision and fragmentation between two giant molecular clouds about 60 million years ago. The effects of the collision can be seen nowadays. There is a big velocity gradient along the declination axis. On the north side of the Orion A Cloud (OMC 2) shows about 12 km s^{-1} but on the south end (L1641) it has velocity about 5 km s^{-1} [6].

II.2 Observation Data

The ^{12}CO ($J = 2 - 1$, 230.538 GHz) data was observed with the IRAM 30 m telescope in Granada, Spain, in 2013. The spatial beamwidth was $11''$, and the spectral resolution was 0.4 km s^{-1} . The noise level was 0.2 K. It only covers the north region of the Orion A cloud [7].

The ^{12}CO ($J = 1 - 0$, 115.271 GHz) data was observed with the NRO 45m telescope in Nobeyama, Japan.

The ^{13}CO ($J = 1 - 0$, 110.201 GHz) and the C^{18}O ($J = 1 - 0$, 109.782 GHz) was observed at Taeduk Radio Astronomy Observatory (TRAO) 13.7 m telescope in 2017. The spatial beamwidth was $45''$, and the spectral resolution was 0.05 km s^{-1} . The noise level was 0.4 K.

^{13}CO and C^{18}O lines are optically thin lines which can trace most of the matter on the line of sight, contrasting to ^{12}CO lines which are so optically oblique that it can only trace the outermost part of the molecular core. In this study, I used TRAO data to determine the protostar's

velocity and linewidth which are the kinematic properties of the envelope. Then, I will trace the outflow jets using ^{12}CO data.

(박기현 코멘트) Figure 1 언급해주세요.




Figure 1. Orion A ^{12}CO ($J = 2 - 1$) integrated intensity map(left) ^{13}CO integrated intensity map(right).

II.3 Identification of Outflows

Data obtained by observing with radio waves are summed over the line of sight, which tells us the distribution of matter which has relative radial velocity to the observer. The envelope around the protostar is static or is contracting slowly to the protostar itself, but outflow jets have big velocity components from each pole. If the inclination of the outflows are not zero, it would be seen as jets are moving closer or further from the observer. In this study, ^{13}CO and C^{18}O lines were used to get the velocity distribution of the protostar. By using Gaussian fitting, I calculated the protostar's central velocity(v_{cen}) and the full width at half maximum (FWHM). The intervals of the red/blue lobes were defined by how far is it from the center velocity and how strong the intensity is.

Because the emission lines of ^{12}CO are optically thicker than other lines, it is appropriate to trace the outflows with ^{12}CO lines. I drew contour maps to find out if bipolar outflows existed with the protostar at its center. To check if the red blue lobes that are found are outflows from the same protostar I checked the ^{12}CO , ^{13}CO , C^{18}O lines from the red peak, blue peak, and the center points. For each outflows confirmed, I calculated the column density and the momentum force.

III. Results

III.1 Outflow Identification

Table 1. Protostars with observed outflows.

Name	coordinates		L_{bol}	T_{bol}
	RA	Dec	L_{\odot}	K
FIR2	05:35:24.3	-05:08:33.3	5.68	100.6
FIR3	05:35:27.5	-05:09:32.5	360.86	71.5
FIR6b	05:35:23.4	-05:12:03.2	21.93	54.1
MMS2	05:35:18.3	-05:00:34.8	20.11	186.3
MMS5	05:35:22.4	-05:01:14.1	15.81	42.4
MMS9	05:35:26.0	-05:05:42.4	8.91	38.1

Table 1 shows the protostars with bipolar outflows observed.

III.1.1 ^{12}CO J = 2 - 1 Observations




Figure 2. The contour map(left) and the line profile(right) of FIR2.




Figure 3. The contour map and the line profile of FIR3.




Figure 4. The contour map and the line profile of FIR6b.




Figure 5. The contour map and the line profile of MMS2.




Figure 6. The contour map and the line profile of MMS5.




Figure 7. The contour map and the line profile of MMS9.

FIR2 - There is a strong bipolar outflow elongated along the N-S direction as shown in Figure 2. The size is about 30 arcsec, which is smaller than other outflows detected. Red and blue contour intervals are 10σ starting from 60σ , and 10σ starting from 100σ , respectively.

FIR3 - A strong bipolar outflow can be seen along NE-SW direction, with red and blue lobes overlapped with each other as shown in Figure ???. This tells us that the outflow axis is almost parallel to the line of sight. Red and blue contour intervals are 20σ starting from 40σ , and 20σ starting from 60σ , respectively.

FIR6b - The contour is not so clear because of other IR sources nearby as shown in Figure ???. The outflow is along the NW-SE direction. Red and blue contour intervals are 10σ starting from 45σ , and 10σ starting from 110σ , respectively.

MMS2 - The contour is in a tricky situation, because both red and blue lobes are in the east side of the protostar as shown in Figure ???. The outflow structure on the SW side is the outflow from another protostar, MMS5. It is possible that the outflow structure changed shape because of the turbulence from other protostars. Red and blue contour intervals are 10σ starting from

30σ , and 10σ starting from 60σ , respectively.

MMS5 - There is an outflow structure along the E-W direction as shown in Figure ?? . This outflow is much smaller than other bipolar outflows. Red and blue contour intervals are 10σ starting from 20σ , and 10σ starting from 40σ , respectively.

MMS9 = There is a strong outflow along the E-W direction as shown in Figure ?? . We can see a smaller red lobe near the center of the blue lobe. Red and blue contour intervals are 10σ starting from 50σ , and 10σ starting from 60σ , respectively.

III.1.2 $^{12}\text{CO J} = 1 - 0$ Observations




Figure 8. The contour map of FIR2(left), FIR3(middle), and MMS9(right).

FIR2 - The red lobe is clear on the NW side of the protostar, but the blue lobe is not that clear as shown in Figure ??.

FIR3 - The outflows are in a similar shape with the $J = 2 - 1$ observations as shown in Figure ?? . The lobe centers are slightly near the protostar.

MMS9 - The outflows are also in a similar shape with the $J = 2 - 1$ observations as shown in Figure ?? . We can also see that there is a small red lobe near the center of the blue lobe.

III.2 Momentum Flux

Table 2. CO outflow parameters.

Name	J = 2 - 1			J = 1 - 0		
	F_R	F_B	F_{CO}	F_R	F_B	F_{CO}
			$(M_\odot \text{ km s}^{-1} \text{ yr}^{-1})$			
FIR2	1.14E-05	3.28E-05	4.42E-05	4.78E-06	-	4.78E-06
FIR3	4.77E-04	7.43E-04	1.22E-03	1.86E-04	3.02E-04	4.88E-04
FIR6b	1.13E-05	1.18E-05	2.31E-05	-	-	-
MMS2	1.14E-05	4.50E-05	5.64E-05	-	-	-
MMS5	5.80E-06	1.55E-05	2.13E-05	-	-	-
MMS9	3.67E-06	1.09E-05	1.46E-05	1.45E-06	6.02E-06	7.47E-06

Table 2 shows the parameters of the outflows detected. F_R and F_B stands for the outflow forces for the red lobe and the blue lobe respectively. F_{CO} is calculated by adding the two forces, which shows the momentum flux of the protostar. We can see that more outflows were detected by using J = 2 - 1 data, and the momentum flux is 2-3 times higher.

III.3 Momentum flux vs. Bolometric luminosity




Figure 9. Momentum flux difference by emission line energy.

Figure 9 shows the relation between the bolometric luminosity and the momentum flux of the outflows from previous studies [8–13].

Since the momentum flux of the same protostar is known to vary somewhat depending on the calculation methods [9], the relation between the bolometric luminosity and the momentum flux is difficult to express with the exact formula and only the degree of tendency can be analyzed. The bolometric luminosity was observed by the Spitzer and Herschel telescopes. Orion A Cloud is a region where stars with medium mass are formed. The fact that the momentum flux of the outflow is proportional to the bolometric luminosity could be checked.

III.4 Momentum flux by emission line energy level




Figure 10. Momentum flux difference by emission line energy.

Figure ?? compares momentum flux calculated by three different emission lines of the same protostar. The ^{12}CO $J = 3 - 2$ observation was made by Takahashi et al [8]. We can see that it is possible to detect more outflows by using a higher energy emission line of ^{12}CO . Using data with smaller beamwidth also enhances detecting outflows. The reason that higher energy lines can detect more outflows can be explained as the following. The excitation temperature is higher for emission lines with higher energy. Outflows drag out matter from the protostar's envelope, which has higher temperature than its surroundings. Lines with higher energy are

emitted, which has an effect that makes column density higher than usual.

IV. Conclusion

The main results of this study are the followings:

1. 6 bipolar outflows were detected from the Orion A Cloud. All outflows were detected by $J = 2 - 1$ data, and 3 outflows were detected by $J = 1 - 0$ data.
2. The well-known correlation between momentum flux and the bolometric luminosity can be checked.
3. It is possible to detect more outflows by using a higher energy emission line of ^{12}CO .
Using data with smaller beamwidth also enhances detecting outflows. The reason that higher energy lines can detect more outflows can be explained as the following. The excitation temperature is higher for emission lines with higher energy. Outflows drag out matter from the protostar's envelope, which has higher temperature than its surroundings. Lines with higher energy are emitted, which has an effect that makes column density higher than usual.

References

- [1] Bontemps, S., André, P., Terebey, S., & Cabrit, S. (1996). Evolution of outflow activity around low mass embedded young stellar objects. In *Disks and Outflows Around Young Stars* (pp. 270–275). Springer.
- [2] Kang, S., Lee, J.-E., Choi, M., & Evans, N. (2013). Outflow properties of digit embedded sources. *한국천문학회보*, 38(1), 51–51.
- [3] Megeath, S., Gutermuth, R., Muzerolle, J., Kryukova, E., Flaherty, K., Hora, J., . . . et al. (2012). The spitzer space telescope survey of the orion a and b molecular clouds. i. a census of dusty young stellar objects and a study of their mid-infrared variability. *The Astronomical Journal*, 144(6), 192.
- [4] Furlan, E., Fischer, W., Ali, B., Stutz, A., Stanke, T., Tobin, J., . . . et al. (2016). The herschel orion protostar survey: spectral energy distributions and fits using a grid of protostellar models. *The Astrophysical Journal Supplement Series*, 224(1), 5.
- [5] Kounkel, M., Hartmann, L., Loinard, L., Ortiz-León, G. N., Mioduszewski, A. J., Rodríguez, L. F., . . . et al. (2017). The gould’s belt distances survey (gobelins). ii. distances and structure toward the orion molecular clouds. *The Astrophysical Journal*, 834(2), 142.
- [6] Schulz, N. S. (2012). *The formation and early evolution of stars: from dust to stars and planets*. Springer Science & Business Media.
- [7] Berne, O., Marcelino, N., & Cernicharo, J. (2014). Iram 30 m large scale survey of 12co (2-1) and 13co (2-1) emission in the orion molecular cloud. *The Astrophysical Journal*, 795(1), 13.

- [8] Takahashi, S., Saito, M., Ohashi, N., Kusakabe, N., Takakuwa, S., Shimajiri, Y., . . . & Kawabe, R. (2008). Millimeter-and submillimeter-wave observations of the omc-2/3 region. iii. an extensive survey for molecular outflows. *The Astrophysical Journal*, 688(1), 344.
- [9] van der Marel, N., Kristensen, L. E., Visser, R., Mottram, J. C., Yıldız, U. A., & van Dishoeck, E. F. (2013). Outflow forces of low-mass embedded objects in ophiuchus: a quantitative comparison of analysis methods. *Astronomy & Astrophysics*, 556, A76.
- [10] Hogerheijde, M. R., van Dishoeck, E. F., Blake, G. A., & van Langevelde, H. J. (1998). Envelope structure on 700 au scales and the molecular outflows of low-mass young stellar objects. *The Astrophysical Journal*, 502(1), 315.
- [11] Nakamura, F., Miura, T., Kitamura, Y., Shimajiri, Y., Kawabe, R., Akashi, T., . . . et al. (2012). Evidence for cloud-cloud collision and parsec-scale stellar feedback within the 11641-n region. *The Astrophysical Journal*, 746(1), 25.
- [12] Aso, Y., Tatematsu, K., Sekimoto, Y., Nakano, T., Umemoto, T., Koyama, K., & Yamamoto, S. (2000). Dense cores and molecular outflows in the omc-2/3 region. *The Astrophysical Journal Supplement Series*, 131(2), 465.
- [13] Zhang, Q., Hunter, T., Brand, J., Sridharan, T., Cesaroni, R., Molinari, S., . . . & Kramer, M. (2005). Search for co outflows toward a sample of 69 high-mass protostellar candidates. ii. outflow properties. *The Astrophysical Journal*, 625(2), 864.

감사의 글

정말 감사합니다.

연구 활동

- 2011학년도 교내 R&E 발표대회에서 장려상 수상
- 2012학년도 교내 R&E 발표대회에서 장려상 수상
- 2013학년도 교내 R&E 발표대회에서 장려상 수상
- 2014학년도 교내 R&E 발표대회에서 장려상 수상
- 2015학년도 교내 R&E 발표대회에서 장려상 수상
- 2016학년도 교내 R&E 발표대회에서 장려상 수상
- 2017학년도 교내 R&E 발표대회에서 장려상 수상
- 2018학년도 교내 R&E 발표대회에서 장려상 수상
- 2019년 노벨 물리학상 수상