

SEOSAW dataset manual

v3.0

John L. Godlee

15th October 2022

Contents

1	Introduction	2
2	SEOSAW code repository	2
2.1	Building the SEOSAW R package for testing	3
3	Description of the SEOSAW dataset	3
3.1	Descriptive statistics and dataset visualisation	4
3.2	Plot data	11
3.2.1	Nesting and subplots	11
3.2.2	Stems or trees	13
3.2.3	Permanent or one-off plot	13
3.3	Subdivided large plots	14
3.3.1	Sampling metadata	14
3.3.2	Landscape metadata	15
3.3.3	Land use metadata	15
3.3.4	Natural disaster metadata	16
3.3.5	Fire and herbivory metadata	16
3.3.6	Floristic clusters	17
3.4	Stem data	17
3.4.1	Taxonomy	17
3.4.2	Biomass	18
3.4.3	Diversity	19
3.5	Spatial data	19
3.5.1	Plot polygons	19

3.5.2	Spatial data extraction	19
3.6	Dataset caveats	21
4	Contributing data	22
4.1	Submission guidelines	22
4.2	Vital data	22
4.3	Additional data	23
5	Cleaning data for the SEOSAW dataset	23
5.1	Cleaning plot data	26
5.2	Plot polygons	27
5.3	Cleaning stem data	27
5.3.1	Multi census data	29
5.3.2	Fix species names	29
5.3.3	Stem biomass	30
5.3.4	Check logical consistency of plots and stems tables	31
5.3.5	Generate plot level statistics from stem data	32
5.4	Compiling datasets	33
5.5	Extracting spatial data	34
6	Dataset table columns	35

1 Introduction

This manual focuses on the **SEOSAW** dataset and provides information on three main topics:

1. The contents of the SEOSAW dataset
2. Contributing data to SEOSAW
3. How data is cleaned for the SEOSAW dataset with the `{seosawr}` R package

For further guidance on collecting SEOSAW data in the field, you should consult the **SEOSAW Field Manual**.

2 SEOSAW code repository

The **SEOSAW code repository** is currently hosted on BitBucket. It contains the `{seosawr}` R package, reproducible code used to clean data from each site, and code for further SEOSAW analyses. Below is a breakdown of

directories in the repository and what they contain:

- `cleaning` - R scripts which clean data for the SEOSAW dataset
- `seosawr` - the SEOSAW R package, used to clean the SEOSAW data and provides convenience functions for further analysis
- `doc` - documents, manuals, and items for the [SEOSAW website](#)

Note that the SEOSAW dataset lives in a separate repository with limited private access, due to the sensitivity of some datasets.

To access the SEOSAW code repository, use `git clone https://bitbucket.org/miombo/seosaw/`, or download the repository directly from [BitBucket](#) as a `.zip` file.

2.1 Building the SEOSAW R package for testing

It is recommended to use the `{devtools}` package to develop the `{seosawr}` R package. To install the package first download the Bitbucket repository using the guidance above.

Open R and set the working directory to `seosaw/seosawr/seosawr/`. Then run the following to install the package on your system:

```
devtools::install()
```

For testing, use the following to load all the package functions into the current R session, including non-exported functions:

```
devtools::load_all()
```

To rebuild the package documentation, run:

```
devtools::document()  
devtools::build_manual()
```

To check the package for common errors, run:

```
devtools::check()
```

Consult the [R Packages book](#) by Hadley Wickham and Jenny Bryan for more guidance on developing R packages.

3 Description of the SEOSAW dataset

This section gives a brief overview of the contents of the latest iteration of the SEOSAW dataset, which is available for research purposes upon approval from the SEOSAW network. At the time of writing the latest version is `v3.0`.

The SEOSAW dataset is built around two key tables:

- `plots.csv` - Plot level data

- `stems.csv` - Stem level data

Each row in the plots table represents a single census in a single plot. Each row in the stems table represents a single stem measurement. Note that a single tree may contain many stems and each stem may be measured multiple times. See [section 6](#) for a breakdown of columns found in the stems table and plots table. All data objects are linked by the `plot_id` column, which serves as the primary key for the dataset.

There are also some auxiliary data objects in the SEOSAW data repository:

- `polys.shp` - Polygons for each plot
- `small_stems.csv` - Count data for small stems (below minimum diameter threshold)
- `cwd.csv` - Coarse Woody Debris (CWD) data
- `herbaceous.csv` - Herbaceous biomass data

There are also some derived datasets available:

- `stems_latest.csv` - Stem data subset for the most recent census of each plot
- `stems_multi.csv` - Stem data subset for plots with multiple censuses
- `plots_latest.csv` - Plot data subset for the most recent census of each plot
- `plots_multi.csv` - Plot data subset for plots with multiple censuses
- `plots_spatial.csv` - Data extracted from various spatial data layers for each plot
- `plots_subdiv.csv` - Plot data for 1 ha subdivisions of plots >2 ha
- `polys_subdiv.shp` - Polygons for 1 ha subdivisions of plots >2 ha
- `plot_clusters.csv` - Floristic clusters for a subset of the plot data
- `cluster_dom.csv` - Dominant species per cluster

3.1 Descriptive statistics and dataset visualisation

- Number of plots: 9992
- Number of permanent plots: 290
- Number of one-off plots: 9702
- Number of stem measurements: 583871
- Number of species: 1249
- Total plot area surveyed (km²): 14.54




Figure 1: Distribution of plots in southern Africa.

Table 1: Country level statistics. Note that none of these statistics are representative of that country, they are only representative of the plots within those countries, which are themselves not evenly distributed.

Country	Mean AGB ($tAGB\ ha^{-1}$)	Mean BA ($m^2\ ha^{-1}$)	N plots
Angola	54.125	10.2768	340
Botswana	0.000	0.0000	40
Congo - Brazzaville	5.811	0.8079	6
Congo - Kinshasa	110.582	18.1995	15
Kenya	15.618	7.3424	11
Malawi	53.490	7.9693	181
Mozambique	58.128	9.3885	4050
Namibia	116.522	26.0773	213
South Africa	86.334	13.9606	441
Tanzania	70.482	11.1304	188
Uganda	567.148	54.2879	54
Zambia	42.375	6.9378	4101
Zimbabwe	33.360	7.4000	352

Table 2: Dataset level statistics. AGB = Above-Ground Biomass, BA = Basal Area. Note that ‘N census’ refers to the total number of censuses conducted within a dataset, the number and dates of censuses per plot may vary. ‘(P)’ next to ‘N plots’ indicates that plots in this dataset are permanent.

Dataset ID	Country	N plots	Min. DBH threshold (cm)	Plot area (ha)	Census range	N census	Mean AGB ($tAGB\ ha^{-1}$)	Mean BA ($m^2\ ha^{-1}$)
ABG	Angola	19 (P)	5	1	2018-2022	3	40.6	7.4
ACM	Angola	60	1	0.1	2013	1	83.8	12.4
AUR	Angola	4 (P)	5	0.25	2022	1	13.0	2.6
CBN	Congo - Brazzaville	6 (P)	5, 11	25	2015-2021	2	5.8	0.8
DKS	Congo - Kinshasa	3 (P)	10	1-10	2012-2014	2	105.9	15.7
DLP	Congo - Kinshasa	12	5	0.03	2017	1	111.8	18.8
KKT	Kenya	3 (P)	5	0.5	2021	1	8.3	2.3
KLA	Kenya	2		1	2021	1	18.7	5.9
KLS	Kenya	6 (P)	0	0.25	2016	1	18.2	10.3
MAR	Mozambique	404	5, 10	0.03-0.2	2014-2015	2	29.1	6.3
MCF	Malawi	93	20	0.05-0.13	2011	1	31.1	5.2
MCL	Malawi	38 (P)	5	0.25	1998-2004	4	55.3	10.2
MGR	Mozambique	15 (P)	5	1	2006-2021	5	53.9	9.0
MLC	Mozambique	51	5	0.03	2011	1	124.3	19.0
MMB	Mozambique	10	5	1	2019	1	115.2	15.0
MMB	Mozambique	10 (P)	5	1	2019	1	115.2	15.0
MNF	Malawi	50	20	0.1-0.13	2011	1	93.8	11.4
MNI	Mozambique	3520	10	0.1	2018	1	60.9	9.7
MNR	Mozambique	50 (P)	5	0.28	2005-2015	2	18.9	2.9
NCC	Namibia, Angola	217	5, 20, 45	0.03-0.07	2011-2014	4	136.3	30.0
SBM	South Africa	5 (P)	5	1	2020	1	2.8	1.3
SBT	South Africa	55	4	0.02	2017	1	19.1	14.6
SHD	South Africa	175	6	0.25	2007	1	60.1	10.5
SHS	South Africa	202	10	0.04	2007	1	131.2	17.4
SSM	South Africa	1 (P)	0	4	2018	1	4.1	1.0
SST	South Africa	3 (P)	5	5	2018	1	1.0	0.2
TIM	Tanzania	45 (P)	5	0.04	2012	1	76.9	13.5
TKM	Tanzania	12 (P)	5	0.09	2020	1	60.9	10.6
TKN	Tanzania	56	5	0.2	2011-2012	2	62.4	10.0
TKW	Tanzania	25 (P)	5	1	2012-2021	4	51.9	7.4
TKX	Tanzania	14 (P)	40	9	2019	1	21.9	2.2
TLA	Tanzania	36	5	0.1-1.9	2016	1	110.1	16.2
UEK	Uganda	54	5, 10	0.5-1	2019	1	567.1	54.3
VAS	Angola, Namibia, Botswana	293	5, 9, 10, 19	0.1	2012-2017	6	26.6	5.6
ZCC	Zambia	4 (P)	1, 3	0.19-1	2001-2021	3	31.7	5.5
ZGF	Zimbabwe	66	5	0.07	2015	1	60.5	11.4
ZHF	Zimbabwe	7	5	0.07	2014	1	40.7	9.6
ZHH	Zimbabwe	18 (P)	5	0.1	2002	1	19.5	13.8
ZIS	Zambia	3886	10	0.1	2014	1	40.0	6.7
ZKS	Zambia	33	5	0.02-0.13	2006	1	28.9	7.3
ZLT	Zambia	178	10	0.07	2015	1	97.0	12.2
ZNF	Zimbabwe	135	10	0.1	2017	1	22.2	4.4
ZPF	Zimbabwe	11 (P)	5	0.25	2006-2021	3	71.3	14.5
ZRF	Zimbabwe	35	5	0.04	2014	1	49.6	10.7
ZWP	Zimbabwe	80	3	0.13	2015	1	19.9	5.1




Figure 2: Diameter-height distribution of all stem measurements. The blue line shows a linear regression and the green line shows a loess line of best fit.




Figure 3: Distribution of plots according to Mean Annual Temperature (MAT) and Mean Annual Precipitation (MAP) of the SEOSAW region. MAT and MAP extracted from the WorldClim dataset gridded at 30 arc second resolution, between 1970 and 2000 (Fick and Hijmans, 2017). Permanent plots are coloured red, non-permanent plots are blue.




Figure 4: Distribution of plots according to herbivore biomass and fire frequency within the SEOSAW region. Herbivore biomass is derived from Hempson, Archibald and Bond (2017) and estimates the biomass of all herbivores. Fire count is derived from the MODIS Burned Area Product (MCD64A1) between 2001 and 2018 (L. Giglio et al., 2015). Both products are gridded at 0.05 degree resolution. Permanent plots are coloured red, non-permanent plots are blue.




Figure 5: Distribution of plots according to human population density and travel time to cities within the SEOSAW region. Human population density data is taken from the WorldPop dataset (1 km) (Linard et al., 2012). Travel time data is estimated from the Malaria Atlas Project 2015 travel time to cities product (Meijer et al., 2018) (30sec). Note that both axes are log-transformed. Permanent plots are coloured red, non-permanent plots are blue.

3.2 Plot data

This section describes important aspects of the plot data in `plots.csv`. See section 6 for a full breakdown of columns in the plots table.

3.2.1 Nesting and subplots

It is common for researchers to stratify their sampling effort across a plot, often according to stem diameter classes, to reduce the work required to measure a plot. This stratification is often referred to as ‘nesting’. See the diagram below for an example of a nested plot. In this example, there are three levels of nesting. Across the whole plot (0.2 ha) stems >10 cm DBH are measured. Within the first level of nesting (N1, 0.1 ha), stems >5 cm DBH are measured, within the second level (N2, 0.05 ha) stems >2 cm DBH, and within the third level (N3, 0.025 ha) stems >1 cm DBH.




Figure 6: Schematic diagram of a nested plot with three levels of nesting.

There are a number of columns in `plots.csv` which record nesting. The columns listed below are only those to describe the first level of nesting, but there are similar columns for the second (`n2_`) and third (`n3_`) levels of nesting:

- `nested_1` - is the plot nested at the first level?
- `n1_min_diam_thresh` - minimum stem diameter threshold within level of nesting
- `n1_min_height_thresh` - minimum stem height threshold within level of nesting
- `n1_plot_shape` - shape of nested plot
- `n1_plot_width` - width of nested plot
- `n1_plot_length` - length of nested plot
- `n1_plot_radius` - radius of nested plot

If plots are nested, with variation in sampling effort for trees of different sizes, the Fractional Probability of Inclusion in the `fpc` column of the stem table describes the sampling effort for each stem. In the example, stems >10 cm DBH are measured across the whole plot, so their `fpc` is 1.0. Stems >5 cm DBH are measured in an area 20x25 m (0.1 ha), which is half the total plot area, so their `fpc` is 0.5. Stems >2cm are measured over 0.05 ha, which is a quarter of the total plot area, so their `fpc` is 0.25, and so on. At present, the SEOSAW dataset only allows up to three levels of nesting, but this may change in the future.

It is important to distinguish between nesting and subplots. Nested plots, as demonstrated above, are smaller parcels within a plot which denote variation in sampling effort. In contrast, subplots are merely administrative sub-divisions of a plot which can help with relocating stems for future censuses, for example. Sampling effort should be the same across all subplots. Below is an example of a plot with four 20x25 m subplots.




Figure 7: Schematic diagram of a plot with subplots, each occupying 25% of the total plot space.

There are a few columns in `plots.csv` which record subplot dimensions. It is assumed that all subplots have the same dimensions and that subplots are not nested:

- `subplot_n` - number of subplots in the plot
- `subplot_shape` - shape of subplots
- `subplot_length` - length of subplots
- `subplot_width` - width of subplots
- `subplot_radius` - radius of subplots

3.2.2 Stems or trees

The ideal data for the SEOSAW dataset is organised so that there is one measurement for each stem, and stems are grouped into trees with a permanent tag to identify each stem. We understand however that not all data matches this ideal, and is dependent on the original data collection protocol. There are a few columns in the plots table to help describe how stem data is organised:

- `tags` - Do stems have permanent tags which allow for remeasurement?
- `tag_tree_stem` - Are tags applied to each tree, or each stem?
- `meas_tree_stem` - Are measurements recorded at the tree or stem level?
- `tree_diff` - If `meas_tree_stem == 'stem'`, were stem measurements grouped into separate trees?

3.2.3 Permanent or one-off plot

A permanent plot in the SEOSAW dataset is defined as one where trees and new recruits can be measured at a later date. The boundaries of a permanent plot must be clearly identifiable, i.e. with corner posts for a rectangular plot or a central marker for a circular plot. Individual trees must also be identifiable in a permanent plot. Normally this is achieved through the use of numbered tags, but can alternatively be achieved with a highly accurate grid coordinate system, a differential GPS, or other tagging methods.

Related to this, often it is possible to track the growth and mortality of individual stems between multiple censuses within a plot. The `stem_timeline_cont` column records whether the current census can be successfully stem-matched with the previous census. The `abandoned` column records if a plot has been abandoned, i.e. will not receive any more censuses in the future.

One-off plots are those where only a single census has been made, with no way of exactly identifying plot boundaries or tree identity in the future.

Both permanent and one-off plots are accepted by SEOSAW. The `permanent` column in the plots table notes whether a plot is permanent or not.

3.3 Subdivided large plots

`plots_subdiv.csv` and `polys_subdiv.shp` hold data on 1 ha square subdivisions of large plots. Large plots are subdivided to make them more readily comparable with other 1 ha plots in the dataset, and to make them easier to work with when conducting spatial analyses. Plots are only subdivided if a number of conditions are met:

- Plot is ≥ 2 ha
- Plot is rectangular
- All plot edges are divisible by 100 m

Additionally, stems are only assigned to a subdivision if they have grid coordinates in `x_grid` and `y_grid`. If your stems are located instead by latitude/longitude coordinates, consider using the `gridCoordGen()` function to approximate grid coordinates originating from the southwest corner of the plot. Subdividing plots is performed by `largePlotSplit()` in the `{seosawr}` R package.

The `subdiv` column in the plots table shows whether a plot has been subdivided. The `subdiv_id` column in the stems table shows which subdivision a stem has been assigned to. `subdiv_id` matches in `plots_subdiv.csv` where subdivision-level calculated values (e.g. `agb_ha`) are stored. The calculated values are calculated for each census per plot, like `plots.csv`. If you want only the most recent census for each plot, use `subsetLatest()`. Subdivision IDs are assigned by appending `_S[1-n]` to the plot ID.

3.3.1 Sampling metadata

There is variation in measurement protocol across the SEOSAW dataset. In some plots for example, only living trees were sampled while dead trees were ignored completely. This does not mean there were no dead trees in those plots. Unless these differences in measurement protocol are accounted for in analysis, erroneous relationships may appear. There are a few columns in the plots table which describe variations in measurement protocol:

- `lianas_sampled`
- `soil_sampled`
- `herbaceous_sampled`
- `dead_stems_sampled`
- `stumps_sampled`
- `coarse_woody_debris_sampled`
- `small_stems_sampled`

- `sampling_notes`
- `sampling_design_code` - Reference to document containing the description of protocol for data collection.
- `xy_method`
- `diam_method`
- `height_method`
- `slope_method`
- `slope_method_notes`
- `canopy_cover_method`
- `canopy_cover_method_notes`

3.3.2 Landscape metadata

The situation of a plot within the landscape may affect vegetation dynamics. The following columns in the plots table provide metadata on the general landscape in the plot:

- `longitude_of_centre`
- `latitude_of_centre`
- `elevation`
- `slope`
- `aspect`
- `catenal_position`
- `termites`

3.3.3 Land use metadata

Vegetation within a plot may differ hugely based on any human land use. The following columns in the plots table provide metadata on human land use and may be used to filter data prior to analysis:

- `pa` - protected area
- `pa_name`
- `pa_type`
- `wdpa_id` - WDPA protected area ID
- `local_community_rights`
- `other_user_rights`
- `timber_harvesting`
- `high_graded_100_years`
- `fuel_wood_harvesting`

- charcoal_harvesting
- other_woody_product_harvesting
- ntfp_harvesting
- farmed_30_years
- clear_cut
- mining
- land_use_notes

3.3.4 Natural disaster metadata

It is hard to measure the impact of natural disasters on a plot over the long term due to their stochastic nature. The following columns in the plots table provide metadata on natural disasters:

- cyclone
- cyclone_last_date
- drought
- drought_last_date
- wind
- wind_last_date
- earthquake
- earthquake_last_date
- landslide
- landslide_last_date
- flood
- flood_last_date
- frost
- frost_last_date

3.3.5 Fire and herbivory metadata

Disturbance by fire and herbivory can greatly affect vegetation dynamics. Users of the SEOSAW dataset may want to filter plots based on disturbance regime. These columns in the plots table provide metadata on disturbances:

- fire_exclusion - Are fires excluded by humans?
- fire_exclusion_success
- fire_treatment - Are fires controlled by humans?
- fire_return_interval

- `fire_last_date`
- `fire_treatment_season`
- `fire_notes`
- `elephants`
- `elephant_density`
- `elephant_data_source`
- `exclosure`
- `small_nonsocial_browsers`
- `large_browsers`
- `medium_sized_social_mixed_diet`
- `nonruminants_excluding_suids`
- `water_dependent_grazers`
- `goat_grazing`
- `sheep_grazing`
- `cattle_grazing`
- `herbivory_notes`

3.3.6 Floristic clusters

A portion of the SEOSAW dataset has been clustered based on Ward's hierarchical clustering method. Data is clustered based on stem abundances per genus in the most recent census per plot. Plots with fewer than 14 stems were excluded from the analysis. Plots with exotic species were also excluded. Data on which cluster each plot is assigned to is found in `plot_clusters.csv` with the dominant species per cluster found in `cluster_dom.csv`.

3.4 Stem data

This section describes aspects of the stem data in `stems.csv`. See [section 6](#) for a full breakdown of columns in the stems table.

3.4.1 Taxonomy

There are a number of columns in the stems table which describe the species of a stem. The list below includes a few of the most important columns:

- `species_orig_binom` - binomial species name supplied by data originator
- `species_orig_local` - local (non-binomial) species name supplied by data originator
- `species_name_sanit` - combined binomial species name after manual cleaning in data cleaning script.
- `species_name_clean` - combined binomial species name after cleaning and synonymy checks. This is the species name to use for most analyses.

- `species_name_gnr` - species name matched by `taxize::gnr_resolve()`
- `family_gnr` - family name matched by `taxize::gnr_resolve()`
- `taxon_rank_gnr` - all rank names of classification from `taxize::gnr_resolve()`
- `taxon_gnr` - all ranks of classification from `taxize::gnr_resolve()`
- `species_name_cjb` - binomial species name matched by the CJB African plants database

The taxonomic ‘backbone’ of the SEOSAW dataset is the [CJB African plants database](#) with some notable exceptions, which are discussed below. The dataset also uses the `taxize::gnr_resolve()` utility to query multiple databases to provide higher taxonomic rank information. The queried databases are:

- [Open Tree of Life Reference Taxonomy](#)
- [The International Plant Names Index](#)
- [Tropicos - Missouri Botanical Garden](#)
- [GBIF Backbone Taxonomy](#)
- [NCBI](#)

The breakup of the *Acacia* genus has been well documented by (Kyalangalilwa et al., 2013). SEOSAW uses their list of synonyms to correct ex-*Acacia* species. The CJB database does not currently acknowledge the *Acacia* split, so queries to CJB retain the *Acacia* name.

All *Rhus* species in SEOSAW’s study region belong to *Searsia* and are transferred in this way. All recent phylogenetic results point to this, but the formal taxonomic work to transfer African *Rhus* species is incomplete.

Previously, *Marquesia* species were placed in the *Trillesanthus* genus, however this has since been reverted. SEOSAW corrects *Trillesanthus* to *Marquesia*.

The latest taxonomy states that *Maytenus* is a strictly Neotropical genus, while acknowledging that African *Maytenus* are a “problem”. The latest taxonomic literature doesn’t tell us what to do with African *Maytenus*. SEOSAW defaults to the CJB taxonomy on *Gymnosporia*/*Maytenus*, but this may change in the future.

There are also some miscellaneous synonym corrections which are maintained throughout the canonical SEOSAW dataset. The vast majority of these corrections reflect previously debunked names which still appear in older datasets.

If a species cannot be identified, it is given the “indet” epithet. If only the genus is known, it is recorded as “*Genus* indet” (e.g. *Brachystegia* indet). If only the family is known, it is recorded as “Family indet” (e.g. Fabaceae indet). As a last resort, if only the local name of the species is known, it is recorded in the `species_orig_local` column.

3.4.2 Biomass

Biomass, specifically aboveground woody biomass (AGB) is stored in the `agb` column in the stems table. Additionally, plot level biomass is stored in the `agb_ha` (AGB per hectare) column in the plots table. It is calculated in the SEOSAW dataset by default using an allometric equation devised by (Chave, Réjou-Méchain et al., 2014) which takes into account species wood density, stem diameter and latitude/longitude. This means that plots with no recorded latitude/longitude will have an AGB of NA.

3.4.3 Diversity

The `{seosawr}` package provides slim wrappers around existing functions in the `{vegan}` package to calculate biodiversity metrics at the plot level:

- `richness` - Species richness
- `rarefy` - Rarefied species richness - `vegan::rarefy()`
- `shannon` - Shannon-Wiener index - $H' = -\sum_{i=1}^s p_i \ln p_i$
- `simpson` - Simpson's index - $D = \sum_{i=1}^s p_i^2$
- `invsimpson` - Inverse Simpson's index - ${}^2D = 1/D$
- `evenness` - Simpson's evenness - $(1/D)/N$

Additionally, each plot's five most dominant species, estimated simply by tree abundance, are shown in `dom_1` etc., with their proportional abundances in `prop_1` etc. The values returned are the proportional abundance of the corresponding species, within the plot.

Note that rarefied species richness cannot be calculated for plots with fewer than 25 stems, and will be `NA` for these plots.

3.5 Spatial data

3.5.1 Plot polygons

Plot outline polygons are provided in `polys.shp`, for each plot where appropriate location and plot dimensions data is available. For rectangular plots, these polygons are normally created from GPS coordinates taken at each corner and used as vertices when transforming to a closed spatial polygon. Plot point information is stored in `points.shp`, and uses the plot centre location. Plot polygons are automatically generated using the `polyEstGen()` function in the `{seosawr}` R package, which uses the plot centre coordinates, plot shape, and plot dimensions to create an approximate polygon if none are provided. If plot corner coordinates are provided, the `polyGen()` function can be used to construct a polygon as the convex hull of the points.

3.5.2 Spatial data extraction

`plots_spatial.csv` provides data extracted from a number of spatial data layers for each plot. See [section 6](#) for a breakdown of these spatial data layers and how values are extracted. Note that values are only point estimates from the raster pixel intersected by the plot centre coordinates, they do not average over plot extent polygons.

The SEOSAW region is provided as a spatial data object in the `{seosawr}` R package (`data(seosaw_region)`), along with a simplified bounding box of the SEOSAW region (`data(seosaw_bbox)`). There is also a useful outline of all countries in Africa (`data(africa)`).




Figure 8: The spatial extent of the SEOSAW region shown in green and the SEOSAW bounding box as a blue rectangle, within Africa.

The SEOSAW region is based largely on White's vegetation map, which defines the locations of savannas and woodlands (White, 1983).

A key issue was defining the north-eastern boundary of the SEOSAW region. It is defined here as the borders of Kenya and Uganda, excluding Ethiopia, Somalia, and South Sudan. While the decision to use country boundaries was largely pragmatic, it is backed up by environmental data. The mountainous Ethiopian highlands on the border between Kenya, Ethiopia and Somalia cause quite a clear change in precipitation, probably due to them blocking dry Arabian air moving further southwest.

The north-western boundary largely follows the rift valley lakes, with everything to the west of them being excluded. Some areas to the east of the lakes are also excluded as they are classed by White's vegetation map as

Congo Forest Transitions and Mosaics.

The south-east of the region is further reduced from White's vegetation map according to the Terrestrial Ecoregions of the World (Dinerstein et al., 2017). White classes some areas as woodland or forest in this area, but for our purposes they are considered too temperate, and are thus excluded from the SEOSAW region.

3.6 Dataset caveats

The SEOSAW dataset is aggregated from many individual datasets contributed by researchers around the world. While effort has been made to ensure that datasets are compatible with each other, particularly with respect to stem data like stem `diam` and `height`, some pieces of metadata should not be compared between datasets. For example, `canopy_cover` is collected with hemispherical photos, different types of densitometer, aerial photos and many other methods. Due to the subjectivity of all these methods, it is not recommended to use this data as a variable in statistical analysis when considering multiple datasets; it should be used only as metadata. Below is a non-exhaustive list of fields we recommend not to compare between sites in analyses:

- `slope` - Variable accuracy and recording techniques
- `aspect` - Variable accuracy and recording techniques
- `catenal_position` - Subject to human bias
- `canopy_cover` - Variable accuracy and recording techniques
- `local_community_rights` - Subject to human bias, sparse data
- `other_user_rights` - Subject to human bias, sparse data
- `timber_harvesting` - Subject to human bias, sparse data
- `high_graded_100_years` - Subject to human bias, sparse data
- `fuel_wood_harvesting` - Subject to human bias, sparse data
- `charcoal_harvesting` - Subject to human bias, sparse data
- `other_woody_product_harvesting` - Subject to human bias, sparse data
- `ntfp_harvesting` - Subject to human bias, sparse data
- `farmed_30_years` - Subject to human bias, sparse data
- `fire_exclusion_success` - Subject to human bias
- `elephant_density` - Variable accuracy

Additionally, SEOSAW takes the stance that metadata should be left blank if it is not known, rather than guessed at. This means that many metadata columns consist mainly of `NA` values. For example, `charcoal_harvesting` is left mostly blank because most data originators did not record whether wood collection for charcoal harvesting had occurred in the plot. This also applies to stem data. If for example, the data originator did not record whether stems were located on termite mounds, `on_termites` is left as `NA`, even if it is more likely that most stems were not located on termite mounds.

4 Contributing data

If you would like to contribute data to the SEOSAW network, it is recommended that you format your data in two tables, one for plots and the other for stems, so it is compatible with the existing SEOSAW dataset. The guidance below details the contents of these two tables, and other metadata.

We provide blank template field data sheets in the SEOSAW Bitbucket repository which you can use when setting up a new plot or re-censusing an existing plot ([blank_plot.pdf](#), [blank_stem.pdf](#)). Additionally, we have definitions for columns in the plots and stems tables ([plot_columns.csv](#), [stem_columns.csv](#)). Finally, we can provide [ODK forms](#) to allow for data collection on an Android device, with helpful hints on data collection protocol. See the [SEOSAW website](#) for more information.

4.1 Submission guidelines

Please submit tabular data as [.csv](#) (comma-separated values) files if possible. If working in Microsoft Excel or a similar spreadsheet program, there is usually an option to export as [.csv](#). [.csv](#) files are preferred as they are simple and store data exactly as entered. [.xls](#) or [.xlsx](#) files are also acceptable.

Spatial data such as plot outlines are preferred as shapefiles ([.shp](#)), though plot corner or plot centre locations as latitude-longitude coordinates, stored in a [.csv](#) file are also acceptable. Make sure to describe the Coordinate Reference System (CRS) used for coordinates, WGS84 Latitude-Longitude (EPSG: 4326) is preferred.

Please provide a description of the contents of each file in a separate document. For tabular data, if not using the SEOSAW column definitions, provide a description of the contents of each column. For example, to describe the contents of a column in the stems table containing stem diameter measurements:

[S diam](#) - Stem diameters, measured in centimetres to one decimal place (i.e. to the nearest mm).

Provide a description of the methodology used to collect the data in a separate document, especially any plot nested-ness or stratified sampling, as this will determine whether any corrections should be made when calculating plot-level statistics like aboveground biomass.

4.2 Vital data

There are many columns which can be filled by data contributors in the stem and plot tables, and you should try to provide data for as many as possible. Understandably however, some data simply won't exist depending on the original goal of your data collection. Our stance is that metadata should be left blank if it is not known, rather than guessed at.

Below is a list of the columns which are absolutely necessary for minimum data quality. Data submitted without all of these columns cannot be added to the SEOSAW dataset. The column names as they appear in the template data tables are shown in brackets:

Plots table:

- Plot name, for matching with plot names in stems table ([plot_name](#))
- Plot longitude and latitude coordinates ([longitude_of_centre](#), [latitude_of_centre](#))
- Plot shape and dimensions ([plot_shape](#), [plot_length](#), [plot_width](#), or [plot_radius](#))
- The Primary Investigator of the data ([prinv](#))

- Information on plot nested-ness, including nested plot dimensions and stem size thresholds (`nested_1`, `n1_*`, etc.).

Stems table:

- Plot name, for matching with plot names in plots table (`plot_name`)
- Tree species name (`species_orig_binom` or `species_orig_local`)
- Tree diameter (`diam`)
- Height of diameter measurement (`pom`)
- Fractional Probability of Inclusion. If plots are nested with varying sampling effort within plots for stems of different size, the probability of a stem this size occurring in the whole plot (`fpc`)
- Is the stem alive, dead or resprouting? (`stem_status`)

If only a few entries from each of these columns is missing due to mistakes during fieldwork, don't worry, you data is probably still usable, but try to fill as many empty values as possible.

4.3 Additional data

Please provide *any* metadata you think might be even slightly useful to gain a more detailed understanding of your plots. For example, if you have meteorological data for a nearby weather station, herbivore dung counts, photos of the plots, land use history timelines, soil core profiles, or fire management notes. As with the plot and stem data, try to provide a detailed description of the data, the methodology used to collect the data, and any caveats.

5 Cleaning data for the SEOSAW dataset

`{seosawr}` is an R package for the SEOSAW dataset. The package serves two main purposes:

1. Formatting data for inclusion in the SEOSAW dataset, or for comparison with the existing SEOSAW dataset.
2. Providing convenience functions for further analysis and manipulation of the SEOSAW dataset.

Most of the functions in `{seosawr}` act on either one of or both the SEOSAW dataset tables, with the column names necessary for that function specified by additional arguments in the function. For example, the function to calculate aboveground biomass using the Chave, Réjou-Méchain et al. (2014) allometry:

```
ex_data$agb <- chave14(x = ex_data$stems, plot_data = ex_data$plots,
  species_name = "species_name_clean",
  diam = "diam",
  stem_plot_id = "plot_id",
  plot_plot_id = "plot_id",
  latitude_of_centre = "latitude_of_centre",
  longitude_of_centre = "longitude_of_centre")
```

The column name arguments default to column names specified by the SEOSAW dataset, but can be altered by the user to allow for some flexibility.

Most top level functions specify a dataframe as the first argument (`x`). Depending on the function, the output is normally either a modified version of `x` with new columns added, or a simple vector with the same length as `nrow(x)` and the same row order as `x`. See the `chave14()` example above for a function which returns a vector, and the species generator function `speciesCheck()` for an example which returns a modified version of the stems table:

```
speciesCheck(ex_data$stems$species_orig_binom,
  ret_unk_seosaw = TRUE, ret_unk_cjb = TRUE, ret_unk_gnr = TRUE)
```

Read the data collection manuals before setting up new plots and collecting data in the field. Follow their protocol to maximise compatibility of newly collected data with existing SEOSAW dataset. The SEOSAW manuals can be found [here](#).

It is suggested that wherever possible you refrain from directly editing raw spreadsheets collected in the field. Instead, use R to create a reproducible script which transforms the raw data into a format which is compatible with the larger SEOSAW dataset. Editing spreadsheets directly is likely to introduce more errors due to mistyping, and your future self will be unsure of the exact data cleaning procedure, which may be important for future analysis and scientific writing.

The following example outlines a generic data cleaning process for a small fictional dataset (`mplots_bad`, `mstems_bad`). Real datasets are likely to be much larger than this, but the cleaning process should be functionally similar. This is the same cleaning process which is followed for all datasets in the SEOSAW dataset. If you want to generate data which behaves exactly like a real SEOSAW dataset, without errors, use the `exampleDataGen()` function in the `{seosawr}` R package.

Table 3: Example plots table before cleaning.

plotcode	plot_cluster	plot_name	dataset_id	census_date	country_iso3	pi	other_authors	plot_shape
MNM_1	MNM_1	a3Uzi	MNM	1979	MOZ	Gyjrs D. O.	Wlzht C.;Boqrx M.	square
MNM_1	MNM_1	a3Uzi	MNM	1982	MOZ	Gyjrs D. O.	Wlzht C.;Boqrx M.	square
MNM_2	MNM_1	AnbJ9	MNM	1979	MOZ	Gyjrs D. O.	Wlzht C.;Boqrx M.	square
MNM_2	MNM_1	AnbJ9	MNM	1982	MOZ	Gyjrs D. O.	Wlzht C.;Boqrx M.	square
MNM_3	MNM_1	8T8Qb	MNM	1979	MOZ	Gyjrs D. O.	Wlzht C.;Boqrx M.	square

Table 4: Example stems table before cleaning.

plotcode	subplot_id	tree	stem	tag_id	census_date	species_orig_binom	species_orig_local	notes_stem
MNM_1	22	MNM_1_2	MNM_1_2_1	1	1979	ZyZy cordat	NA	NA
MNM_1	22	MNM_1_2	MNM_1_2_2	2	1979	Lannear diScolor	NA	NA
MNM_1	25	MNM_1_3	MNM_1_3_3	3	1979	UNKNOWN	NA	NA
MNM_1	21	MNM_1_5	MNM_1_5_4	4	1979	Terminalia prunioides	NA	NA
MNM_1	21	MNM_1_5	MNM_1_5_5	5	1979	Terminalia prunioides	NA	NA

See [Figure 9](#) for a flow diagram of the recommended cleaning process, which outlines the functions used and their order.




Figure 9: Flow diagram of dataset cleaning. “Bascule” refers to the private repository of the SEOSAW dataset. Access granted after consultation with SEOSAW.

5.1 Cleaning plot data

Rename, remove and create columns based on the accepted names found in `plotColClass()`. This function contains a named list, where the names of each list item are the names of columns in the SEOSAW dataset, and the list item is a column constructor function which provides checks and conservatively fixes common errors. While the `{dplyr}` package isn't necessary for this workflow, it excels at dataframe manipulation on medium-sized dataframes and makes some parts of the process much more concise to write and easier to interpret by the reader. The `colGen()` function can be used to generate empty columns to remind you which data fields should be provided:

```
plots_clean <- mplots_bad %>%
  dplyr::rename(
    plot_id = plotcode,
    prinv = pi) %>%
  dplyr::select(
    -soil_carbon) %>%
  colGen(., colClass = plotColClass())
```

Then, replace some values which are incompatible with the SEOSAW dataset's column rules:

```
plots_clean_val <- plots_clean %>%
  mutate(
    plot_shape = case_when(
      plot_shape == "square" ~ "rectangle",
      TRUE ~ plot_shape),
    min_diam_thresh = case_when(
      min_diam_thresh == "<5cm" ~ "5",
      min_diam_thresh == "<10cm" ~ "10",
      TRUE ~ min_diam_thresh))
```

Use `plotTableCheck()` to check whether column names and contents are SEOSAW compatible, then generate new columns with derived values. This checking function can be used iteratively with the cleaning process above until the function returns no errors.

```
plots_check <- plotTableCheck(plots_clean_val)
```

As there are always exceptions to perfect data collection practise, the checking functions return warnings rather than errors by default, indicating which rows or IDs are logically inconsistent. If `output = TRUE` in `plotTableCheck()`, the checking functions return a list of row index vectors, which can then be used to subset the original data to locate potential errors. The user is encouraged to consider the warnings and check whether these columns are legitimately inconsistent, or if there has been an error during data entry.

The default checking functions, and their checks, are shown below:

- `nestedCheck()`
 - If a nesting level is FALSE (e.g. `nested_1 = FALSE`), make sure nesting descriptors (e.g. `n1_min_diam_thresh`) are NAs, and vice versa.
 - If a lower level of nesting is TRUE (e.g. `nested_2 = TRUE`), make sure higher levels are also TRUE (e.g. `nested_1 = TRUE`).
- `paCheck()`
 - If a plot is in a protected area (i.e. `pa = TRUE`), make sure protected area name (`pa_name`) and type (`pa_type`) are filled in.

- `shapeCheck()`
 - If a plot is a circle or rectangle (e.g. `plot_shape = "circle"`), are the corresponding plot dimension columns filled (i.e. `plot_radius`, `plot_length`, `plot_width`).
- `fireCheck()`
 - If a plot has a fire treatment or fire exclusion protocol (e.g. `fire_treatment = TRUE` or `fire_exclusion = TRUE`), are the descriptor fields for those treatments filled (e.g. `fire_treatment_season`, `fire_treatment_success`)?
- `isoCheck()`
 - Does the country described in `country_iso3` agree with the plot coordinates provided in `longitude_of_centre` and `latitude_of_centre`?

The extra columns generated by `plotTableCheck()` are:

- `censusYearGen()` - Generate the year of the census (`census_year`) by extracting from `census_date`
- `plotCensusIDGen()` - Generate the census ID (`census_id`) by concatenating `plot_id` and `census_year`
- `plotAreaGen()` - Generate the area of the plot (`plot_area`) in hectares using `plot_shape`, `plot_radius`, `plot_length` and `plot_width`
- `subplotAreaGen()` - Generate the area of the subplots (`subplot_area`), in hectares, using `subplot_shape`, `subplot_radius`, `subplot_length` and `subplot_width`
- `plotPerimeterGen()` - Generate the perimeter of the plot (`plot_perimeter`) in metres using `plot_shape`, `plot_radius`, `plot_length` and `plot_width`

5.2 Plot polygons

Generate plot polygons either using plot corner coordinates and the `polyGen()` function, or approximate the polygon using the `polyEstGen()` function and the values in `plot_shape`, `plot_width`, `plot_length`, `plot_radius`, `latitude_of_centre`, `longitude_of_centre`. Note that `polyEstGen()` assumes the plot is orientated perfectly north-south, which may not be true:

```
plot_polys <- polyEstGen(plots_check)
```

5.3 Cleaning stem data

At this point, it is recommended to clean the stem data, using a similar procedure to above:

```
stems_clean <- mstems_bad %>%
  dplyr::rename(plot_id = plotcode,
    tree_id = tree,
    stem_id = stem) %>%
  mutate(height = height / 100,
    stem_status = case_when(
      stem_status == "alive" ~ "a",
      stem_status == "dead" ~ "d",
      TRUE ~ stem_status),
    fpc = 1) %>%
  colGen(., stemColClass())
```

Then check that the contents of each column conforms with the guidelines in the `stemTableCheck()` function:

```
stems_check <- stemTableCheck(stems_clean)
```

The default checking functions for `stemTableCheck()`, and their checks, are shown below:

- `diamHeightCheck()`
 - Uses a linear model to flag diameter:height ratios that are clearly unusual, based on a σ standard deviation threshold specified by the user.
- `stemLogicCheck()`
 - If a stem is recorded as broken (i.e. `stem_mode = "P"`), is `broken_height` filled?
 - If a stem is recorded as damaged by ‘other’ (i.e. `damage_cause = "o"`), is `damage_cause_notes` filled?
 - If a stem is recorded as wounded (i.e. `stem_mode = "y"`), is `wound_area` filled?
 - Are both sets of stem coordinates (lat-long, XY, or angle-distance) provided?
 - If a stem is dead (i.e. `stem_status = "d"`), are all other stems on the tree dead?
 - Are all stems on a single tree the same species?
 - If a stem is vanished (i.e. `stem_mode = "v"`), are all other pertinent measures left blank?
- `diamDecCheck()`
 - Identifies diameter values which are unusually common. The function bins diameter measurements and runs a linear regression on the ranked diameter bin frequencies, then checks if any bin frequencies are outside a σ standard deviation threshold.
- `measurementUniqueCheck()`
 - Identifies measurement IDs which are duplicated in the stems table, which are forbidden.
- `stemNCheck()`
 - Checks whether trees have an unusually high number of stems, which might indicate a problem with tree assignment.
- `tagUniqueCheck()`
 - Checks whether tag IDs are duplicated within a plot, which is discouraged

`stemTableCheck()` adds some new columns too:

- `treeIDGen` - New tree ID (`tree_id`) created by concatenating `plot_id` and `tree_id`.
- `stemIDGen` - New stem ID (`stem_id`) created by concatenating `plot_id`, `tree_id` and `stem_id`.
- `censusYearGen()` - Generate the year of the census (`census_year`) by extracting from `census_date`
- `censusIDGen()` - Generate the census ID (`census_id`) by concatenating `plot_id` and `census_year`
- `measurementIDGen()` - Generate the unique measurement ID (`measurement_id`) by concatenating `plot_id`, `census_year`, `tree_id` and `stem_id`.
- `crownArea()` - Crown area (`crown_area`) modelled as an ellipse if both `crown_x` and `crown_y` are provided, otherwise modelled as a circle if only one provided, in m^2 .
- `basalArea()` - Stem basal area (`ba`) in square metres, calculated using DBH measurements (`diam`), assuming a circular trunk cross-section.
- `diamAdj()` - If the DBH was taken at a POM (`pom`) other than 1.3 m, an estimated DBH at 1.3 m (`diam_adj`), assuming a degree of stem taper with height.

5.3.1 Multi census data

If the stem data is from the re-measurement of a previously measured plot, you may want to match up the tree IDs and stem IDs with a previous census, using the tag IDs as a way to match the two datasets. To make this process easier, use `stemIDGet()`:

```
stemIDGet(new_data, old_data,
  plot_id = "plot_id", tag_id = "tag_id",
  plot_id_y = "plot_id", tag_id_y = "tag_id",
  tree_id_y = "tree_id", stem_id_y = "stem_id")
```

You may also want to identify recruits, i.e. new stem IDs, using `recruitGen()`:

```
recruitGen(stems_check, plots_check,
  stem_plot_id = "plot_id", plot_plot_id = "plot_id",
  stem_id = "stem_id", stem_census_date = "census_date",
  plot_census_date = "census_date")
```

Then you can check the multi-census data for logical inconsistencies using `multiCensusCheck()`:

```
multiCensusCheck(stems_clean)
```

- `speciesPlotNewCheck()`
 - Checks if a species is found in previous censuses within each plot.
- `diamChangeCheck()`
 - Identify stems where diameter measurement changes drastically.
- `speciesChangeCheck()`
 - Checks if a stem changes species between censuses
- `resurrectCheck()`
 - Checks if a stem resurrects between censuses. I.e. `stem_status` changes from “d” to “a”

5.3.2 Fix species names

Next fix the species names. `speciesCheck()` generates clean species names by querying `taxize::gnr_resolve()`. Firstly the function checks whether a species has been observed in the larger SEOSAW dataset. Given that the SEOSAW dataset contains in excess of 500,000 stem measurements, if a species is new to the dataset it is likely to be a mistake. Secondly the function corrects common orthographic errors and synonymy issues based on a number of lookup tables, which can be provided by the user or from data objects in the `{seosawr}` package. Thirdly, the function checks `taxize::gnr_resolve()` using the `gnrQuery()` wrapper function to extract taxonomic rank information, and thirdly it checks the CJB database to retrieve the species name. At any of these steps it is possible for the user to specify that the function should return unmatched species names. This is especially useful when cleaning a new dataset where species names may be misspelled or misclassified and need to be manually corrected.

There are a number of lookup tables provided as data objects in the `{seosawr}` package which deal with synonymy issues that aren’t handled appropriately by CJB and/or `taxize::gnr_resolve()`. By default these lookup tables are fed to `speciesCheck()` and species names are automatically corrected. You can change which lookup tables are used by changing the `ortho` parameter in `speciesCheck()`, which takes a list of lookup tables:

- `data(acacia)`
- `data(maytenus)`
- `data(trillesanthus)`
- `data(synonymy)`
- `data(rhus)`

```
stems_species <- speciesCheck(stems_check$species_orig_binom,
  ret_unk_seosaw = TRUE, ret_unk_cjb = TRUE, ret_unk_gnr = TRUE)
```

If species names are not found, `stems_species` will contain a vector of species names not matched. The user can then use this vector to create a lookup table which corrects the species names. If you are sure the species name is correct, just replace the old name with the same name in the lookup table:

```
lookup_table <- data.frame(
  original = stems_species,
  corrected = case_when(
    stems_species == "ZyZy cordat" ~ "Syzygium cordatum",
    stems_species == "Lannear diScolor" ~ "Lannea discolor",
    TRUE ~ stems_species))
```

Then run `speciesCheck()` for a second time, this time passing the lookup table as an argument. This will replace the species name with the corrected version before they are passed to the checking routines.

```
stems_species <- speciesCheck(stems_check$species_orig_binom,
  ret_unk_seosaw = FALSE, ret_unk_cjb = FALSE, ret_unk_gnr = FALSE,
  lookup_user = lookup_table)
stems_species <- cbind(stems_check, stems_species)
```

If `speciesCheck()` is successful, `stems_species` will now be filled with a dataframe containing various taxonomic information.

The function `abMatGen()` can be used to create an abundance matrix from a stem measurements table. An abundance matrix is a table where each column is a species, each row is a plot, and each cell contains abundance values for each species in each plot. The most basic use of `abMatGen()` uses tree abundance as a measure of abundance, but other columns can be used as alternative abundance measures by providing them as parameters to `abundance`, such as aboveground biomass (`agb`) or basal area (`ba`). The function also accounts for variation in sampling effort across nested plots using the `fpc` columns. You can alter the abundance unit e.g. tree or stem, by supplying a parameter to `abundance_unit`.

```
abMatGen(stems_species, census_id = "census_id", abundance_unit = "tree_id",
  species_name = "species_name_clean", fpc = "fpc")[,1:4]
```

5.3.3 Stem biomass

Calculate aboveground biomass using the Chave, Réjou-Méchain et al. (2014) `chave14()` allometric equation, which requires both plot level longitude (`longitude_of_centre`) and latitude (`latitude_of_centre`) as well as stem diameter (`diam`) and species names.

Wood density data used by `chave14()` and `chave05()` can be generated separately using `woodDensity()`, which is a wrapper around `BIOMASS::getWoodDensity()`.

```

stems_species$wood_density <- woodDensity(stems_species)
stems_species$agb <- chave14(stems_species, plots_check)
stems_species$subdiv_id <- NA_character_

```

```

head(woodDensity(stems_species))

```

There are a number of other allometric equations available in the `{seosawr}` package which the user can use at their discretion (Table 5). There is also the option for the user to use their own allometric equation on the SEOSAW dataset.

The SEOSAW dataset contains `local_allometry` column in the plots table which allows data originators to suggest their own local allometry, though these allometries aren't available as functions in the `{seosawr}` package as they are numerous.

Function	Inputs	Usage	Reference
<code>ryan11()</code>	Diameter	Generic miombo woodland	(Ryan, Williams and Grace, 2011)
<code>mugasha14()</code>	Diameter	Tanzanian miombo woodland	(Mugasha et al., 2013)
<code>chidumayo90()</code>	Diameter	Zambian woodlands	(Chidumayo, 1990)
<code>mutakele09()</code>	Diameter	Mopane woodlands in Botswana	(Mutakela, 2009)
<code>chamshama04()</code>	Diameter	Tanzanian miombo woodlands	(Chamshama, Mugasha and Zahabu, 2004)
<code>tietema93()</code>	Diameter	Mopane woodlands in Botswana	(Tietema, 1993)
<code>goodman13()</code>	Diameter	Palms only, originally for the Amazon	(Goodman et al., 2013)
<code>tiepolo02()</code>	Height	Tree ferns (Cyatheaceae) only	(Tiepolo, Calmon and Feretti, 2002)
<code>chave05()</code>	Wood density Height Diameter	Global dry forests	(Chave, Andalo et al., 2005)
<code>chave14()</code>	Wood density Latitude Longitude Diameter	Global dry forests	(Chave, Réjou-Méchain et al., 2014)

Table 5: The various allometric equations to estimate aboveground woody biomass (AGB) provided in the `{seosawr}` R package.

The `{seosawr}` package relies on functions from the `{BIOMASS}` package to calculate the `chave14()` and `chave05()` allometries, which uses wood density data from (Chave, Coomes et al., 2009). All biomass allometry functions return aboveground biomass per stem in tonnes of dry matter.

5.3.4 Check logical consistency of plots and stems tables

Next, we are in a position to compare the logical consistency of the stem table with the plot table. Much like `stemTableCheck()` and `plotTableCheck()`, `stemPlotTableCheck()` conducts a number of checking procedures and flags rows in the stems table which don't agree with the plot level data, such as if a stem diameter is smaller than the minimum diameter threshold recorded for the plot.

```

stemPlotTableCheck(stems_species, plots_check)

```

The functions run by `stemPlotTableCheck()` are:

- `stemDiamThreshCheck()` - Are there stem measurements outside of the minimum or maximum stem diameter thresholds. Note that if a plot is nested with diameter thresholds, the highest of these thresholds is used and so may return warnings for stems in nested subplots.
- `censusIDMatchCheck()` - Check if all census IDs are present in both tables.
- `deadStemsSampledCheck()` - If `dead_stems_sampled = "FALSE"` are all stems alive (`stem_status = "a"`)?
- `stumpsSampledCheck()` - If `stumps_sampled = "FALSE"` are all stems not stumps (`stem_mode = "t"`)?

5.3.5 Generate plot level statistics from stem data

Then, you can calculate per census plot level statistics from the stems table. This includes calculations like the plot level aboveground biomass, the number of stems at different size thresholds, and the most dominant species in each plot:

```
plots_table_stat <- censusStemColGen(stems_species, plots_check)
plots_stat <- cbind(plots_check, plots_table_stat)
```

The statistics calculated are:

- `diam_max` - Maximum stem diameter
- `diam_min` - Minimum stem diameter
- `diam_mean` - Mean stem diameter
- `diam_median` - Median stem diameter
- `diam_sd` - Standard deviation of stem diameter distribution
- `height_max` - Maximum height
- `height_min` - Minimum height
- `height_mean` - Mean height
- `height_median` - Median height
- `height_sd` - Standard deviation of height
- `n_stems` - Number of stems
- `n_diam` - Number of stems with diameter measurements
- `n_height` - Number of stems with height measurements
- `lorey_height` - Lorey height
- `n_determ_genus` - Number of stems determined to genera
- `n_determ_species` - Number of stems determined to species

Additionally, diversity statistics are calculated using `diversityGen()` and the dominant species are identified using `domSpeciesGen()`:

- `richness` - Species richness
- `rarefy` - Rarefied species richness

- `shannon` - Shannon-Wiener diversity index H'
- `simpson` - Simpson diversity index D
- `invsimpson` - Inverse Simpson diversity index
- `evenness` - Species abundance evenness

`diamWeibullGen()` and `diamScaleGen()` are used to calculate diameter scaling factors, which describe the distribution of diameter values in a plot. `diamScaleGen()` bins diameters into 5 cm bins and takes the slope of a linear model of log-transformed bin frequencies as the scaling factor. `diamWeibullGen()` uses the scale and shape factors of a Weibull distribution of diameter measurements:

- `weibull_shape` - Shape parameter of Weibull distribution
- `weibull_scale` - Scale parameter of Weibull distribution
- `weibull_shape_se` - Standard error on shape parameter of Weibull distribution
- `weibull_scale_se` - Standard error on scale parameter of Weibull distribution
- `diam_scale` - Slope parameter of diameter bin frequency linear model
- `diam_scale_se` - Standard error on slope parameter of linear model

A number of statistics calculated as per-hectare variants:

- `n_stems_ha` - Number of stems per ha
- `ba_ha` - Basal area per ha
- `agb_ha` - Aboveground biomass per ha

Many of the above are replicated with subsets of data at various minimum stem diameter thresholds: >0, >3, >5, >10, >20, >40, >60 cm.

If you want to add any column from the plot data into the stem data, matching by `census_id`, use `plotToStem()`:

```
stems_analysis <- cbind(stems_species,
  plotToStem(stems_species, plots_stat, cols = c("agb_ha", "ba_ha"),
    stem_census_id = "census_id", plot_census_id = "census_id"))
```

Finally, the data is now in the SEOSAW format, with all metadata attached and some preliminary analysis already completed. You can export data using the `exportData()` function, or using the normal `write.csv()`, `st_write()` etc.. The advantage of `exportData()` is that it performs one last round of checking to make sure all required columns are present and correct in their values, even those that have been calculated during the cleaning process.

```
exportData("data_clean/example_data", stems = stems_species,
  plots = plots_stat, polys = plot_polys)
```

5.4 Compiling datasets

Once multiple datasets have been cleaned according to the process outlined above, they can be compiled to a single dataset using `compileAll()`:

```
compileAll(dirs = c("data_1", "data_2"),
  files = c("stems.csv", "plots.csv", "polys.shp"),
  out_dir = "../path/to/out/dir")
```

There are also two functions for extracting subsets of a compiled dataset. The first, `subsetLatest()` extracts the most recent census data for each plot, and `subsetMulti()` extracts measurements only for those plots which have multiple censuses:

```
subsetLatest(stems_species, plots_check,
  stem_plot_id = "plot_id", stem_census_date = "census_date",
  plot_census_date = "census_date")
```

`versionCheck()` allows you to check for differences in versions of a dataset. This is mostly used internally when creating the SEOSAW dataset, but might be useful elsewhere. The function takes two tables of plot data, and a vector of column names within which to check the values of each plot:

```
version_diff <- versionCheck(new_plots, old_plots, cols = c("agb_ha", "n_stems"))
```

5.5 Extracting spatial data

The `{seosawr}` package contains generic functions for extracting plot level data from arbitrary raster (`rasterExtract()`) and vector (`vectorExtract()`) spatial objects.

`rasterExtract()` can be used to extract field values from other user supplied raster datasets, with an optional circular buffer around the point of interest. `rasterExtract()` also allows for extraction from factor rasters using the `Mode()` function, which returns the modal value from raster cells within the buffer radius.

`vectorExtract()` is a similar function for extracting attributes from spatial vector datasets for points of interest.

```
plots_un <- ex_data$plots %>%
  dplyr::select(plot_id, longitude_of_centre, latitude_of_centre) %>%
  unique()
mplots_africa <- vectorExtract(plots_un, africa, attrib = "iso3")
```

The `{seosawr}` package also contains various convenience functions for manipulating spatial data. `{seosawr}` primarily uses `{sf}` spatial data objects, as this is likely the future of spatial analysis in R, but some functions such as `reprojectPoints()` will alternatively return a `SpatialPoints` object if requested.

A common scenario is that data originators like to store their plot coordinates as cartesian UTM coordinates for analysis in their local region, but SEOSAW requires all coordinates as WGS84 decimal latitude/longitude values (EPSG: 4326). As an example, imagine a dataset where plot coordinates straddle both UTM 37S and UTM 38S. We want to convert the plot coordinates to WGS84 latitude/longitude:

```
plots_un$utm_zone <- latLong2UTM(plots_un$longitude_of_centre, plots_un$latitude_of_centre)
plots_un_list <- split(plots_un, plots_un$utm_zone)

utm_crs <- unlist(lapply(plots_un_list, function(x) {
  UTMProj4(unique(x$utm_zone))
})))
```

```
plots_un_list_wgs <- lapply(1:length(plots_un_list), function(x) {  
  reprojectPoints(plots_un_list[[x]]$longitude_of_centre,  
    plots_un_list[[x]]$latitude_of_centre,  
    from = utm_crs[[x]], to = 4326)  
})  
  
plots_un_wgs <- do.call(rbind, plots_un_list_wgs)  
  
head(plots_un_wgs)
```

6 Dataset table columns

This section provides a comprehensive description of every column found in the SEOSAW dataset, across the plots table, stems table, small stems table, coarse woody debris table, herbaceous table, and spatial data table.

Table 6: Column names for plots table.

Column	Class	Description	Units	Example
plot_id	chr	Standardised method of identifying plots. A three letter code, followed by a number referring to the particular plot. Three letter codes are normally, but not always, created like this: Angola (A), in Bicuar National Park (B), by John Godlee (G). The three letter code should be conserved within a dataset, i.e. a group of plots owned by a particular researcher within a particular locale.	NA	SSM_1
census_date	chr	The date of the census. This can be formatted as either YYYY, YYYY-MM, or YYYY-MM-DD	date	2008
plot_cluster	chr	If plots occur in a spatial cluster, a single plot_id used to identify the cluster	NA	ABG_1
plot_name	chr	Plot name as designated by those who originally collected the data. 1, Wedza_01, MHR-04b are all valid. No spaces please.	NA	Wedza_01
country_iso3	fct	ISO 3166-1 alpha-3 code to refer to the country in which the plot is situated. Only countries within Africa. See 'data(africa_iso3)' for available country codes. use countrycode::countrycode() to convert a country name to an ISO3 code	NA	AGO
prinv	chr	Name of the Principal Investigator. This can be an organisation, a person's name, a surname, initials, etc.	NA	Godlee J. L.
other_authors	chr	Names of other authors who contributed to the collection of the dataset, separated by semi-colons	NA	Godlee J. L.; Maiato F.; Dexter K. G.
plot_shape	fct	Shape of the plot. Possible values are: "rectangle", "circle" or "irregular". Note that a square is a special case of "rectangle".	NA	rectangle
plot_width	int	If plot_shape == "rectangle", the shorter side length, in metres to the nearest metre	m	100
plot_length	int	If plot_shape == "rectangle", the longer side length, in metres to the nearest metre	m	50
plot_radius	int	If plot_shape == "circle", the radius of the plot, measured in metres to the nearest metre	m	25
subplot_n	int	Number of subplots within the larger plot. Subplots (as opposed to nested plots) have the same sampling effort as the entire plot, generally used for ease of relocating trees.	n	10
subplot_shape	fct	Shape of the subplots. Possible values are the same as plot_shape	NA	circle
subplot_length	int	If subplot_shape == "rectangle", the longest side length, in metres to the nearest metre	m	50
subplot_width	int	If subplot_shape == "rectangle", the shorter side length, in metres to the nearest metre	m	10

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
subplot_radius	int	If subplot_shape == "circle", the radius of the plot, measured in metres to the nearest metre	m	10
min_diam_thresh	int	Minimum stem diameter value considered within this plot, in centimetres to the nearest centimetre. If sampling is nested this should be the minimum value outside of all nested subplots.	cm	5
min_height_thresh	num	Minimum stem height value considered within this plot, in metres to the nearest 10 centimetres. If sampling is nested this should be the maximum value outside of all nested subplots.	m	1.5
nested_1	log	Does the plot have at least one level of nested sampling?	NA	TRUE
n1_min_diam_thresh	int	If nested_1 == TRUE, the minimum stem diameter value considered within the first nesting level, in centimetres to the nearest centimetre.	cm	10
n1_min_height_thresh	num	If nested_1 == TRUE, the minimum stem height value considered within the first nesting level, in metres to the nearest metre.	m	20
n1_plot_shape	fct	If nested_1 == TRUE, the shape of the subplot, same values as plot_shape	NA	circle
n1_plot_width	int	If nested_1 == TRUE & n1_plot_shape == "rectangle" the shorter side length, in metres to the nearest metre	m	20
n1_plot_length	int	If nested_1 == TRUE & n1_plot_shape == "rectangle" the longer side length, in metres to the nearest metre	m	10
n1_plot_radius	int	If nested_1 == TRUE & n1_plot_shape == "circle" the radius of the subplot, in metres to the nearest metre	m	14
nested_2	log	Does the plot have at least two levels of nested sampling?	NA	TRUE
n2_min_diam_thresh	int	If nested_2 == TRUE, the minimum stem diameter value considered within the second nesting level, in centimetres to the nearest centimetre.	cm	5
n2_min_height_thresh	num	If nested_2 == TRUE, the minimum stem height value considered within the second nesting level, in metres to the nearest metre.	m	2.5
n2_plot_shape	fct	If nested_2 == TRUE, the shape of the subplot, same allowed values as plot_shape	NA	rectangle
n2_plot_width	int	If nested_2 == TRUE & n2_plot_shape == "rectangle" the shorter side length, in metres to the nearest metre	m	30
n2_plot_length	int	If nested_2 == TRUE & n2_plot_shape == "rectangle" the longer side length, in metres to the nearest metre	m	20
n2_plot_radius	int	If nested_2 == TRUE & n2_plot_shape == "circle" the radius of the subplot, in metres to the nearest metre	m	5
nested_3	log	Does the plot have at least three levels of nested sampling?	NA	TRUE

Table 6: Column names for plots table. (*continued*)

Column	Class	Description	Units	Example
n3_min_diam_thresh	int	If nested_3 == TRUE, the minimum stem diameter value considered within the third nesting level, in centimetres to the nearest centimetre.	cm	5
n3_min_height_thresh	num	If nested_3 == TRUE, the minimum stem height value considered within the third nesting level, in metres to the nearest metre.	m	2.5
n3_plot_shape	fct	If nested_3 == TRUE, the shape of the subplot, same allowed values as plot_shape	NA	rectangle
n3_plot_width	int	If nested_3 == TRUE & n3_plot_shape == "rectangle" the shorter side length, in metres to the nearest metre	m	30
n3_plot_length	int	If nested_3 == TRUE & n3_plot_shape == "rectangle" the longer side length, in metres to the nearest metre	m	20
n3_plot_radius	int	If nested_3 == TRUE & n3_plot_shape == "circle" the radius of the subplot, in metres to the nearest metre	m	5
lianas_sampled	log	Were lianas sampled? I.e. woody stems that are not self-supporting and use other trees to grow towards the canopy	NA	TRUE
soil_sampled	log	Were soil samples collected?	NA	FALSE
herbaceous_sampled	log	Was the herbaceous layer sampled for grasses, forbs, shrubs, or small trees that did not fit the size thresholds for the plot?	NA	TRUE
dead_stems_sampled	log	Were dead stems sampled in this plot?	NA	FALSE
stumps_sampled	log	Were tree stumps sampled?	NA	TRUE
coarse_woody_debris_sampled	log	Was coarse woody debris sampled?	NA	FALSE
small_stems_sampled	log	Were woody stems smaller than the minimum size thresholds counted?	NA	TRUE
tags	log	Are stems within the plot tagged?	NA	FALSE
permanent	log	Is the plot a permanent plot (TRUE) or a one-off census (FALSE)? Could the plot be re-measured in the future?	NA	TRUE
abandoned	log	Is this the last census and the plot is now abandoned? Common reasons for abandonment: Clear-felling or nearly all trees removed. Often combined with tag loss and irreconcilable damage to the plot which would make it hard to match the stems with a previous census.	NA	TRUE
stem_timeline_cont	log	If tags == TRUE and permanent == TRUE, is it impossible to link stems between this census and previous censuses?	NA	FALSE
sampling_notes	chr	Extra notes relating to the sampling process	NA	Only stumps greater than 2 m tall were sampled

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
sampling_design_code	chr	File name for document containing the description of your methods/protocol for data collection, provided to SEOSAW, or a reference to such a document. If not documented then NA	NA	SSAW_V1.gdoc
plot_notes	chr	Include any long form comments about the plot here, if they don't fit into other columns.	NA	It looks like this plot is a rubbish dump for household waste.
longitude_of_centre	num	Longitude of the centre of gravity of the plot. If you have GPS points of the plot corners, send those and we will calculate this. Measured in decimal degrees, WGS84 EPSG:4326	deg	-50.425622
latitude_of_centre	num	Latitude of the centre of gravity of the plot. If you have GPS points of the plot corners, send those and we will calculate this. Measured in decimal degrees, WGS84 EPSG:4326	deg	30.446453
elevation	num	Elevation of the plot. Measured in metres above International Mean sea level to 1 decimal place (nearest 10 cm). In order of preference: 1) Average of GPS readings across the plot, 2) a single GPS reading, 3) Manual lookup on topographic map, 4) NA. We can query SRTM for missing values.	m asl	1234
slope_method	fct	Method used to determine plot slope. Allowed values: "rangefinder" if a laser or sonic rangefinder was used, "trig" if slope was estimated with tape measures and-or a clinometer, "eye" if estimated by eye, "other" if another method.	NA	trig
slope_method_notes	chr	Extra notes relating to the slope method	NA	20 points taken with 20 m minimum distance, randomly located
slope	num	Plot slope, measured in degrees from 0 (flat) to 90 (vertical)	deg	4.4
aspect	num	Aspect of the plot, i.e. which direction is the slope facing, in degrees (0-360)	deg	54
polygons	log	Polygon data of the plot shape provided by the data originator?	NA	TRUE
local_allometry	fct	Most appropriate aboveground woody biomass allometry for the plot. In order of preference: (1) DOI, (2) reference, (3) other URL, (4) shorthand allometry name	NA	ryan11
allometry_notes	chr	Extra notes relating to the appropriate local allometry	NA	ryan11 should be used for everything except the eucalyptus trees present in plot ABC_5

Table 6: Column names for plots table. (*continued*)

Column	Class	Description	Units	Example
catenal_position	fct	Where on the catena is plot? Possible values are:"ridge_crest", "concave_slope", "midslope", "convex_slope", "valley_floor", "flat"	NA	midslope
termites	int	If termites were counted, the number of termite mounds found on the plot	n	34
canopy_cover	prop	The proportion of the ground (periscope, aerial photo, canopy dimensions), or sky hemisphere (hemi-photo, mirror densitometer), covered by a tree canopy. Measured as a proportion between 0 and 1, e.g. 60% = 0.6	proportion	0.28
canopy_cover_method	fct	If canopy cover was estimated, what method was used? Possible values are: "periscope_densitometer", "mirror_densitometer", "hemispherical_photo", "aerial_photo", "canopy_dimensions", "other"	NA	hemispherical_photo
canopy_cover_method_notes	chr	Extra notes relating to the canopy cover method	NA	40 hemispherical photos in a regular grid with 20 m spacing. Analysed with Hemiphot.
xy_method	fct	Method used to estimate stem location within the plot, either grid or GPS coordinates. Possible values: "tape", "gps", "dgps", "angle" (angular direction and distance from plot centre in circular plot), "other".	NA	tape
manipulation_experiment	log	Has any aspect of the plot's land use or environment been manipulated by researchers?	NA	FALSE
experiment_ref	chr	If manipulation_experiment == TRUE, a link, citation, or contact details relating to the experimental manipulation of this plot. In order of preference: (1) DOI, (2) reference, (3) other URL, (4) contact details	NA	10.1890/06-1664
manipulation_experiment_notes	chr	If manipulation_experiment == TRUE, notes relating to the experimental treatment applied to the plot	NA	Tilling of soil 15-20 cm throughout
pa	log	Is the plot in a protected area?	NA	TRUE
pa_name	chr	If pa == TRUE, the name of the protected area. If the protected area is listed by the WDPA, please use the name provided by the WDPA	NA	Bicuar National Park
pa_type	chr	If pa == TRUE, the type of protected area. If the protected area is listed by the WDPA, use their type classification.	NA	Forest reserve

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
wdpa_id	chr	If pa == TRUE and the protected area is listed by the WDPA, the WDPA ID value for that protected area. See https://www.protectedplanet.net	NA	370
local_community_rights	chr	Explain the rights of the local community in using the plot for resource extraction, grazing, farming etc.	NA	Dead wood collection by local village
other_user_rights	chr	Explain the rights of any others in using the plot for resource extraction, grazing, farming etc.	NA	Yearly game drive in early May
timber_harvesting	log	Is there evidence of tree harvesting for timber products, e.g. planks?	NA	TRUE
high_graded_100_years	log	Is it likely that high quality timber trees were removed from the plot the last 100 years?	NA	FALSE
clear_cut	log	Has any part of the plot ever been clear cut (all trees removed)?	NA	TRUE
mining	log	Has any part of the plot ever been used for mining?	NA	FALSE
fuel_wood_harvesting	log	Is there evidence of fuel wood collection, excluding charcoal production?	NA	TRUE
charcoal_harvesting	log	Is there evidence of trees being harvested for charcoal production?	NA	FALSE
other_woody_product_harvesting	log	Is there evidence of people harvesting woody material for any other use, e.g. string, beehives, canoes?	NA	TRUE
ntfp_harvesting	log	Is there evidence of people harvesting non-wood tree products, e.g. mushrooms honey, fruit etc.?	NA	FALSE
farmed_30_years	log	Is it likely that the plot was used for farming (or cleared for another reason), in the last 30 years?	NA	TRUE
land_use_notes	chr	Extra notes relating to land use in the plot	NA	FALSE
meas_tree_stem	fct	Are measurements taken for each "tree", or each "stem"?	NA	tree
tag_tree_stem	fct	Are tag IDs at the "tree" or "stem" level?	NA	tree
tree_diff	log	Were stems grouped into trees, where trees have multiple stems?	NA	TRUE
fire_exclusion	log	Is there a fire exclusion experiment in this plot? If TRUE, manipulation_experiment should also = TRUE.	NA	TRUE
fire_exclusion_success	fct	If fire_exclusion == TRUE, to what extent were fires successfully excluded from the plot? Possible values: "all", "most", "few"	NA	all
fire_treatment	log	Is or was there a fire treatment experiment in this plot, e.g. controlled fires every two years? If TRUE, manipulation_experiment should also = TRUE.	NA	FALSE
fire_treatment_season	fct	If fire_treatment == TRUE, the season in which fires are set. Possible values: "early_dry_season", "late_dry_season", "early_wet_season", "late_wet_season"	NA	early_dry_season

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
fire_return_interval	int	Number of years between fires. If there has been some sort of fire control experiment, the treatment fire return interval measured in years, e.g. 1 for annual burning. If no treatment, the estimated mean interval between fires.	year	5
fire_last_date	chr	Date of the last fire.	date	2022
fire_cause	chr	What commonly causes fires in the plot? E.g. Management, accidental, lightning, agriculture.	NA	Adjacent field fires
fire_fuel	chr	What do fires on this plot typically burn? E.g. Canopy, understorey, litter	NA	Tree canopies
fire_notes	chr	Extra notes relating to fire regime or fire treatment	NA	Fires are set with drip torches in the late evening, after a heavy rain
elephants	log	Do elephants frequent the plot	NA	TRUE
elephant_density	num	If elephants are present, at what approximate density do they occur within the area? Measured as elephants per km ²	n km ⁻²	5
elephant_data_source	chr	A link, citation, or contact details relating to elephant density estimates. In order of preference: (1) DOI, (2) reference, (3) other URL, (4) contact details	NA	10.6088/ijes.002
exclosure	fct	Are herbivores excluded from the plot by a fence? If so, what sizes? Possible values are: "extra_large" (>1000 kg), "large" (100-1000 kg), "medium" (30-100 kg), "small" (<30 kg). Multiple values possible separated by semi-colon.	NA	extra_large
small_nonsocial_browsers	log	Do small non-social browsers frequent this plot? e.g. dik-diks, duikers and grysbok, steenbok (these categories are based on Hempson et al 2015). TRUE or FALSE	NA	TRUE
large_browsers	log	Do large browsers frequent this plot? e.g. giraffe, okapi, kudu	NA	FALSE
medium_sized_social_mixed_diet	log	Do medium sized social mixed diet herbivores frequent this plot? e.g. gazelle, oryx	NA	TRUE
nonruminants_excluding_suids	log	Do non-ruminant herbivores frequent this plot? e.g. zebra, hippo, rhino	NA	FALSE
water_dependent_grazers	log	Do water-dependent grazers frequent this plot? e.g. roan, sable, bluebuck, waterbuck, lechwe	NA	TRUE
goat_grazing	log	Do goats feed in this plot?	NA	FALSE
sheep_grazing	log	Do sheep feed in this plot?	NA	TRUE
cattle_grazing	log	Do domestic cattle feed in the plot?	NA	FALSE

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
herbivore_notes	chr	Extra notes relating to herbivores in the plot.	NA	Dik Diks, palankas, Olongo, elephants like to knock over the B. massaiensis
cyclone	log	Do cyclones (category 3+) ever occur in this plot?	NA	TRUE
cyclone_last_date	chr	When was the last cyclone?	NA	2022
drought	log	Do major droughts ever occur in this plot?	NA	FALSE
drought_last_date	chr	When was the last major drought?	NA	2021
wind	log	Do wind storms or tornadoes ever occur in this plot?	NA	TRUE
wind_last_date	chr	When was the last wind storm or tornado?	NA	2020
earthquake	log	Do earthquakes ever occur in this plot?	NA	FALSE
earthquake_last_date	chr	When was the last earthquake?	NA	2019
landslide	log	Do landslides ever occur in this plot?	NA	TRUE
landslide_last_date	chr	When was the last landslide?	NA	2018
flood	log	Do major floods ever occur in this plot?	NA	FALSE
flood_last_date	chr	When was the last major flood?	NA	2017
frost	log	Do major frost events (<-2C for three consecutive nights) ever occur in this plot?	NA	TRUE
frost_last_date	chr	When was the last major frost event?	NA	2016
data_ref	chr	A link, citation, or contact details relating to the publication of this plot's data. In order of preference: (1) DOI, (2) reference, (3) other URL, (4) contact details	NA	10.3390/d12040140
photos	chr	Filenames of photos of the plot, separated by semi-colons	NA	DSC_134.jpg;DSC_-135.jpg
date_metadata	num	Year that dataset metadata values, i.e. herbivory, land use, plot location, fire regime etc. were last updated	NA	2020
file_plots	chr	Filepath of the raw plot metadata file provided to SEOSAW. Include multiple files in a semi-colon separated list.	NA	archibald_-kruger.xlsx;Kruger_-2024.csv
file_stems	chr	Filepath of the raw stem data file provided to SEOSAW. Include multiple files in a semi-colon separated list.	NA	Archibald_-kruger.csv
subdiv	log	Is the plot subdivided into smaller 1 ha square parcels? If TRUE, plot-level calculated values can be found in plots_subdiv.csv	NA	TRUE
dataset_id	chr	The dataset ID, normally defined as the three letter code in plot_id	NA	SSM
plot_area	num	Flat area of the plot, in hectares	ha	0.25
subplot_area	num	Flat area of the subplots, in hectares	ha	0.005
plot_perimeter	num	Perimeter of the plot, in metres	m	400

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
census_year	chr	The year of the census (YYYY)	NA	2022
census_id	chr	A standardised method of identifying censuses. A composite of 'plot_id' 'census year'	NA	ABG_23_2022
diam_max	num	Maximum stem diameter found in the plot	cm	50.4
diam_min	num	Minimum stem diameter found in the plot	cm	5
diam_mean	num	Mean stem diameter found in the plot	cm	5.6
diam_median	num	Median stem diameter found in the plot	cm	10.3
diam_sd	num	Standard deviation of stem diameter found in the plot	cm	1.5
diam_ge5_mean	num	Mean stem diameter found in the plot, greater than 5 cm diameter	cm	5.4
diam_ge5_median	num	Median stem diameter found in the plot, greater than 5 cm diameter	cm	10.6
diam_ge5_sd	num	Standard deviation of stem diameter found in the plot, greater than 5 cm diameter	cm	1.5
diam_ge10_mean	num	Mean stem diameter found in the plot, greater than 10 cm diameter	cm	15.7
diam_ge10_median	num	Median stem diameter found in the plot, greater than 10 cm diameter	cm	15.9
diam_ge10_sd	num	Standard deviation of stem diameter found in the plot, greater than 10 cm diameter	cm	2.5
diam_ge20_mean	num	Mean stem diameter found in the plot, greater than 20 cm diameter	cm	15.7
diam_ge20_median	num	Median stem diameter found in the plot, greater than 20 cm diameter	cm	15.9
diam_ge20_sd	num	Standard deviation of stem diameter found in the plot, greater than 20 cm diameter	cm	2.5
n_diam	num	Number of stems for which diameter was measured	n	524
n_height	num	Number of stems for which height was measured	n	532
height_max	num	Maximum stem height found in the plot	m	10.5
height_min	num	Minimum stem height found in the plot	m	1.6
height_mean	num	Mean stem height found in the plot	m	1.6
height_median	num	Median stem height found in the plot	m	5.4
height_sd	num	Standard deviation stem height found in the plot	m	4.5
height_ge5_mean	num	Mean stem height found in the plot, greater than 5 cm diameter	m	3.5
height_ge5_median	num	Median stem height found in the plot, greater than 5 cm diameter	m	5.4
height_ge5_sd	num	Standard deviation of stem height found in the plot, greater than 5 cm diameter	m	1.5
height_ge10_mean	num	Mean stem height found in the plot, greater than 10 cm diameter	m	3.5
height_ge10_median	num	Median stem height found in the plot, greater than 10 cm diameter	m	6.6
height_ge10_sd	num	Standard deviation of stem height found in the plot, greater than 10 cm diameter	m	1.4

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
height_ge20_mean	num	Mean stem height found in the plot, greater than 20 cm diameter	m	3.5
height_ge20_median	num	Median stem height found in the plot, greater than 20 cm diameter	m	6.6
height_ge20_sd	num	Standard deviation of stem height found in the plot, greater than 20 cm diameter	m	1.4
height_per95	num	95th percentile of stem height, i.e. the height below which 95% of stems can be found	m	4.3
n_stems	num	Number of stems in the plot	n	5025
n_stems_ge5	num	Number of stems greater than 5 cm diameter	n	5010
n_stems_ge10	num	Number of stems greater than 10 cm diameter	n	2040
n_stems_ge20	num	Number of stems greater than 20 cm diameter	n	505
n_stems_ge40	num	Number of stems greater than 40 cm diameter	n	20
n_stems_ge60	num	Number of stems greater than 60 cm diameter	n	1
lorey_height	num	Lorey height of the plot. Calculated as the sum of basal areas multiplied by height, divided by the total basal area. This variant is adjusted for nested sampling with fpc	NA	3.5
n_determ_genus	num	Number of stems where genera successfully identified	n	20
n_determ_species	num	Number of stems where species epithets successfully identified	n	49
dom_1	chr	Most dominant species in the plot, by stem abundance	NA	Brachystegia spiciformis
dom_2	chr	Second most dominant species in the plot, by stem abundance	NA	Ochna pulchra
dom_3	chr	Third most dominant species in the plot, by stem abundance	NA	Burkea africana
dom_4	chr	Fourth most dominant species in the plot, by stem abundance	NA	Indet indet
dom_5	chr	Fifth most dominant species in the plot, by stem abundance	NA	Pseudolachnostylis maprouneifolia
prop_1	num	Proportional abundance of the most dominant species in the plot	proportion	20.4
prop_2	num	Proportional abundance of the second most dominant species in the plot	proportion	14.5
prop_3	num	Proportional abundance of the third most dominant species in the plot	proportion	12.7
prop_4	num	Proportional abundance of the fourth most dominant species in the plot	proportion	8.4
prop_5	num	Proportional abundance of the fifth most dominant species in the plot	proportion	3.6
richness	num	Species richness of plot	n	53
rarefy	num	Rarefied species richness at 25 individuals	NA	42.65
shannon	num	Shannon diversity index of plot	NA	4.3
simpson	num	Simpson diversity index of plot	NA	2.5

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
invsimpson	num	Inverse Simpson diversity index of plot	NA	2.6
evenness	num	Abundance evenness of plot	NA	2.6
stocking_ha	num	Number of stems per hectare, adjusted for nested sampling with fpc	n ha ⁻¹	50.2
stocking_ge5_ha	num	Number of stems greater than 5 cm diameter per hectare, adjusted for nested sampling with fpc	n ha ⁻¹	45.5
stocking_ge10_ha	num	Number of stems greater than 10 cm diameter per hectare, adjusted for nested sampling with fpc	n ha ⁻¹	32.5
stocking_ge20_ha	num	Number of stems greater than 20 cm diameter per hectare, adjusted for nested sampling with fpc	n ha ⁻¹	32.5
ba_ha	num	Basal area per hectare, adjusted for nested sampling with fpc	m ² ha ⁻¹	3.6
agb_ha	num	Aboveground woody biomass per hectare, for living stems only (stem_status == "a" is.na(stem_status)), adjusted for nested sampling with fpc	t DM ha ⁻¹	42.6
agb_ge5_ha	num	Aboveground woody biomass per hectare for stems greater than 5 cm diameter, adjusted for nested sampling with fpc	t DM ha ⁻¹	42.6
agb_ge10_ha	num	Aboveground woody biomass per hectare for stems greater than 10 cm diameter, adjusted for nested sampling with fpc	t DM ha ⁻¹	42.6
agb_ge20_ha	num	Aboveground woody biomass per hectare for stems greater than 20 cm diameter, adjusted for nested sampling with fpc	t DM ha ⁻¹	42.6
diam_scale	num	Diameter distribution scaling parameter. The slope of a linear model of log-transformed diameter class frequency values in 5 cm diameter bins.	NA	1.55
diam_scale_se	num	Standard error on the slope parameter of the diameter distribution scaling parameter.	NA	0.544
weibull_scale	num	Scale parameter of a two-parameter diameter Weibull probability distribution.	NA	1.47
weibull_shape	num	Shape parameter of a two-parameter diameter Weibull probability distribution.	NA	0.63
weibull_scale_se	num	Standard error on the scale parameter of the diameter class distribution	NA	0.444
weibull_shape_se	num	Standard error on the shape parameter of the diameter class distribution	NA	0.135
baobab	log	Do members of the Adansonia genus comprise ≥ 10% of the aboveground biomass? This column is included because Adansonia spp. Peculiar growth form can drastically inflate the aboveground biomass of a plot	NA	TRUE
stocking	num	Number of stems per plot adjusted for nested sampling with fpc	n	54.23

Table 6: Column names for plots table. *(continued)*

Column	Class	Description	Units	Example
stocking_ge5	num	Number of stems greater than 5 cm diameter per plot, adjusted for nested sampling with fpc	n	43.2
stocking_ge10	num	Number of stems greater than 10 cm diameter per plot, adjusted for nested sampling with fpc	n	34.5
stocking_ge20	num	Number of stems greater than 20 cm diameter per plot, adjusted for nested sampling with fpc	n	3.24
ba	num	Basal area per plot, adjusted for nested sampling with fpc	m ² ha ⁻¹	3.2
agb	num	Aboveground woody biomass per plot, adjusted for nested sampling with fpc	t DM ha ⁻¹	5.4
agb_ge5	num	Aboveground woody biomass per plot for stems greater than 5 cm diameter, adjusted for nested sampling with fpc	t DM ha ⁻¹	42.6
agb_ge10	num	Aboveground woody biomass per plot for stems greater than 10 cm diameter, adjusted for nested sampling with fpc	t DM ha ⁻¹	42.6
agb_ge20	num	Aboveground woody biomass per plot for stems greater than 20 cm diameter, adjusted for nested sampling with fpc	t DM ha ⁻¹	42.6

Table 7: Column names for stems table.

Column	Class	Description	Units	Example
plot_id	chr	Unique identifier for plots. A three letter code, followed by a number referring to the particular plot. Three letter codes are normally, but not always, created like this: Angola (A), in Bicuar National Park (B), by John Godlee (G). The three letter code should be conserved within a dataset, i.e. a group of plots owned by a particular researcher in a particular locale.	NA	ABG_1
subplot_id	chr	If the plot is divided into multiple parcels, but not "nested", i.e. sampling effort is equal across all parcels, the parcel within which the stem is found.	NA	A
tree_id	chr	Unique identifier for trees within a plot, normally the first stem ID within the tree.	NA	45
stem_id	chr	Unique identifier for stems within a plot.	NA	6
tag_id	chr	The code of the tag attached to the tree or stem in the field. If tags were not used, leave this column blank	NA	BP094
census_date	chr	The date of the census. This can be either YYYY, YYYY-MM, or YYYY-MM-DD.	NA	26/06/2022
species_orig_binom	chr	The binomial name of the tree species, as provided by the data originator	NA	Burkea africana
species_orig_local	chr	The local name of the tree species/morphospecies, as provided by the data originator	NA	mumwei
notes_stem	chr	Miscellaneous notes related to the tree or stem	NA	Leaning around 45 degrees and with a beehive in the fluted trunk
diam	num	Stem diameter measurement in cm	cm	6.4
pom	num	The height at which diam was taken, in metres to one decimal place (nearest 10 cm)	m	1.3
x_grid	num	The x coordinate of the stem within the plot if a grid method was used. In metres to nearest 10 cm	m	54.2
y_grid	num	The y coordinate of the stem within the plot if a grid method was used. In metres to nearest 10 cm	m	35.6
latitude	num	The latitude coordinate of the stem within the plot if a GPS method was used. WGS84 EPSG4326	decimal degrees	-50.425622
longitude	num	The longitude coordinate of the stem within the plot if a GPS method was used. WGS84 EPSG4326	decimal degrees	30.446453

Table 7: Column names for stems table. (*continued*)

Column	Class	Description	Units	Example
angle	num	If 'plot_shape' == "circle", and angular stem coordinates were used, the angle (0 = North) direction of the stem from the plot centre, in degrees (0-360)	degrees	43.2
distance	num	If 'plot_shape' == "circle", and angular stem coordinates were used, the distance of the stem from the plot centre. In metres to the nearest 10 cm	m	3.5
height	num	The height of the tree, in metres to one decimal place (nearest 10 cm)	m	3.5
bole_height	num	The height from the ground to the base of the lowest branch also called 'clear length'	m	6.3
crown_x	num	If two crown diameters are taken perpendicular to one another, the first of these measurements. If one crown diameter was taken, that crown diameter, in metres to one decimal place (nearest 10 cm).	m	5.2
crown_y	num	If two crown diameters are taken perpendicular to one another, the second of these measurements. If one crown diameter was taken, this remains blank, in metres to one decimal place (nearest 10 cm).	m	3.5
stump_diam_top	num	If the stem is a stump, the diameter at the top of the stump, in cm.	cm	10.4
stump_diam_base	num	If the stem is a stump, the diameter at the base of the stump (30 cm), in cm.	cm	12.5
stem_status	fct	What is the growth status of the stem. Possible values are: "r" if the stem is topkilled and resprouting, "d" if the stem and the rest of the tree is completely dead (i.e. base appears dead and all stems are dead), "a" if the stem is alive with no abnormal growth status.	NA	r
broken_per_remain	num	If the stem is broken, approximately how much is remaining? As a decimal, e.g. 60% of stem remaining = 0.6, to one decimal place (nearest 10%)	proportion	0.2
broken_height	num	If the stem is broken, at what height is it broken. In metres to one decimal place (nearest 10 cm)	m	4.5
wound_area	num	If the stem is wounded, the area of the wound.	cm ²	6

Table 7: Column names for stems table. (*continued*)

Column	Class	Description	Units	Example
stem_mode	fct	Various flags denoting stem mode. Possible values are: "b" - broken (material above break is detached), "c" - has climbing or strangling or liana plants in canopy, "f" - fallen, "g" - hollow, "j" - bark peeling or lost, "k" - stem has bracket fungi, "l" - leaning (>15deg), "m" - on a termite mound, "n" - a new recruit, "p" - snapped (material above break is attached), "q" - the stem location is unknown and was not found (contrast with "v"), "s" - standing, "t" - stump i.e. broken below POM, "u" - uprooted, "v" - vanished (location of tree found but stem missing), "w" - stem is a liana, "y" - wounded, "z" - sick (i.e. lost most leaves, discoloured foliage, drooping branches) or rotten wood	NA	P
damage_cause	fct	If the stem is damaged, what caused it? Possible values are: "a" if animals (not elephants, not insects), "e" if elephants, "f" if fire, "h" if humans (e.g. cut, ringbarked, broken or beehive), "i" if frost, "l" if lightning, "m" if termites, "n" if neighbouring tree, "o" if other (describe in damage_cause_notes), "q" if unknown, "w" if wind	NA	N
damage_cause_notes	chr	If damage_cause = "o", describe the specific method of damage.	NA	chainsaw marks and machete on small branches
cambium	num	The proportion of trunk surface area with exposed vascular cambium, between the ground and 2 m.	proportion	0.22
decay	num	What level of decay is the stem? Scale runs from 1 (no decay) to 5 (very decayed). Follow the "Decay class" table in the SEOSAW protocol. To 0.5 of a decay level (e.g. 1.5, 1, 3.5)	NA	4.5
diam_method	fct	What method was used to derive the stem diameter measurement? Possible values are: "dbh_tape" if a tape measure with a diameter conversion scale was used, "tape" if a normal tape measure was used and the circumference converted to diameter, "caliper_1" if a single caliper measurement was used, "caliper_2" if two caliper measurements at 90 degrees with the mean taken, "other".	NA	dbh_tape

Table 7: Column names for stems table. (*continued*)

Column	Class	Description	Units	Example
height_method	fct	What method was used to derive the height measurement? Possible values are: "eye" if estimated by eye, "rangefinder" if estimated using laser or ultrasonic apparatus with an electronic tilt sensor e.g. Nikon Forestry Pro, "clinometer" if estimated using a mechanical Clinometer, "hyposometer" if estimated using a laser vertex hypsometer from below the tree using last return, "direct" if estimated by climbing the tree or adjacent tower and measuring the height with tape or stick, "other"	NA	eye
voucher_id	chr	If a herbarium voucher has been collected, what is the code to identify it? 1, Wedza_01, MHR-04b are all valid. No spaces please.	NA	KEW_FK_42
next_census_check	chr	If any measured values seem wrong or, note checks to be performed during the next census.	NA	Appears to have resurrected check mortality
fpc	num	Fractional Probability of Inclusion. If plots are nested with varying sampling effort within plots for stems of different size, the probability of a stem this size occurring in the whole plot.	proportion	0.35
census_year	chr	The year of the census (YYYY)	NA	2022
census_id	chr	A standardised method of identifying censuses. A composite of 'plot_id' 'census_year'	NA	ABG_23_2022
measurement_id	chr	Unique identifier for stem measurements. A composite of 'plot_id' 'tree_id' 'stem_id' 'census_year'	NA	ABG_1_40_3_1990
crown_area	num	Area of the tree crown, measured as an ellipse using either one or two crown diameter measurements, if present. In m ²	m ²	4.5
agb	num	Aboveground biomass of stem, in tonnes of dry matter	t DM	0.093
wood_density	num	Wood density of stem, in g cm ⁻³	g cm ⁻³	0.042
ba	num	Basal area of stem, in m ²	m ²	0.46
diam_adj	num	Adjusted stem diameter at 1.3 m height accounting for tree taper	cm	5.3
species_name_sanit	chr	Species name after automated taxonomy cleaning by sanitizeTaxa()	NA	Protea cf. petiolaris
species_name_clean	chr	Species name after taxize checks and synonymy corrections	NA	Burkea africana
genus_clean	chr	Cleaned genus name, taken from 'species_name_clean'	NA	Burkea
species_clean	chr	Cleaned species epithet, taken from 'species_name_clean'	NA	africana
subspecies_clean	chr	Cleaned subspecies epithet, taken from 'species_name_clean'	NA	africana
confer_clean	chr	Cleaned cf. part of species epithet, taken from 'species_name_clean'	NA	polyantha
variety_clean	chr	Cleaned var. part of species epithet, taken from 'species_name_clean'	NA	micrantha
family_gnr	chr	Matched family name using taxize::gnr_resolve()	NA	Fabaceae

Table 7: Column names for stems table. (*continued*)

Column	Class	Description	Units	Example
species_name_gnr	chr	Name matched during the GNR resolve	NA	Burkea africana
taxon_rank_gnr	chr	All rank names of classification from GNR resolve, separated by " "	NA	superkingdom kingdom phylum subphylum
taxon_gnr	chr	All ranks of classification from GNR resolve, separated by " "	NA	Eukaryota Viridiplantae Streptophyta Stratiotales Burkeales Burkeaceae Burkea makondensis
base_rank_gnr	chr	The lowest rank identified in GNR resolve	NA	species
species_name_cjb	chr	Species name matched with the CJB African Plants database	NA	Burkea africana
subdiv_id	chr	If the plot has been divided into smaller 1 ha subdivisions, the names of 1 ha plot subdivisions to which the stem belongs.	NA	ABG_1_S4

Table 8: Column names for small stems table.

Column	Class	Description	Units	Example
plot_id	chr	Standardised method of identifying plots. A three letter code, followed by a number referring to the particular plot. Three letter codes are normally, but not always, created like this: Angola (A), in Bicuar National Park (B), by John Godlee (G). The three letter code should be conserved within a dataset, i.e. a group of plots owned by a particular researcher within a particular locale.	NA	SSM_1
census_date	chr	The date the measurement was taken. This can be formatted as either YYYY, YYYY-MM, or YYYY-MM-DD	NA	05/05/2022
transect_id	chr	ID of transect	NA	T1
transect_length	num	Length of transects, in m to nearest 10 cm. If ‘transect_id’ is not provided, the total length of all transects in the plot.	m	100
transect_width	num	Width of belt transect, in m to nearest 10 cm	m	2
species_orig_binom	chr	The binomial name of the tree species, as provided by the data originator	NA	Burkea africana
species_orig_local	chr	The local name of the tree species/morphospecies, as provided by the data originator	NA	mumwei
n_alive	int	Number of living stems found for a given species	NA	50
n_dead	int	Number of dead stems found for a given species	NA	4
notes_small_stems	chr	Miscellaneous notes on each species of small stem	NA	Could be Combretum collinum
census_year	chr	The year of the census (YYYY)	NA	2022
census_id	chr	A standardised method of identifying censuses. A composite of ‘plot_id’_‘census year’	NA	ABG_23_2022

Table 9: Column names for coarse woody debris table.

Column	Class	Description	Units	Example
plot_id	chr	Standardised method of identifying plots. A three letter code, followed by a number referring to the particular plot. Three letter codes are normally, but not always, created like this: Angola (A), in Bicuar National Park (B), by John Godlee (G). The three letter code should be conserved within a dataset, i.e. a group of plots owned by a particular researcher within a particular locale.	NA	SSM_1
census_date	chr	The date the measurement was taken. This can be formatted as either YYYY, YYYY-MM, or YYYY-MM-DD	NA	05/05/2022
transect_id	chr	ID of transect	NA	NW
transect_length	num	Length of transects, in m to nearest 10 cm. If ‘transect_id’ is not provided, the total length of all transects in the plot.	m	100
piece_id	chr	ID of piece of coarse woody debris. A piece is a single contiguous bit.	NA	1
section_id	chr	ID of section of coarse woody debris. A section is part of a piece.	NA	BP094
tag_id	chr	The code of any tags attached to the piece in the field.	NA	BP094
species_orig_binom	chr	The binomial name of the tree species, as provided by the data originator	NA	Burkea africana
species_orig_local	chr	The local name of the tree species/morphospecies, as provided by the data originator	NA	mumwei
section_diam	num	Diameter of section of CWD at point of intersection with transect, in cm.	cm	6.8
section_length	num	Length of section of CWD, in m to nearest cm	m	4.5
decay	num	What level of decay is the stump? Scale runs from 1 (no decay) to 5 (very decayed). Follow the "Decay class" table in the SEOSAW protocol. To 0.5 of a decay level (e.g. 1.5, 1, 3.5)	NA	4.5
notes_cwd	chr	Miscellaneous notes related to the piece of CWD	NA	Approximately half of the trunk has broken off
census_year	chr	The year of the census (YYYY)	NA	2022
census_id	chr	A standardised method of identifying censuses. A composite of ‘plot_id’_‘census year’	NA	ABG_23_2022

Table 10: Column names for herbaceous biomass table.

Column	Class	Description	Units	Example
plot_id	chr	Standardised method of identifying plots. A three letter code, followed by a number referring to the particular plot. Three letter codes are normally, but not always, created like this: Angola (A), in Bicular National Park (B), by John Godlee (G). The three letter code should be conserved within a dataset, i.e. a group of plots owned by a particular researcher within a particular locale.	NA	SSM_1
census_date	chr	The date the measurement was taken. This can be formatted as either YYYY, YYYY-MM, or YYYY-MM-DD	NA	11/06/2022
quadrat_id	chr	Unique identifier for quadrats within a plot	NA	Q1.1
quadrat_area	num	Area covered by the quadrat.	m ²	0.25
sward_height	num	Height of the grass sward to nearest cm	cm	25
bare_cover	num	Proportional bare ground cover. Bare ground has no litter, just bare soil.	proportion	0.45
rock_cover	num	Proportional bare rock cover.	proportion	0.73
dung	log	Is there animal dung larger than a human fist inside the quadrat?	NA	TRUE
fire	log	Is there evidence of a previous fire inside the quadrat, e.g. charcoal.	NA	FALSE
under_canopy	log	Is the quadrat directly under a tree canopy?	NA	TRUE
canopy_cover	prop	The proportion of the ground (periscope, aerial photo, canopy dimensions), or sky hemisphere (hemi-photo, mirror densitometer), covered by a tree canopy. Measured as a proportion between 0 and 1, e.g. 60% = 0.6	proportion	0.28
canopy_cover_method	fct	Possible values are: "periscope_densitometer", "mirror_densitometer", "hemispherical_photo", "aerial_photo", "canopy_dimensions", "other"	NA	periscope_densitometer
lai_above	num	Leaf Area Index above the grass sward, measured with an AccuPAR Ceptometer.	m ² m ⁻²	1.72
lai_below	num	Leaf Area Index below the grass sward, measured with an AccuPAR Ceptometer.	m ² m ⁻²	4.55
litter_depth	num	Depth of leaf litter inside the quadrat, to nearest cm.	cm	12
grass_wet_mass	num	The wet mass of grass material collected inside the quadrat.	g	52.5
other_leaf_wet_mass	num	The wet mass of non-grass leaf material collected inside the quadrat.	g	25.6
wood_wet_mass	num	The wet mass of woody litter collected inside the quadrat.	g	42.8
litter_wet_mass	num	The wet mass of leaf litter collected inside the quadrat.	g	24.7
succ_wet_mass	num	The wet mass of succulent material collected inside the quadrat.	g	35.6
grass_dry_mass	num	The dry mass of grass material collected inside the quadrat.	g	5.2
other_leaf_dry_mass	num	The dry mass of non-grass leaf material collected inside the quadrat.	g	2.5
wood_dry_mass	num	The dry mass of woody litter collected inside the quadrat.	g	4.2

Table 10: Column names for herbaceous biomass table. *(continued)*

Column	Class	Description	Units	Example
litter_dry_mass	num	The dry mass of leaf litter collected inside the quadrat.	g	2.4
succ_dry_mass	num	The dry mass of succulent material collected inside the quadrat.	g	3.5
notes_quadrat	chr	Miscellaneous notes on each quadrat	NA	Pig diggings inside quadrat
census_year	chr	The year of the census (YYYY)	NA	2022
census_id	chr	A standardised method of identifying censuses. A composite of ‘plot_id’ ‘census year’	NA	ABG_23_2022
grass_moisture	num	Moisture content of grass	g	3.6
other_leaf_moisture	num	Moisture content of other leaves	g	2.8
wood_moisture	num	Moisture content of woody litter	g	2.6
litter_moisture	num	Moisture content of leaf litter	g	7.4
succ_moisture	num	Moisture content of succulent material	g	7.3

Table 11: Column names for plot spatial data table.

Column	Class	Description	Resolution	Citation	Units
plot_id	chr	A standardised method of identifying plots. For example, the first plot in a set of plots installed in Angola (A), in Bicular National Park (B), by John Godlee (G), would be referred to as ABG_1. The three letter code should be conserved within a dataset, i.e. a group of plots owned by a particular researcher within a particular locale.	NA	NA	NA
bio1	num	WorldClim - Mean Annual Temperature	30s	Fick et al. 2017	degC
bio2	num	WorldClim - Mean Diurnal Range (mean monthly max. temp. - min. temp.)	30s	Fick et al. 2017	degC
bio3	num	WorldClim - Isothermality (bio2 / bio7)	30s	Fick et al. 2017	degC
bio4	num	WorldClim - Temperature Seasonality (standard deviation of bio1)	30s	Fick et al. 2017	degC
bio5	num	WorldClim - Max Temperature of Warmest Month	30s	Fick et al. 2017	degC
bio6	num	WorldClim - Min Temperature of Coldest Month	30s	Fick et al. 2017	degC
bio7	num	WorldClim - Temperature Annual Range (mean of annual bio5 - bio6)	30s	Fick et al. 2017	degC
bio8	num	WorldClim - Mean Temperature of Wettest Quarter	30s	Fick et al. 2017	degC
bio9	num	WorldClim - Mean Temperature of Driest Quarter	30s	Fick et al. 2017	degC
bio10	num	WorldClim - Mean Temperature of Warmest Quarter	30s	Fick et al. 2017	degC
bio11	num	WorldClim - Mean Temperature of Coldest Quarter	30s	Fick et al. 2017	degC

Table 11: Column names for plot spatial data table. (*continued*)

Column	Class	Description	Resolution	Citation	Units
bio12	num	WorldClim - Mean Annual Precipitation	30s	Fick et al. 2017	mm y ⁻¹
bio13	num	WorldClim - Precipitation of Wettest Month	30s	Fick et al. 2017	mm
bio14	num	WorldClim - Precipitation of Driest Month	30s	Fick et al. 2017	mm
bio15	num	WorldClim - Precipitation Seasonality (Coefficient of Variation of bio12)	30s	Fick et al. 2017	mm
bio16	num	WorldClim - Precipitation of Wettest Quarter	30s	Fick et al. 2017	mm
bio17	num	WorldClim - Precipitation of Driest Quarter	30s	Fick et al. 2017	mm
bio18	num	WorldClim - Precipitation of Warmest Quarter	30s	Fick et al. 2017	mm
bio19	num	WorldClim - Precipitation of Coldest Quarter	30s	Fick et al. 2017	mm
phen_avg_vi	num	Mean annual EVI (Enhanced Vegetation Index)	5600 m	Didan and Barretto 2016	NA
phen_cum_vi	num	Cumulative annual EVI (Enhanced Vegetation Index)	5600 m	Didan and Barretto 2016	NA
phen_day_peak_1	num	Mean day of year of peak EVI of growing season 1	5600 m	Didan and Barretto 2016	NA
phen_day_peak_2	num	Mean day of year of peak EVI of growing season 2	5600 m	Didan and Barretto 2016	NA
phen_day_peak_3	num	Mean day of year of peak EVI of growing season 3	5600 m	Didan and Barretto 2016	NA
herbiv_total	num	Mean total herbivore biomass	0.5deg	Hempson et al. 2017	kg km ⁻²
herbiv_small_nonsocial_browser	num	Mean total Small non-social browsers biomass	0.5deg	Hempson et al. 2017	kg km ⁻²
herbiv_medium_sized_social	num	Mean total Medium-sized social mixed herbivore biomass	0.5deg	Hempson et al. 2017	kg km ⁻²
herbiv_large_browser	num	Mean total Large browser biomass	0.5deg	Hempson et al. 2017	kg km ⁻²
herbiv_water_dependent	num	Mean total Water-dependent grazers biomass	0.5deg	Hempson et al. 2017	kg km ⁻²
herbiv_non_ruminant	num	Mean total Non-ruminants biomass	0.5deg	Hempson et al. 2017	kg km ⁻²

Table 11: Column names for plot spatial data table. (*continued*)

Column	Class	Description	Resolution	Citation	Units
herbiv_elephant	num	Mean elephant biomass	0.5deg	Hempson et al. 2017	kg km ⁻²
elevation	num	The elevation of the plot. Measured in metres above International Mean sea level to 1 decimal place. From SRTM	3s	Jarvis et al. 2008	m asl
tpi	num	Topographic Position Index - difference between cell elevation and mean value of 8 surrounding cells	1000 m	Amatulli et al. 2018	m asl
tri	num	Terrain Ruggedness Index - mean of absolute difference between cell elevation and value of 8 surrounding cells	1000 m	Amatulli et al. 2018	m asl
roughness	num	Topographic Roughness Index - difference between maximum and minimum cell elevation and its 8 surrounding cells	1000 m	Amatulli et al. 2018	m asl
slope	num	Slope	1000 m	Amatulli et al. 2018	degrees
admin_region	chr	Administrative area at level 1 according to rnatuarearth data	NA	South 2017	NA
soil_bulk_density	num	Bulk density of soil (fine earth)	250 m	Hengl et al. 2017	kg m ⁻³
soil_cation_ex_cap	num	Cation Exchange Capacity of soil	250 m	Hengl et al. 2017	cmolc kg ⁻¹
soil_clay	num	Clay content of soil (0-2 um) mass fraction	250 m	Hengl et al. 2017	proportion
soil_sand	num	Sand content of soil (50-2000 um) mass fraction	250 m	Hengl et al. 2017	proportion
soil_silt	num	Silt content of soil (2-50 um) mass fraction	250 m	Hengl et al. 2017	proportion
soil_coarse_frag	num	Coarse fragments volumetric in soil	250 m	Hengl et al. 2017	proportion
soil_ph	num	Soil pH x 10 in H2O	250 m	Hengl et al. 2017	NA
soil_org_c	num	Soil organic carbon stock	250 m	Hengl et al. 2017	Mg ha ⁻¹
soil_org_c_densit	num	Soil organic carbon density	250 m	Hengl et al. 2017	kg m ⁻³
soil_nitrogen	num	Soil nitrogen content	250 m	Hengl et al. 2017	g kg ⁻¹
teow_eco	chr	Dinerstein et al. 2017 - Terrestrial Ecoregion	NA	Dinerstein et al. 2017	NA

Table 11: Column names for plot spatial data table. *(continued)*

Column	Class	Description	Resolution	Citation	Units
teow_nnh	chr	Dinerstein et al. 2017 - Nature Needs Half classification	NA	Dinerstein et al. 2017	NA
teow_realm	chr	Dinerstein et al. 2017 - Terrestrial Realm	NA	Dinerstein et al. 2017	NA
teow_biome	chr	Dinerstein et al. 2017 - Terrestrial Biome	NA	Dinerstein et al. 2017	NA
whites_veg_major	chr	Whites vegetation map of Africa (1983) - major units	NA	Dinerstein et al. 2017	NA
whites_veg_minor	chr	Whites vegetation map of Africa (1983) - minor units	NA	Dinerstein et al. 2017	NA
frost_days	num	Mean annual number of frosty days (1990-2019)	0.5deg	Harris et al. 2020	NA
fire_count	int	MODIS burned area - Number of fires between 2001 and 2018	1000 m	Giglio et al. 2015	n
fire_intensity	num	MODIS active fires - mean fire intensity	1000 m	Giglio et al. 2016	NA
forest_height	num	Woody canopy height, estimated from GLAD GEDI Global Forest Height	30 m	Potapov et al. 2020	m
travel_time_city	num	Malaria Atlas Project, travel time to nearest city (>50,000 people)	30s	Nelson et al. 2019	min
travel_time_settlement	num	Malaria Atlas Project, travel time to nearest settlement (>5000 people)	30s	Nelson et al. 2019	min
distance_road	num	GRIP4 distance to nearest road of any size	NA	Meijer et al. 2018	m
type_nearest_road	chr	GRIP4 type of the nearest road	NA	Meijer et al. 2018	NA
pop_density	num	Worldpop Population density - people per km2	1000 m	Linard et al. 2012	n km ²

References

- Amatulli, Giuseppe et al. (2018). 'A suite of global, cross-scale topographic variables for environmental and biodiversity modeling'. In: *Scientific Data* 5.1. DOI: [10.1038/sdata.2018.40](https://doi.org/10.1038/sdata.2018.40).
- Chamshama, S. A. O., W. A. Mugasha and E. Zahabu (2004). 'Stand biomass and volume estimation for Miombo woodlands at Kitulungalo, Morogoro, Tanzania'. In: *The Southern African Forestry Journal* 200.1, pp. 59–70. DOI: [10.1080/20702620.2004.10431761](https://doi.org/10.1080/20702620.2004.10431761).
- Chave, J., C. Andalo et al. (2005). 'Tree allometry and improved estimation of carbon stocks and balance in tropical forests'. In: *Oecologia* 145.1, pp. 87–99. DOI: [10.1007/s00442-005-0100-x](https://doi.org/10.1007/s00442-005-0100-x).
- Chave, J., David Coomes et al. (2009). 'Towards a worldwide wood economics spectrum'. In: *Ecology Letters* 12.4, pp. 351–366. DOI: [10.1111/j.1461-0248.2009.01285.x](https://doi.org/10.1111/j.1461-0248.2009.01285.x).
- Chave, J., Maxime Réjou-Méchain et al. (2014). 'Improved allometric models to estimate the aboveground biomass of tropical trees'. In: *Global Change Biology* 20.10, pp. 3177–3190. DOI: [10.1111/gcb.12629](https://doi.org/10.1111/gcb.12629).
- Chidumayo, E.N. (1990). 'Above-ground woody biomass structure and productivity in a Zambezian woodland'. In: *Forest Ecology and Management* 36.1, pp. 33–46. DOI: [10.1016/0378-1127\(90\)90062-g](https://doi.org/10.1016/0378-1127(90)90062-g).
- Didan, K. and A. Barreto (2016). 'NASA MEaSUREs Vegetation Index and Phenology (VIP) Phenology NDVI Yearly Global 0.05 Deg CMG [Data set]'. In.
- Dinerstein, Eric et al. (2017). 'An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm'. In: *BioScience* 67.6, pp. 534–545. DOI: [10.1093/biosci/bix014](https://doi.org/10.1093/biosci/bix014).
- Fick, S. E. and R. J. Hijmans (2017). 'WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas'. In: *International Journal of Climatology* 37.12, pp. 4302–4315. DOI: <http://dx.doi.org/10.1002/joc.5086>.
- Giglio, L. et al. (2015). *MCD64A1 MODIS/Terra+Aqua Burned Area Monthly L3 Global 500m SIN Grid V006 [Data set]*. NASA EOSDIS Land Processes DAAC. [Accessed 2020-04-20].
- Giglio, Louis, C. Justice et al. (2015). 'MCD64A1 MODIS/Terra+Aqua Burned Area Monthly L3 Global 500m SIN Grid V006 [Data set]'. In: DOI: <https://doi.org/10.5067/MODIS/MCD64A1.006>.
- Giglio, Louis, Wilfrid Schroeder and Christopher O. Justice (2016). 'The collection 6 MODIS active fire detection algorithm and fire products'. In: *Remote Sensing of Environment* 178, pp. 31–41. DOI: [10.1016/j.rse.2016.02.054](https://doi.org/10.1016/j.rse.2016.02.054).
- Goodman, Rosa C. et al. (2013). 'Amazon palm biomass and allometry'. In: *Forest Ecology and Management* 310, pp. 994–1004. DOI: [10.1016/j.foreco.2013.09.045](https://doi.org/10.1016/j.foreco.2013.09.045).
- Harris, Ian et al. (2020). 'Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset'. In: *Scientific Data* 7.1. DOI: [10.1038/s41597-020-0453-3](https://doi.org/10.1038/s41597-020-0453-3).
- Hempson, Gareth P., Sally Archibald and William J. Bond (Dec. 2017). 'The consequences of replacing wildlife with livestock in Africa'. In: *Scientific Reports* 7.1. DOI: [10.1038/s41598-017-17348-4](https://doi.org/10.1038/s41598-017-17348-4).
- Hengl, Tomislav et al. (2017). 'SoilGrids250m: Global gridded soil information based on machine learning'. In: *PLOS ONE* 12.2. Ed. by Ben Bond-Lamberty, e0169748. DOI: [10.1371/journal.pone.0169748](https://doi.org/10.1371/journal.pone.0169748).
- Jarvis, A. et al. (2008). *Hole-filled SRTM for the globe : version 4 : data grid*.
- Kyalangalilwa, Bruce et al. (2013). 'Phylogenetic position and revised classification of *Acacia* s.l. (Fabaceae: Mimosoideae) in Africa, including new combinations in *Vachellia* and *Senegalia*'. In: *Botanical Journal of the Linnean Society* 172.4, pp. 500–523. DOI: [10.1111/boj.12047](https://doi.org/10.1111/boj.12047).
- Linard, Catherine et al. (2012). 'Population Distribution, Settlement Patterns and Accessibility across Africa in 2010'. In: *PLoS ONE* 7.2. Ed. by Guy J-P. Schumann, e31743. DOI: [10.1371/journal.pone.0031743](https://doi.org/10.1371/journal.pone.0031743).
- Meijer, Johan R et al. (2018). 'Global patterns of current and future road infrastructure'. In: *Environmental Research Letters* 13.6, p. 064006. DOI: [10.1088/1748-9326/aabd42](https://doi.org/10.1088/1748-9326/aabd42).
- Mugasha, W. A. et al. (2013). 'Allometric models for prediction of above- and belowground biomass of trees in the miombo woodlands of Tanzania'. In: *Forest Ecology and Management* 310, pp. 87–101. DOI: [10.1016/j.foreco.2013.08.003](https://doi.org/10.1016/j.foreco.2013.08.003).
- Mutakela, P. S. (2009). 'Biomass prediction models for *Colophospermum mopane* (Mopane) in Botswana'. MA thesis. Stellenbosch, South Africa: Stellenbosche University.
- Nelson, Andy et al. (2019). 'A suite of global accessibility indicators'. In: *Scientific Data* 6.1. DOI: [10.1038/s41597-019-0265-5](https://doi.org/10.1038/s41597-019-0265-5).
- Potapov, Peter et al. (2020). 'Mapping global forest canopy height through integration of GEDI and Landsat data'. In: *Remote Sensing of Environment*, p. 112165. DOI: [10.1016/j.rse.2020.112165](https://doi.org/10.1016/j.rse.2020.112165).

- Ryan, Casey M., Mathew Williams and John Grace (2011). ‘Above- and Belowground Carbon Stocks in a Miombo Woodland Landscape of Mozambique’. In: *Biotropica* 43.4, pp. 423–432. DOI: [10.1111/j.1744-7429.2010.00713.x](https://doi.org/10.1111/j.1744-7429.2010.00713.x).
- South, Andy (2017). *rnaturalearth: World Map Data from Natural Earth*. URL: <https://CRAN.R-project.org/package=rnaturalearth>.
- Tiepolo, Gilberto, Miguel Calmon and André Feretti (2002). ‘Measuring and Monitoring Carbon Stocks at the Guaraqueçaba Climate Action Project, Paraná, Brazil’. In: *Extension Serie Taiwan Forestry Research Institute* 153.
- Tietema, T. (1993). ‘Biomass determination of fuelwood trees and bushes of Botswana, Southern Africa’. In: *Forest Ecology and Management* 60.3-4, pp. 257–269. DOI: [10.1016/0378-1127\(93\)90083-y](https://doi.org/10.1016/0378-1127(93)90083-y).
- White, F. (1983). *The Vegetation of Africa: A descriptive memoir to accompany the UNESCO/AETFAT/UNSO vegetation map of Africa*. Paris, France: UNESCO. DOI: <http://dx.doi.org/10.2307/2260340>.
- Zanne, A. E. et al. (2009). *Global wood density database*. <http://hdl.handle.net/10255/dryad.235>. Dryad.