

PPOC pediatric EHR snapshot: an exploratory data analysis

A project-neutral reference for anyone analysing this extract. Every figure is measured from the delivered bundle; the report states what the data support, what they do not, and which checks this extract cannot answer at all.

0. How to use this report

Three ways in, depending on what you came for.

0.1 Three ways in

This report describes one snapshot of one pediatric primary-care EHR extract. It belongs to no project: it states what the data are, what they support, and what they cannot answer, and it leaves the research question to you.

Where to start

IF YOU ARE	START HERE
New to this extract	Read Part 1, then the not-applicable table in Part 2. Twenty minutes, and it will save days.
About to use a specific field	Find it in the Part 6 field index, then follow the finding it links to.
Explaining a number that looks wrong	Check the Part 7 artifact catalogue before assuming a bug in your code.
Planning a study	Part 1.4 first. The cohort selection invalidates several whole classes of question, and it is not visible in any field.

Every number here was measured from the delivered bundle for snapshot `2026-08-24` ; none is copied from another document without being recomputed. The cohort is pinned to 31 Dec 2024 and the extract was cut on 03 Feb 2025.

What this report is not. It is not a clinical validation, not a registered analysis, and not a statement about any individual child. Every figure is an aggregate, and any cell resting on fewer than 10 records is suppressed.

1. The snapshot

What this extract contains, how it was built, and what its construction forecloses.

1.1 Package identity and integrity

Everything in this report was computed from the typed DuckDB bundle of package `ppoc-pediatric-ehr` 1.0.0, snapshot `2026-08-24`, sha256 `425c6f873cefcd149344570561a03b33c69a6a6af7fa18bc777c0429579507116`. The bundle is opened read-only and is never copied into this repository.

Three independent sources state how large this extract should be: the bundle manifest, the PPOC delivery documents committed under `docs/`, and the data itself. They are reconciled here before any other figure is computed, so that a bundle drawn from a different extract would be visible rather than silently profiled.

Row counts, measured against both declared sources

RESOURCE	MEASURED	BUNDLE MANIFEST	PPOC DOCUMENT	AGREES
patients	250,588	250,588	250,588	yes
patients_augmented	250,588	250,588	—	yes
visits	6,494,473	6,494,473	6,494,473	yes
visits_augmented	6,494,473	6,494,473	—	yes
labs	17,230,681	17,230,681	17,230,681	yes
medications	3,823,049	3,823,049	3,823,049	yes
problem_list	1,709,584	1,709,584	1,709,584	yes
referrals	349,827	349,827	349,827	yes

A dash in the PPOC column means the resource was not part of the delivery: `patients_augmented` and `visits_augmented` are generated locally from the delivered files, so PPOC states no count for them. See 1.3.

Distinct patients per resource, against the PPOC counts

RESOURCE	MEASURED	PPOC DOCUMENT	AGREES
visits	250,588	250,588	yes
problem_list	238,823	238,823	yes
labs	247,271	247,271	yes
medications	236,323	236,323	yes
referrals	138,071	138,071	yes

The lab resource carries a second stated figure: 6,578,838 distinct lab orders behind the resulted components. The bundle holds 6,578,838. Across every count above, 0 disagree.

1.2 Resource map, grain, and keys

The package is 8 tables carrying 254 columns between them, but they do not share a provenance: 6 were delivered by PPOC and 2 are generated locally (1.3). Grain matters more than row count here: three of the resources are keyed on something other than the patient or the visit, and one of them needs two columns to be unique.

The eight resources

RESOURCE	SOURCE	ROWS	COLS	GRAIN	PRIMARY KEY	LINKS TO
patients	PPOC	250,588	11	one row per patient	patient_id	—
patients_augmented	scripts/augment.py	250,588	87	one row per patient	patient_id	patients
visits	PPOC	6,494,473	43	one row per patient per encounter	visit_id	patients
visits_augmented	scripts/augment.py	6,494,473	82	one row per patient per encounter	visit_id	visits
labs	PPOC	17,230,681	12	one row per resulted component of a lab order	lab_order_id + result_line_num	patients; visits (partial)
medications	PPOC	3,823,049	8	one row per medication order or historical record	med_record_id	patients; visits (partial)
problem_list	PPOC	1,709,584	5	one row per problem-list entry	problem_list_id	patients
referrals	PPOC	349,827	6	one row per referral order	referral_id	patients; visits (partial)

Six resources were delivered by PPOC; the two augmented ones are generated locally from them. 1.3 explains why that distinction matters.

`visit_id` on labs, medications, and referrals is a partial link by design, not a defect: an order placed outside a visit carries an identifier that resolves to no encounter in this extract. Section 3.2 measures how partial.

1.3 Two layers with different provenance

Only 6 of the 8 resources in this package came from PPOC. The delivery comprised patients, visits, problem list, medications, labs, and referral orders; the data dictionary and the extract diagram committed under `docs/` describe those 6 and no others. The remaining 2 — `patients_augmented` and `visits_augmented` — are **generated locally** by `scripts/augment.py` from the delivered files, using CDC LMS reference tables, velocity rules, and outlier detection.

That distinction decides who can fix what. A defect in a delivered resource is the source system's and can only be worked around; a defect in the augmented layer belongs to a script in this repository and can be corrected by re-

running it. Everything this report labels a *derivation* artifact — the truncated height z-score of 4.6, the double-converted head circumference of 4.7, the interval rule behind the velocity fields of 4.8 — is a property of that local step, not of the data PPOC sent.

Because the augmented layer is derived from the delivered one, the fields they share should agree exactly. Across all 6,494,473 joined visit rows, five of the six do.

Shared visit fields, raw against augmented


FIELD	ROWS DIFFERING	SHARE
height_in	0	0.00%
weight_oz	0	0.00%
head_circ_cm	0	0.00%
encounter_type	0	0.00%
age_in_days	0	0.00%
BMI / bmi	3,658,277	56.33%

BMI is the exception, and the disagreement is structured rather than noisy. 1,703,005 visits carry a raw BMI where the augmented bmi is null, at a median age of 0.51 years; the augmented layer withholds BMI below age 2, where a CDC BMI-for-age reference does not apply, while the raw value is computed inside the source EHR at every age. A further 41 rows go the other way, and 536 carry both values differing by more than 0.01.

Implications for analysis. Reading visits.BMI silently yields infant BMI values that the augmented layer deliberately suppresses, and the two layers will not reproduce each other's descriptive statistics. Choose a layer for a stated reason and record which; do not mix them within one analysis. The 536 rows where both are present and disagree are small enough to screen individually.

1.4 How this cohort was built

This is the most consequential section of the report, because it describes a property of the data that no field exposes and that no amount of analysis can recover. The 250,588 patients here are what remains after four successive exclusions applied by PPOC to an active-patient registry of 437,996. The final count reconciles against the delivered data exactly: 250,588 patient rows.



Cohort construction, registry to delivered extract

The four exclusions

STEP	CRITERION	EXCLUDED	REMAINING
0	On the PPOC active-patient registry		437,996
1	Age under 18 as of 31 Dec 2024	76,670 excluded	361,326
2	Excluding 2 practices that declined participation	9,309 excluded	352,017
3	At least 5 growth measurements of one type on distinct dates, spanning over 1095 days, last measurement within 400 days	61,842 excluded	290,175
4	Carrying no rare diagnosis, medication, or lab	39,587 excluded	250,588

"Active" on that registry means living status alive, not flagged as a test or inactive record, an active PPOC primary-care association, and either a visit in the last three years or one scheduled in the next fifteen months. The cohort is pinned to 31 Dec 2024 and the extract was cut on 03 Feb 2025.

The fourth exclusion is the one most likely to be missed, because it removed *patients* rather than codes. A diagnosis, medication, or lab occurring fewer than 11 times in the data set was classed rare, and every patient carrying one was dropped.

What the rarity exclusion removed

VOCABULARY	DISTINCT VALUES	CLASSSED RARE	SHARE
ICD-10 diagnosis codes	30,493	18,604	61%
Simple generic medications	2,503	1,391	56%
Lab procedures	13,402	9,621	72%

Implications for analysis. Rare conditions, rare exposures and uncommon labs are absent by construction, not merely sparse: a study of any of them returns a confident low rate rather than an obviously missing population. 61% of diagnosis codes, 56% of medications and 72% of lab procedures left with their patients. Because the registry requires living status alive, there are no deceased patients and mortality is not an available outcome. Because entry required at least five growth measurements, trajectory richness is an entry criterion and not a finding about pediatric care. And because the last measurement had to fall within 400 days of the cohort date, the panel is right-censored by design. No frequency in this extract is a population prevalence.

Two ambiguities in the source documents are recorded rather than silently resolved. The cohort workbook describes the under-three exemption as applying to the span requirement for children who already have five measurements, while the extract diagram describes it as age under three with at least one measurement. The same two documents give the rarity threshold as "fewer than 11 occurrences" and "under 10 patients".

1.5 The de-identification envelope

`age_in_days` is the only clock. The extract carries no calendar date, no time of day, no site, practice, provider, or geography, and no free text from any note. That is stated once here and referenced from Part 2 rather than re-argued at each check it rules out.

Checks this extract forecloses, and why

STANDARD CHECK	WHY IT CANNOT BE RUN
Duplicate-patient detection	no name, birth date, or linkage key survives
Batch-entry clustering	ages are integer days; there is no time of day
System downtime gaps	no calendar axis on which a void could appear
Missingness by site or provider	no such column exists in any resource
Calendar trend breaks and policy shifts	no calendar axis
Copy-forward of note text	no note text is included
Documentation timing	no timestamps

One qualification, because "no calendar axis" is easy to overstate: the cohort itself is pinned to 31 Dec 2024 and the extract was cut shortly after. Ages are relative to each child's birth, but the *window* is fixed and known, which is what makes the recency criterion in 1.4 a right-censoring rule rather than an unknown.

2. Checklist coverage

Every item of the general EHR EDA checklist, mapped to what this snapshot can and cannot support.

2.1 The checklist, item by item

This part exists so that nobody has to wonder whether a standard check was skipped or was impossible. Of 44 items in the general EHR exploratory-analysis checklist, 31 are covered here, 4 are partially covered, and 9 cannot be run against this extract at all.

Checklist coverage

CHECKLIST SECTION	ITEM	STATUS	WHERE, OR WHY NOT
0 Provenance	Extraction window	covered	Cohort and extract dates recovered from the delivery documents — 1.4
0 Provenance	Inclusion/exclusion logic	covered	The full four-step funnel — 1.4
0 Provenance	Vendor, version, migration events	covered	Epic against converted legacy records — 3.7
0 Provenance	Data dictionary present	covered	Committed under docs/, reconciled field by field — 1.1
0 Provenance	Raw vs CDM vs custom extract	covered	A custom extract plus a derived augmentation layer — 1.3
1 Structural	Row and table counts	covered	Against the manifest and the vendor's own counts — 1.1
1 Structural	Primary key uniqueness	covered	All eight resources — 3.1
1 Structural	Referential integrity	covered	3.2
1 Structural	Duplicate patient detection	not applicable	No name, birth date, or linkage key survives de-identification — 1.5
1 Structural	Schema drift	covered	Live schema against the dictionary; three documented fields absent — 1.1
1 Structural	Grain per table	covered	Including that patient and age is not unique in visits — 3.1
2 Temporal	Timestamp semantics	covered	3.3
2 Temporal	Impossible sequences	covered	3.3
2 Temporal	Batch-entry clustering	not applicable	Ages are integer days; there is no time of day — 1.5
2 Temporal	System downtime gaps	not applicable	No calendar axis — 1.5
2 Temporal	Coding or vendor transition	partial	Epic against converted is computable; ICD-9 to ICD-10 is not, without dates
2 Temporal	Age sanity	covered	3.3
3 Missingness	Missingness per field	covered	3.4 and the field index

CHECKLIST SECTION	ITEM	STATUS	WHERE, OR WHY NOT
3 Missingness	Missingness pattern	covered	By age, sex, and encounter — 3.4
3 Missingness	Sentinel values	covered	3.5
3 Missingness	Not measured vs measured negative	covered	Two fields whose nulls carry meaning — 3.5
3 Missingness	Missingness by site or provider	not applicable	No site, department, or provider column exists — 1.5
4 Distributional	Univariate distributions	covered	4.3
4 Distributional	Unit inconsistencies	covered	4.3 and 4.4
4 Distributional	Digit preference and rounding	covered	4.2
4 Distributional	Categorical value counts	covered	3.6
4 Distributional	Outlier detection	covered	Bounds reported before any exclusion is recommended — 4.3
4 Distributional	Cross-field plausibility	covered	Raw against augmented layers — 1.3
5 Terminology	Code system vintage	covered	3.6
5 Terminology	Granularity consistency	covered	3.6
5 Terminology	Problem list staleness	covered	3.5
5 Terminology	Free text vs structured	covered	Laboratory result values are semi-structured text — 3.6
5 Terminology	Local or custom codes	covered	3.6
6 Workflow	Copy-forward detection	partial	Detectable on measurements; no note text is included
6 Workflow	Template or boilerplate detection	not applicable	No note text — 1.5
6 Workflow	Documentation timing	not applicable	No timestamps — 1.5
6 Workflow	Order/result reconciliation	covered	3.6
7 Population	Cohort representativeness	partial	The cohort is not representative and 1.4 says exactly how; no external benchmark ships with this repository
7 Population	Encounter type mix	covered	3.7
7 Population	Follow-up time distribution	covered	4.1
7 Population	Site or provider volume	not applicable	No such field — 1.5
8 Longitudinal	Calendar trend breaks	not applicable	No calendar axis. Age-axis profiles are reported instead and are not the same thing — 1.5
8 Longitudinal	Guideline or policy shift	not applicable	Requires calendar time — 1.5
8 Longitudinal	Vendor changeover effects	partial	The Epic against converted contrast only

The not-applicable list is the part worth reading before you start. Every entry is a consequence of de-identification or of what the extract simply does not carry, and no amount of analysis recovers any of them.

Checks this extract cannot support

CHECK	WHY
Duplicate patient detection	No name, birth date, or linkage key survives de-identification — 1.5
Batch-entry clustering	Ages are integer days; there is no time of day — 1.5
System downtime gaps	No calendar axis — 1.5
Missingness by site or provider	No site, department, or provider column exists — 1.5
Template or boilerplate detection	No note text — 1.5
Documentation timing	No timestamps — 1.5
Site or provider volume	No such field — 1.5
Calendar trend breaks	No calendar axis. Age-axis profiles are reported instead and are not the same thing — 1.5
Guideline or policy shift	Requires calendar time — 1.5

Implications for analysis. Treat the second table as a design constraint rather than a gap to work around. A protocol that depends on provider variation, time-of-day effects, calendar trends, or deceased patients cannot be run on this extract, and discovering that after cohort construction is expensive.

3. Integrity

Keys, linkage, the age axis, missingness, terminology, and capture.

3.1 Keys, grain, and uniqueness

Every declared primary key holds. The labs resource needs all three of its declared columns to be unique, which is worth stating because joining on order and component alone will multiply rows.

Declared keys, measured

RESOURCE	KEY	ROWS	DISTINCT KEYS	UNIQUE
patients	patient_id	250,588	250,588	yes
patients_augmented	patient_id	250,588	250,588	yes
visits	visit_id	6,494,473	6,494,473	yes
visits_augmented	visit_id	6,494,473	6,494,473	yes
medications	med_record_id	3,823,049	3,823,049	yes
problem_list	problem_list_id	1,709,584	1,709,584	yes
referrals	referral_id	349,827	349,827	yes
labs	lab_order_id + component + line	17,230,681	17,230,681	yes

What is *not* a key is the combination a longitudinal analysis reaches for first. 5,478 patient-days (0.08% of 6,488,911) carry more than one visit, covering 11,040 visit rows (0.17% of all visits). `age_in_days` is therefore not unique within a patient.

Implications for analysis. Any trajectory ordered by age alone has ties, and any window function partitioned by patient and ordered by age will resolve them arbitrarily unless you say how. Decide whether to take the first row, the mean, or the non-null value, and apply it before the analysis rather than inside it.

3.2 Referential integrity and cross-resource linkage

`patient_id` resolves everywhere. `visit_id` does not, and the shortfall is large enough that treating it as a complete foreign key will quietly drop or duplicate rows.

Visit linkage by resource

RESOURCE	ROWS	VISIT_ID NULL	POPULATED BUT UNRESOLVED	SHARE OF POPULATED	UNRESOLVED PATIENT_ID
labs	17,230,681	0.00%	5,201,657	30.19%	0

RESOURCE	ROWS	VISIT_ID NULL	POPULATED BUT UNRESOLVED	SHARE OF POPULATED	UNRESOLVED PATIENT_ID
medications	3,823,049	0.00%	1,592,437	41.65%	0
referrals	349,827	7.10%	98,623	30.35%	0

This is documented behaviour rather than corruption. The data dictionary states for each of these resources that the visit link "may not match to all" when the order was placed or the record documented outside a visit. The trap is that the column is populated on nearly every row, so a required-looking key silently fails to join.

Implications for analysis. Join to visits with an explicit outer join and count what fails, rather than an inner join that hides the loss. Anything computed per visit — encounter type, visit-level anthropometrics — is unavailable for the unresolved share, and that share is not random: it concentrates in orders placed outside encounters.

3.3 Age-axis consistency and impossible sequences

Age in days is the only clock, so ordering violations within a resource are visible directly. Counts below 10 are suppressed.

Ordering and range checks

CHECK	VIOLATING ROWS	ROWS CHECKED	SHARE
Lab result age earlier than lab order age	583,055	14,947,495	3.901%
Medication start age earlier than order age	329,107	3,539,983	9.297%
Medication end age earlier than start age	12,709	3,179,759	0.400%
Problem resolved age earlier than noted age	0	754,996	0.000%
Problem noted before birth	650	1,702,300	0.038%
Lab ordered before birth	47	17,230,681	0.000%
Medication ordered before birth	—	3,823,049	—
Visit recorded before birth	0	6,494,473	0.000%

An em dash in the violating-rows column means the count is nonzero but below the suppression threshold.

The lab and medication violations are the substantial ones, and both are documented at source. For a historically documented medication the order date is the date the record was *written*, not when the drug was started, and a charted approximation such as a month with no day is stored as the first of that month. End dates may sit in the future while a medication is active. Lab result and order ages derive from different source timestamps.

Implications for analysis. Differences between two age fields in these resources are not reliable durations. Where you need an interval, take it from a single field across rows rather than between two fields on one row, and exclude historically documented medication records from any start-to-end calculation.


3.4 Missingness, by field and by age

Population was measured for all 176 columns in the extract, counting the repeated diagnosis and race families once each. 0 columns are entirely empty. The full table is Part 6; the sixteen least-populated columns are below.

The least-populated columns

RESOURCE	FIELD	POPULATED ROWS	MISSING
patients_augmented	dx_age_years_e24	1	100.0%
patients_augmented	dx_age_years_e72_11	1	100.0%
patients_augmented	dx_age_years_n25_0	1	100.0%
patients_augmented	dx_age_years_q78_1	2	100.0%
patients_augmented	dx_age_years_e22_0	3	100.0%
patients_augmented	dx_age_years_q78_0	10	100.0%
patients_augmented	dx_age_years_q77	15	100.0%
patients_augmented	dx_age_years_q87_4	17	100.0%
patients_augmented	dx_age_years_q98_5	17	100.0%
patients_augmented	dx_age_years_q98_0	26	100.0%
patients_augmented	dx_age_years_e23_6	31	100.0%
patients_augmented	dx_age_years_q87_2	32	100.0%
patients_augmented	dx_age_years_q96	36	100.0%
patients_augmented	dx_age_years_q98_4	42	100.0%
patients_augmented	dx_age_years_q87_3	46	100.0%
patients_augmented	dx_age_years_p04_3	53	100.0%

A single missingness rate hides the thing that matters most for a longitudinal extract: whether a field is missing *at random* or missing *by age*. For the measurement channels it is emphatically the latter.



Share of visits carrying each measurement, by age band

Implications for analysis. Head circumference is an infant measurement and effectively disappears after age 2; BMI and its percentile are withheld below age 2 by the augmentation; height is recorded far less often than weight at every age. Any cohort defined by "has a complete measurement row" is therefore an age-selected cohort, and any model that drops incomplete rows inherits that selection. Report availability by age band before interpreting any age-stratified contrast.

3.5 Nulls that are not missing, and sentinels that are not data

Two of the largest null populations in this extract are not missing data at all, and reading them as missing throws away the majority of the signal in their columns.

Laboratory result flags

RESULT_FLAG	ROWS	MEANING
null	15,550,985	normal result
Abnormal	704,327	abnormal
High	513,650	abnormal
Low	361,300	abnormal
Sensitive	62,794	abnormal
Resistant	10,278	abnormal
High Panic	9,744	abnormal
(NONE)	5,881	abnormal
Normal	4,273	abnormal
Panic	3,056	abnormal
Intermediate	1,704	abnormal
Low Panic	1,406	abnormal
Critical	373	abnormal
Negative	188	abnormal
High Off-Scale	134	abnormal
Susceptible-Dose Dependent	123	abnormal
Abnormal High	99	abnormal
Abnormal Low	92	abnormal
Invalid High	84	abnormal
Sig Change Up	68	abnormal
Positive	35	abnormal
Critical High	23	abnormal
Low Off-Scale	17	abnormal
Critical Low	13	abnormal
Class 0: Absent Allergen Specific IgE	—	abnormal

RESULT_FLAG	ROWS	MEANING
Delta Abnormal High	—	abnormal
Invalid Low	—	abnormal
Better	—	abnormal
Class 2: Moderate Level Allergen Specific IgE	—	abnormal
Delta Critical High	—	abnormal
In Process	—	abnormal
Class 3: High Level Allergen Specific IgE	—	abnormal
Sig Change Down	—	abnormal
Delta Abnormal Low	—	abnormal
Moderately Sensitive	—	abnormal
Worse	—	abnormal

All 36 distinct values are listed.

The data dictionary defines `result_flag` as an HL7 abnormality category in which the value `(NONE)` means a normal result and anything else means abnormal. This extract contains 5,881 literal `(NONE)` values and 15,550,985 nulls — 90.3% of all lab rows. The sentinel became a null somewhere between the source system and delivery, so **a null flag means normal, not unknown**.

`problem_list.resolved_date_age_in_days` behaves the same way: the dictionary defines null as "problem currently active". 951,677 of 1,709,584 entries (55.7%) are null, which is a statement about 56% of problems being open, not about missing dates.

Zero and blank values checked as possible sentinels

RESOURCE	FIELD	PATTERN	ROWS
visits_augmented	height_in	zero height	0
visits_augmented	weight_oz	zero weight	0
visits_augmented	head_circ_cm	zero head circumference	—
labs	result_value	empty result string	0
patients	sex	blank sex	0
patients	race_1	blank race_1	8,818
patients	ethnicity	blank ethnicity	5,464

Implications for analysis. Never impute or drop on `result_flag` or `resolved_date_age_in_days` nullity. An abnormal-result rate computed as "non-null flags over non-null flags" will read as 100%; the correct denominator is all resulted rows. A problem-list resolution rate must count nulls as unresolved rather than excluding them.

3.6 Code systems, free text, and categorical hygiene

Diagnosis coding is almost entirely well-formed ICD-10. Of 14,714,503 filled encounter-diagnosis slots across 8,029 distinct codes, 145,992 (0.99%) do not match the ICD-10 shape; of 1,709,584 problem-list entries across 4,739 codes, 39,860 (2.33%).

The non-conforming diagnosis values

VALUE	SLOTS
IMO0002	116,950
U07.1	25,483
IMO0001	3,443
U09.9	77
U07.0	39

These are proprietary placeholders the source EHR emits when a clinical term has no ICD-10 equivalent. They carry no diagnostic meaning and should be excluded from code-based cohort definitions rather than treated as unmapped diagnoses.

Laboratory results are the opposite case. `result_value` is a text column: of 17,230,681 rows, 7,621,449 (44.2%) parse as a number and 2,494,261 (14.5%) are empty. Among the rest, 487,168 are censored results carrying a comparator prefix, and the remainder are qualitative results, specimen descriptors, and administrative non-results. A LOINC code is present on only 7.8% of rows.

The declared key holds, but 33,879 order-and-component pairs appear on more than one result line and 23,679 of those (69.9%) carry disagreeing values. The data dictionary records the cause: a result may fail to link back to its original order, which duplicates the record.

Categorical vocabularies before and after normalising case and internal whitespace

RESOURCE	FIELD	DISTINCT VALUES	AFTER NORMALISING	COLLAPSED
labs	lab_procedure_name	3,742	3,739	3
medications	med_simple_generic_name	1,073	1,073	0
referrals	requested_specialty	119	119	0

Implications for analysis. A naive numeric cast on `result_value` silently discards more than half the populated values and turns a left-censored result into a missing one rather than a bound. Join labs on order, component *and* line number, or the duplicate lines will multiply rows and pick a value arbitrarily. The categorical vocabularies barely collapse under normalisation, so grouping by them is safe after trimming.

3.7 Capture: measurement presence is not measurement occurrence

Completeness by age says how often a column is filled. Encounter type says whether filling it could have meant a measurement.

Measurement and diagnosis presence by encounter type

ENCOUNTER TYPE	VISITS	WEIGHT PRESENT	HEIGHT PRESENT	FIRST DIAGNOSIS
Office Visit	4,725,643	99.9%	52.4%	95.8%
Well Visit (Conv.)	778,452	99.9%	98.7%	94.4%
Sick	580,991	99.9%	16.9%	95.4%
Follow-Up	92,370	99.8%	24.9%	93.1%
Walk-In	79,679	99.9%	8.7%	96.9%
Consult	32,355	99.9%	53.7%	99.6%
Conversion Encounter	32,007	99.9%	66.9%	17.3%
Newborn	31,142	99.9%	87.5%	99.1%
Telemedicine	25,658	97.4%	44.8%	99.8%
Telephone	22,053	99.3%	50.1%	34.8%
Weight Check	16,295	99.9%	37.7%	96.4%
Clinical Support	15,347	99.3%	15.4%	83.4%
Documentation	13,107	98.3%	84.4%	8.6%
Immunization	11,774	97.6%	30.7%	93.4%
New Patient	11,410	99.9%	79.8%	98.1%
Nutrition	5,406	99.8%	73.8%	99.5%
Medication Management	4,345	99.9%	75.9%	93.3%
Nurse Only	3,786	96.9%	36.2%	69.1%
Abstract	3,672	99.4%	83.1%	9.0%
Flu	2,104	96.4%	13.7%	99.4%
Lactation Consult	1,384	100.0%	11.6%	99.7%
Lab	1,351	97.0%	37.3%	78.7%
Lactation Encounter	1,337	99.8%	1.9%	49.7%
Procedure visit	678	97.8%	56.3%	96.0%
Pre-op/Pre-procedure Orders	575	100.0%	84.3%	96.3%
Erroneous Encounter	555	99.8%	45.9%	30.5%
Orders Only	273	99.6%	78.4%	51.6%
External Contact	211	100.0%	99.5%	2.4%
Patient Message	209	95.7%	30.6%	18.2%
Evaluation	143	99.3%	74.1%	94.4%
Lab Requisition	63	96.8%	14.3%	90.5%
Scanned Document	23	100.0%	95.7%	0.0%
Letter (Out)	18	100.0%	94.4%	0.0%
Refill	12	100.0%	0.0%	66.7%
History	11	100.0%	45.5%	0.0%
Ophth Exam	11	100.0%	0.0%	100.0%
Hospital	—	—	—	—

ENCOUNTER TYPE	VISITS	WEIGHT PRESENT	HEIGHT PRESENT	FIRST DIAGNOSIS
Routine Prenatal	—	—	—	—
Transcribe Orders	—	—	—	—
Patient Care Review	—	—	—	—
Episode Changes	—	—	—	—
Erroneous Telephone Encounter	—	—	—	—
ED	—	—	—	—
OurPractice Advisory	—	—	—	—
Treatment	—	—	—	—

All 45 distinct values are listed. 9 carry too few visits to show a count.

Telephone encounters carry a weight on 99.3% of 22,053 visits. A weight cannot be measured over the telephone, so those values were produced some other way — reported by a caregiver, carried from a nearby in-person encounter, or attached to an encounter whose type label does not describe how the patient was seen. Which of those it is cannot be determined from this extract.

Recording completeness by source system

ENCOUNTER SOURCE	VISITS	HEIGHT PRESENT	FIRST DIAGNOSIS
Epic	4,149,865	51.4%	99.7%
converted from a legacy system	2,344,608	58.7%	86.1%

The source-system split is the migration signal. Records converted from the practice network's previous EHR carry a first diagnosis on only 86.1% of encounters, which the data dictionary anticipates: converted encounters may be missing diagnosis information depending on the quality of the conversion.

Implications for analysis. A visit-level indicator that a measurement is present is not evidence that a measurement was taken at that encounter. If your design counts measurement occasions — visit density, monitoring intensity, follow-up adherence — restrict to encounter types where physical measurement is possible rather than relying on presence. And any diagnosis-based rate computed across the whole extract mixes two populations with very different coding completeness.

3.8 Same-day measurements that disagree

Section 3.1 shows that a patient-day can carry more than one visit. Where those visits each carry the same measurement, they often do not agree, and the size of the disagreement is a direct estimate of how far two records of the same child on the same day can sit apart.

Patient-days carrying more than one value of a channel

CHANNEL	PATIENT-DAYS WITH 2 OR MORE	OF WHICH THEY DISAGREE	SHARE DISAGREEING	MEDIAN SPREAD	95TH PERCENTILE	MAXIMUM
---------	-----------------------------------	------------------------------	----------------------	------------------	--------------------	---------

CHANNEL	PATIENT-DAYS WITH 2 OR MORE	OF WHICH THEY DISAGREE	SHARE DISAGREEING	MEDIAN SPREAD	95TH PERCENTILE	MAXIMUM
height	2,958	942	31.8%	3.175	12.16	34.92
weight	5,319	2,648	49.8%	0.118	3.64	36.57
head circumference	1,475	837	56.7%	0.510	27.80	397.76

Spread columns are in each channel's own unit and describe only the disagreeing days, not the panel.

The height spread is the notable one. A median disagreement of 3.17 cm between two heights recorded for the same child on the same day is far larger in relative terms than the weight equivalent, and it is the size of difference expected when recumbent length and standing height are mixed, or when one value is carried from an earlier note. Section 4.5 finds the same effect across the length-to-height transition age, seen there across months rather than within a day.

Implications for analysis. These days need a tie rule chosen before the analysis, not left to whatever order the query returns. Taking the minimum, the maximum, the mean, or the first row are all defensible and they give different answers; what is not defensible is not knowing which one you took. Deduplicate the patient-day before any window function, since 4.8 shows the derivation layer's own ambiguity on exactly these rows.

3.9 Counting diagnosis codes: ICD-10 is a hierarchy

ICD-10 is a tree, not a list. `E10` is type 1 diabetes and `E10.9` is type 1 diabetes without complications; a chart may carry either, and which one it carries is a coding decision rather than a clinical one. **A query that matches a code exactly therefore counts one node of the tree, not the concept.** This is the single most common way to undercount a diagnosis in this extract, and it fails silently — the query returns a number, just the wrong one.

The extract carries 8,965 distinct codes across its two diagnosis resources, of which only 123 are bare three-character categories. Rolling every code up to its category gives 1,327 categories, and **1,204 of those (90.7%) never appear as a bare code at all.** For those, an exact-match query returns zero while the condition is present.

The effect is large enough to reorder a frequency table. Below, the six most common literal codes beside the six most common categories after rollout.

The most common diagnoses, counted flat and rolled up

RANK	LITERAL CODE	PATIENTS	CATEGORY	PATIENTS
1	Z00.129	247,963	Z00	250,354
2	Z23	244,617	Z23	244,617
3	Z13.0	135,783	Z13	188,758
4	J06.9	134,337	J06	134,368
5	Z13.88	128,880	H66	132,693
6	R50.9	116,527	J02	131,194

H66 is the clearest case. Counted literally it has 0 patients, because clinicians code the laterality-specific children instead. Counted as a subtree it has 132,693 — enough to place it among the most common conditions in the extract, where a flat count makes it invisible.

The children a flat count misses

CODE	PATIENTS
H66.001	49,713
H66.002	44,040
H66.003	35,569
H66.90	34,331
H66.91	26,536


Implications for analysis. Match on a prefix (`code LIKE 'E10%'`) or roll up to the level you actually mean before counting, and say which level that is. Two cautions on prefixes: a code is a string, so compare against the code with its decimal point as stored, and a prefix of a prefix will over-match — `E1` is not a category. Where a frequency table is the deliverable rather than an input, report the rolled-up count and the literal one side by side, since the gap between them is itself a description of local coding practice.

4. Anthropometrics

The richest and most artifact-prone measurements in the extract.

4.1 Trajectory supply: how many heights each child has

250,267 of 250,588 patients (99.9%) carry at least one derived height, 235,651 (94.0%) carry five or more, and 182,037 (72.6%) carry ten or more.



Patients retaining at least k height observations

Height observations per patient


AT LEAST K HEIGHTS	PATIENTS	SHARE OF COHORT
1	250,267	99.9%
3	245,449	97.9%
5	235,651	94.0%
10	182,037	72.6%
15	104,838	41.8%
20	43,501	17.4%
25	16,040	6.4%

Implications for analysis. Read this against 1.4 before treating it as a fact about pediatric care. Cohort entry required at least five growth measurements of *some* type, so a dense height series here is partly the selection rule and partly the underlying practice; the two cannot be separated within this extract. What the curve does support is a feasibility estimate: how many children remain if your design needs k observations.

4.2 Recording units and the measurement grid

Height and weight are captured in imperial units, and the metric columns are exact conversions of them — measured, not assumed. Across 3,491,662 visits carrying both a raw and a derived height, 0 disagree with `height_in` times 2.54 by more than 0.01 cm; across 6,483,007 weight pairs, 0 disagree with `weight_oz` times 0.0283495. The arithmetic is clean, which matters because a value keyed in the wrong unit survives an exact conversion unchanged — 4.4 takes that up.

The recorded values are heaped on human-readable fractions: of 3,509,633 heights, 31.0% fall on a whole inch, 54.9% on a half inch and 80.0% on a quarter inch. Of 6,488,028 weights, 54.4% fall on a whole ounce and 24.6% on a whole pound.



Share of measurements falling on the coarse grid, by age

The two channels age in opposite directions. Height stays on its quarter-inch grid throughout childhood, while weight moves from ounce-level precision in infancy to whole pounds in adolescence, so the effective resolution of the weight channel degrades as children get older.


Implications for analysis. One quarter inch is 0.635 cm, and the derived `height_cm` carries two decimals it has not earned. Any change smaller than roughly half the rounding interval is not distinguishable from the rounding itself, which sets a floor on the smallest trajectory deflection that can be detected at all. State the assumed precision wherever a measurement is written out, and set detection thresholds at or above the grid.

4.3 Distributions and plausibility bounds


The four measurement channels, summarised on the derived metric columns. The final column counts values outside a conventional review range; those are reported, not removed, because the decision to exclude belongs to the analysis rather than to this report.

Measurement channels


CHANNEL	UNIT	VALUES	MIN	1ST PCT	MEDIAN	99TH PCT	MAX	OUTSIDE REVIEW RANGE
height	cm	3,491,662	37.47	48.26	92.71	173.41	196.60	0
weight	kg	6,483,007	0.01	2.81	14.52	78.65	513.92	217
BMI	kg/m^2	1,955,339	8.12	13.29	16.70	32.42	219.51	44
head circumference	cm	1,635,690	0.00	33.00	44.00	53.00	505.46	13,742




Distribution of height (cm)



Distribution of weight (kg)



Distribution of BMI (kg/m^2)



Distribution of head circumference (cm)

Implications for analysis. Head circumference is the channel whose tails are worst, and 4.4 shows why. For the others the extremes are sparse but the bulk is clinically ordinary. Bound the raw imperial columns rather than the derived metric ones when screening, since a wrong unit survives an exact conversion unchanged.

4.4 Transcription-error signatures in the typed fields


Method. Each measurement is anchored by linear interpolation between the same child's previous and next measurement. Both neighbours must themselves be plausible and span no more than four years, so a bad neighbour cannot manufacture an anomaly. A height is anomalous more than 3 inches from that anchor, a weight more than 50% from it. A mechanism *reconciles* an anomaly when applying it to the recorded value lands back at the anchor.

The null. Each anomaly's anchor is replaced by the recorded value plus a deviation drawn from another anomaly in the same year-of-age band, 20 times. That preserves the distribution of deviations exactly and destroys only the arithmetic relationship between the recorded digits and the anchor, which is the thing under test. A mechanism that reconciles anomalies no more often than it reconciles these scrambled pairs has no evidence behind it, however many hits it returns.

Height. 7,443 anomalies in the testable interior. Mechanisms are tested one at a time and are not mutually exclusive, so the rows do not sum to the total.

Height: mechanisms against the null

MECHANISM	RECONCILED	SHARE	NULL	RATIO
height recorded in whole feet	686	9.22%	1.34%	6.9x
centimetre value in the inch field	91	1.22%	0.67%	1.8x
inch value where a centimetre is expected	30	0.40%	1.00%	0.4x
decimal point misplaced	11	0.15%	2.80%	0.1x
adjacent digit transposition	40	0.54%	0.69%	0.8x
one digit omitted	407	5.47%	1.14%	4.8x
one digit wrong (calibration class)	4,024	54.06%	50.15%	1.1x



Height: observed against null, by mechanism

Adjacent digit transposition — the classic keying error, and the one most often assumed — reconciles fewer height anomalies than chance alone. The unit error is real and it is directional: a centimetre value in the inch field is enriched, while the arithmetically opposite reading sits at or below the null. That asymmetry is what a one-way data-entry confusion looks like; a spurious mechanism would be symmetric.

The dropped-digit row does not survive inspection, and it is worth showing why. Inserting a digit into a two-digit inch value always produces a three-digit one, which is never a plausible height, so the class can only fire on a value with a single-digit integer part. Among the 6,619 height anomalies whose integer part has two or more digits it reconciles 0. Its entire 5.47% is the whole-foot family reached by another route.

Two clusters are visible without any anchor at all. 1,371 visits record a `height_in` of 1 to 6 as an exact integer, median age 5.2 years — a height of 3 or 4 for a child three or four feet tall. And 143 record a `height_in` between 90 and 115, which read as inches is implausible and read as centimetres is an ordinary preschool stature at a median age of 3.1 years. The recording grid decides between the two readings: 35.0% of that cluster falls on the quarter-inch grid against 80.0% of all heights, so those values never passed through the inch-typing workflow.

Weight. 6,196 anomalies in the testable interior.

Weight: mechanisms against the null

MECHANISM	RECONCILED	SHARE	NULL	RATIO
pound value in the ounce field	39	0.63%	0.26%	2.4x
ounce value where a pound is expected	25	0.40%	0.09%	4.7x
kilogram value in the ounce field	13	0.21%	0.09%	2.5x
gram value in the ounce field	2	0.03%	0.08%	0.4x
decimal point misplaced	1,208	19.50%	1.13%	17.2x
adjacent digit transposition	77	1.24%	6.65%	0.2x
one digit omitted	927	14.96%	8.63%	1.7x
one digit wrong (calibration class)	795	12.83%	25.39%	0.5x

Transposition is again below chance, so neither channel shows evidence of digit swapping. A misplaced decimal point, which the height channel does not show at all, is the dominant weight artifact: 1,208 anomalies at 17 times the null rate, the strongest enrichment measured anywhere in this report. An ounce value has more digits than an inch value and no natural decimal point, so a factor of ten is both easy to key and hard to notice.

The calibration row is why the null is not optional. Allowing any single digit to be wrong reconciles about half of all height anomalies and reconciles almost exactly as many randomly paired values. Reported without a null it would look like the largest finding here.

How strong is the transposition negative? Only as strong as the share of transpositions the anomaly gate could have caught. Applying every adjacent digit swap to a sample of measurements in the testable interior gives that share directly: 69.7% of height swaps would displace a value past the gate, against 31.6% of weight swaps. The height negative is well powered; the weight negative rules out only large swaps, since a four-digit ounce value can absorb a swap without moving far.

What the mechanisms account for

CHANNEL	ANOMALIES	A NAMED MECHANISM FITS	ONLY THE CALIBRATION CLASS	NOTHING FITS
height	7,443	939	3,989	2,515
weight	6,196	1,987	742	3,467

Implications for analysis. Digit transposition can be dropped from the checklist for this extract at the magnitude that displaces a measurement from its own trajectory; for weight the same test is only about a third sensitive, so a small swap is not ruled out. Unit confusion and decimal placement do matter, and both are cheap to screen because both produce values implausible on their face. Bound `height_in` and `weight_oz` before any conversion, and check the recording grid rather than the value alone — the grid separates a tall adolescent from a centimetre in the wrong field where magnitude cannot. Note also that 1,371 of the whole-foot entries and 143 of the centimetre cluster already carry a null `height_cm`: the derived layer's own bound removes them as a side effect, so anyone reading the derived channels is protected and anyone reading the raw ones is not.

4.5 Repeated measurements: zero growth and apparent height loss

Children do not shrink, so a recorded decrease is recording behaviour rather than physiology. That much is easy. What matters is that the behaviour is not one thing, and the interval between measurements separates the mechanisms.


Repeat height pairs at age 2 or later, by interval

INTERVAL	PAIRS	EXACTLY ZERO CHANGE	ANY DECREASE	MEDIAN LOSS
up to 7 days	25,968	47.86%	21.31%	0.69 cm
8 to 30 days	82,652	32.00%	19.03%	0.64 cm
31 to 90 days	170,521	16.38%	10.51%	0.64 cm
91 to 180 days	205,153	5.15%	4.22%	0.81 cm

INTERVAL	PAIRS	EXACTLY ZERO CHANGE	ANY DECREASE	MEDIAN LOSS
181 to 365 days	368,248	1.57%	1.44%	0.79 cm
over 365 days	893,691	0.64%	0.66%	0.64 cm

At short intervals a child genuinely has not grown a measurable amount and the quarter-inch grid absorbs the rest. At long intervals both effects should vanish, and they do not entirely. The residue is small and its median size is about one grid step of 0.635 cm, which is the first clue that it is rounding rather than error.

Holding the interval fixed and varying age identifies the mechanisms directly.




Apparent height loss by age, over 181-365 day intervals

Apparent loss by age at the earlier measurement

AGE BAND (MONTHS)	PAIRS	ANY DECREASE	MEDIAN LOSS	MEAN CHANGE
18-24	63,685	0.53%	1.27 cm	5.51 cm
24-30	52,826	1.21%	0.99 cm	5.20 cm
30-36	19,242	3.65%	1.25 cm	3.75 cm
36-42	32,650	0.44%	1.63 cm	6.19 cm
48-60	37,302	0.37%	1.91 cm	5.68 cm
60-84	64,090	0.33%	1.92 cm	5.14 cm
84-120	76,194	0.36%	1.60 cm	4.70 cm
120-144	37,388	0.47%	1.25 cm	4.87 cm
144-168	26,514	2.78%	0.64 cm	4.04 cm
168-192	14,518	11.39%	0.64 cm	1.79 cm
192-216	2,566	23.69%	0.64 cm	0.54 cm

Two separate excesses, with different signatures. The first is a narrow spike at 30 to 36 months carrying a median loss of over a centimetre — the age at which recumbent length gives way to standing height, and a standing height

genuinely is shorter than a recumbent length for the same child. It is a change of measurement protocol recorded in a field that does not name the protocol.



Apparent loss in adolescence, by sex

Adolescent bands split by recorded sex

AGE BAND (MONTHS)	FEMALE PAIRS	FEMALE DECREASE	FEMALE MEAN CHANGE	MALE PAIRS	MALE DECREASE	MALE MEAN CHANGE
144-168	12,858	5.29%	2.53 cm	13,656	0.41%	5.46 cm
168-192	7,127	18.59%	0.70 cm	7,391	4.45%	2.83 cm
192-216	1,346	28.83%	0.22 cm	1,220	18.03%	0.89 cm

The second excess is the adolescent rise, and the sex split identifies it. Girls reach the high rates about two years before boys, in the same order as growth cessation, while the mean change over the same interval falls towards zero. Once annual growth drops below the recording grid, re-measuring a child who has stopped growing returns a lower value about as often as a higher one. Restricting to ages 2 to 10, where growth is unambiguously ongoing, collapses the long-interval decrease rate from 0.663% to 0.083% — 565 pairs of 681,114.

What survives both explanations divides again. Of 1,188 decreases over a centimetre across more than a year that are followed by a further measurement, 725 (61.0%) are followed by a value back at or above the earlier level, and 463 (39.0%) by one that stays below it. In the first the low value is the suspect; in the second it is corroborated and the earlier, higher measurement is the candidate error.

Implications for analysis. Most apparent shrinkage here is not error and should not be filtered as an outlier: it is the recording grid acting on a flattened trajectory, plus a protocol change at two to three years. A synthetic or smoothed trajectory that lacks both will not resemble this panel. Where a decrease does need adjudication, 39% of long-interval losses persist into the next measurement, so a rule that always discards the lower value is wrong on that share.


4.6 Derived z-scores and percentiles: bounds and saturation

The derived channels are not a neutral restatement of the measurements. Each carries its own support, and they do not share one.

Z-score channels

CHANNEL	VALUES	MINIMUM	MAXIMUM	BEYOND 15I
height z	3,491,616	-4.9992	3.0000	0
weight z	6,482,932	-4.9991	4.9995	0
BMI z	1,955,337	-18.7803	6.7026	400
head circ z	1,635,640	-17,485.9110	306,212.6000	16,663
weight-for-length z	2,027,317	-145.6016	7.6285	1,123
weight-for-stature z	1,371,347	-14.3445	7.4762	246

The height z-score is bounded above at exactly 3.00 while its lower tail runs past -4.99. The truncation leaves no pile-up at the boundary, so it is invisible in a summary: only 21 visits sit at or above +3. The asymmetry is what exposes it. In the lower tail 45.4% of the mass beyond $|z| = 2.5$ continues past 3; if the upper tail behaved the same way roughly 15,800 visits would sit above +3.



Height z-score, both tails

Percentile channels and their saturation points

CHANNEL	VALUES	EXACTLY 0	SHARE	EXACTLY 100	SHARE
height	3,491,616	2,801	0.080%	0	0.000%
weight	6,482,932	3,584	0.055%	5,221	0.081%
BMI	1,955,337	1,599	0.082%	1,590	0.081%
weight-for-length	2,027,317	6,151	0.303%	1,203	0.059%
weight-for-stature	1,371,347	1,509	0.110%	856	0.062%


Implications for analysis. The height channel cannot support any question about tall stature: its upper tail is absent, and a trajectory approaching the bound from below is distorted too. The percentile channels carry point masses at exactly 0 and 100 that are saturated rather than measured, so they are not continuous and should not be modelled as such. Because the four z channels do not share a support, a model consuming several of them together inherits the inconsistency silently. Recomputing from the raw measurement against a stated reference avoids all of this.

4.7 Head circumference: a recoverable conversion defect

15,025 visits carry a head circumference outside the conventional review range of 25 to 65 cm. Read as a distribution that looks like a badly behaved channel. Read as clusters, it looks like arithmetic.

Head-circumference values by band

BAND	VISITS	MEDIAN	MINIMUM	MAXIMUM
below 10 cm	174	4.10 cm	0.00 cm	9.65 cm
10 to under 25 cm	943	18.00 cm	10.00 cm	24.77 cm
25 to 65 cm (within review range)	1,620,665	44.00 cm	25.00 cm	65.00 cm
over 65 to 200 cm	13,472	110.49 cm	65.50 cm	193.04 cm
above 200 cm	436	252.22 cm	202.57 cm	505.46 cm



Where head-circumference values fall

The cluster between 65 and 200 cm is not noise. It holds 13,472 visits, and 13,467 of them — 99.96% — fall back inside the review range when divided by 2.54, with a median of 43.5 cm. That is an ordinary infant head circumference. These are centimetre values that were put through an inch-to-centimetre conversion a second time. A further 436 visits sit above 200 cm, of which 356 become plausible after dividing by 2.54 twice, consistent with the same conversion applied again.

This one defect explains most of the damage. Of 16,663 visits with an absolute head-circumference z-score above 5, 14,899 (89.4%) sit on a measurement outside the review range, and the double-converted cluster alone

accounts for 13,467 of them. The remaining 1,764 visits carry a plausible measurement and still produce an extreme z, so the z transform is independently defective and repairing the units would not fully fix the channel. That is why 4.6 shows this channel with a maximum no measurement could produce.

Implications for analysis. A declared plausible range deletes all 15,025 out-of-range values, but 13,467 of those (89.6%) are ordinary infant measurements that one documented division restores. Repairing before bounding is strictly better than bounding alone, and it recovers most of the only measurement channel whose declared range removes a non-trivial share of values. The derived z-score is a separate matter: it stays unusable on 1,764 visits even after the units are fixed, so recompute it from the repaired measurement rather than consuming it as distributed.


4.8 The distributed delta and velocity fields

The augmented visit layer distributes `delta_height_cm`, `delta_age_in_days_height`, and the velocity fields derived from them. These are **not** a lag over successive measurements, and reading them as one is the error this subsection exists to prevent. For each measurement the pipeline walks backwards to the most recent earlier measurement whose age gap meets an age-dependent minimum, skipping every measurement in between.

The interval rule, inferred from the data

AGE BAND	CONDITION ON CURRENT AGE	MINIMUM INTERVAL
birth to 12 months	up to 365 days	90 days
1 to 2 years	up to 730 days	180 days
2 to 12 years	up to 4380 days	335 days
13 years and over	beyond 4380 days	180 days

Applying that rule reproduces the distributed fields. Across 2,786,770 visits carrying a nonmissing `delta_height_cm`, the recomputed age gap matches the distributed one on 2,786,770 rows (100.00%), the recomputed delta matches within one hundredth of a centimetre on 2,786,432 rows (99.988%), and the recomputed velocity matches on 99.68% of rows. A naive lag over successive height-bearing visits matches only 1,218,842 rows (43.7%) — which is what makes these fields look unreproducible when the rule is not known.



The most common recorded measurement intervals

Two residuals are worth recording. 372,482 rows differ by exactly one hundredth of a centimetre, because the pipeline rounds half to even while this check rounds half away from zero; heights come from a quarter-inch grid, so exact halfway cases are common rather than rare. Only 338 rows (0.012%) differ by more than that, and they sit on the duplicate patient-days of 3.1, where which earlier height was used is ambiguous.

Implications for analysis. The velocity channels are usable as distributed, which a distributional summary alone could not establish. What must travel with them is the definition: a velocity here is computed over an interval of at least 90 to 335 days depending on age, not between adjacent visits, so it is already smoothed relative to a visit-to-visit rate and cannot be compared with one. Any recomputation, and any synthetic series carrying a velocity, must use the same rule or the two are not on the same scale. The rounding to two decimals is part of what the distributed values are.


4.9 Age- and sex-stratified growth profile

A reference table for anyone who needs to know what ordinary looks like in this extract before deciding what is unusual. Mean z-scores run from -0.00 to 0.36 across the age and sex cells, so the cohort sits close to the reference population on average even though it is not a sample of one.

Measurements and derived z-scores by age band and sex

AGE BAND (YEARS)	SEX	VISITS	PATIENTS	MEAN HEIGHT	MEAN HEIGHT Z	HEIGHT Z SD	MEAN WEIGHT Z	MEAN BMI Z
0-2	F	1,294,946	95,419	66.3 cm	0.363	1.002	-0.144	—
0-2	M	1,397,984	99,226	68.0 cm	0.341	0.992	-0.167	—
2-5	F	677,359	92,851	96.0 cm	0.354	0.974	0.285	0.150
2-5	M	716,626	96,604	97.3 cm	0.338	0.995	0.369	0.148
5-10	F	745,589	82,246	122.5 cm	0.119	0.981	0.362	0.418
5-10	M	784,028	85,364	123.6 cm	0.208	0.989	0.451	0.428
10-15	F	366,624	50,775	151.9 cm	0.218	0.993	0.489	0.451

AGE BAND (YEARS)	SEX	VISITS	PATIENTS	MEAN HEIGHT	MEAN HEIGHT Z	HEIGHT Z SD	MEAN WEIGHT Z	MEAN BMI Z
10-15	M	375,601	52,556	152.7 cm	0.312	1.012	0.575	0.475
15-18	F	72,447	17,530	162.5 cm	-0.000	0.991	0.452	0.453
15-18	M	63,194	18,049	173.9 cm	0.147	0.950	0.646	0.476



Mean height z-score by age band and sex

Implications for analysis. Read the height-z column against 4.6 before using it: its upper tail is truncated at +3, so every mean here is pulled very slightly downward relative to an untruncated reference, and the effect grows in the bands where tall children are most numerous. The SD column is the more useful one for scaling, and it is close to 1 by construction of the z transform rather than as a finding.

4.10 Within-child dependence in the height channel

Repeated measurements of one child are not independent observations, and the size of that dependence decides how much information a visit count actually carries. Measured on the height z-score at age 2 or later, across 199,727 patients with at least two values.

Variance components and serial correlation

QUANTITY	VALUE	WHAT IT SAYS
between-child SD of patient means	0.9140	how far children sit from one another
within-child SD about a patient's own mean	0.4262	how much one child's channel moves
implied intraclass correlation	0.8214	share of variance that is between children
lag-1 autocorrelation	0.9249	correlation of successive values, 1,744,631 pairs

A child's height z-score is strongly self-similar: successive values correlate at 0.925, and 82.1% of the total variance is between children rather than within them. The design-effect consequence is blunt: in the limit of many

measurements a child contributes about 1.2 independent observations, not one per visit, however many visits are recorded.


Implications for analysis. Resample and model at the patient level, not the visit level: a visit-level standard error on any quantity aggregated across this panel will be far too small. And treat these as sample statistics rather than the parameters of a process that would generate them — a patient's mean carries residual variation as well as the child's own level, so the between-child SD of patient means overstates the underlying channel SD, while the sample SD within a positively autocorrelated series understates its marginal SD. Calibrate a generative model against these by simulation rather than by setting its parameters equal to them.

4.11 BMI: recomputation and recorded categories

BMI is the one derived channel that can be checked against its own inputs. Across 1,955,339 visits carrying a BMI together with both a weight and a height, recomputing weight in kilograms over height in metres squared gives a median absolute difference of 9.7e-07 and a 95th percentile of 3.0e-06 — floating-point noise, nothing more. 0 visits differ by more than 0.1. The channel is internally consistent, so a BMI here disagreeing with your own calculation means you used a different height or weight, not that the field is wrong.

Recorded BMI categories

CATEGORY	VISITS	SHARE OF CATEGORISED VISITS	DISTINCT PATIENTS
underweight	83,602	4.3%	33,608
normal	1,338,418	68.4%	193,723
overweight	280,226	14.3%	84,669
obese	253,091	12.9%	49,998



Distribution of recorded BMI categories

The category is present only where a BMI percentile is, which 1.3 and 3.4 show means age 2 or later. Of 1,955,337 categorised visits, 14.3% are overweight and 12.9% obese.


Implications for analysis. This is a distribution over recorded visits, not a prevalence: children with more visits contribute more rows, BMI is missing selectively by age and encounter type, and 1.4 shows the cohort is not a population sample. Aggregate to the patient before quoting any proportion, state the age window, and prefer the continuous percentile to the category where the analysis allows it, since the cut points discard most of the information.

5. Other clinical domains

Diagnoses, laboratory results, medications, referrals, and demographics.

5.1 Diagnoses

Diagnoses arrive two ways: up to 33 coded slots per encounter, and a problem list that is not visit-linked. 14,714,503 encounter slots are filled across 250,563 patients, and 6,154,801 visits (94.8%) carry at least a first diagnosis.



Coded diagnoses per visit

Most frequently recorded encounter diagnoses

ICD-10	DESCRIPTION	SLOTS	PATIENTS
Z00.129	Encounter for routine child health examination without abnormal findings	2,494,658	247,896
Z23	Encounter for immunization	1,438,556	244,583
J06.9	Acute upper respiratory infection, unspecified	363,785	132,869
J02.9	Acute pharyngitis, unspecified	323,876	113,944
Z13.88	Encounter for screening for disorder due to exposure to contaminants	295,071	128,827
Z13.0	Encounter for screening for diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	292,612	135,760
R50.9	Fever, unspecified	265,363	115,401
Z71.3	Dietary counseling and surveillance	230,303	81,930
Z71.82	Exercise counseling	209,541	78,003
R05.9	Cough, unspecified	199,083	91,945
Z00.121	Encounter for routine child health examination with abnormal findings	163,509	72,084
Z20.822	Contact with and (suspected) exposure to COVID-19	160,041	65,268

ICD-10	DESCRIPTION	SLOTS	PATIENTS
J02.0	Streptococcal pharyngitis	129,310	71,029
B34.9	Viral infection, unspecified	125,662	68,886
Z00.110	Health examination for newborn under 8 days old	124,627	105,062
IMO0002	[not in the ICD-10 lookup]	116,950	42,737
F90.2	Attention-deficit hyperactivity disorder, combined type	112,292	15,694
Z29.3	Encounter for prophylactic fluoride administration	107,178	52,897
Z68.53	Body mass index [BMI] pediatric, 85th percentile to less than 95th percentile for age	105,889	52,088
J45.20	Mild intermittent asthma, uncomplicated	93,013	24,479
R21	Rash and other nonspecific skin eruption	92,028	62,699
K21.9	Gastro-esophageal reflux disease without esophagitis	90,172	31,748
F41.9	Anxiety disorder, unspecified	81,476	24,519
K59.00	Constipation, unspecified	80,647	39,182
H66.001	Acute suppurative otitis media without spontaneous rupture of ear drum, right ear	76,772	49,374

The 25 most frequent of 8,029 distinct values, covering 52.8% of filled slots; the remaining 8,004 values hold the rest. Every count here is a recorded frequency within a selected cohort. Patients carrying any code that occurred fewer than 11 times were removed before delivery (1.4), so rare entries are absent by construction and nothing in this table is a population rate.

The problem list holds 1,709,584 entries for 238,823 patients, of which 44.3% carry a resolved age. As 3.5 shows, the remainder are open problems rather than missing dates.

Most frequently recorded problem-list diagnoses

ICD-10	DESCRIPTION	ENTRIES	PATIENTS
U07.1	COVID-19	26,260	26,260
Z28.21	Immunization not carried out because of patient refusal	21,189	21,189
F41.9	Anxiety disorder, unspecified	20,950	20,950
K59.00	Constipation, unspecified	17,491	17,491
Z00.129	Encounter for routine child health examination without abnormal findings	17,348	17,348
K21.9	Gastro-esophageal reflux disease without esophagitis	17,184	17,184
Z86.16	Personal history of COVID-19	16,007	16,007
J45.20	Mild intermittent asthma, uncomplicated	15,088	15,088
L30.9	Dermatitis, unspecified	14,868	14,868
R46.89	Other symptoms and signs involving appearance and behavior	14,789	14,789
F80.9	Developmental disorder of speech and language, unspecified	13,661	13,661
F80.1	Expressive language disorder	13,650	13,650
L20.83	Infantile (acute) (chronic) eczema	13,282	13,282
F90.2	Attention-deficit hyperactivity disorder, combined type	12,918	12,918
IMO0002	[not in the ICD-10 lookup]	12,915	12,915

ICD-10	DESCRIPTION	ENTRIES	PATIENTS
R62.51	Failure to thrive (child)	12,503	12,503
J30.9	Allergic rhinitis, unspecified	11,383	11,383
Z91.018	Allergy to other foods	11,308	11,308
J06.9	Acute upper respiratory infection, unspecified	11,197	11,197
B08.1	Molluscum contagiosum	10,781	10,781
Z38.00	Single liveborn infant, delivered vaginally	10,304	10,304
R63.39	Other feeding difficulties	10,088	10,088
R01.1	Cardiac murmur, unspecified	10,081	10,081
G47.9	Sleep disorder, unspecified	9,772	9,772
J45.909	Unspecified asthma, uncomplicated	9,502	9,502

The 25 most frequent of 4,739 distinct values, covering 20.7% of entries; the remaining 4,714 values hold the rest.

Both tables above count literal codes, which is the right unit for describing what gets typed but the wrong one for counting a condition. Rolling the same data up to the three-character category changes which diagnoses appear at all — see 3.9, and note that 1,204 of the 1,327 categories in this extract never appear as a bare code, so an exact-match query for them returns zero.

The same diagnoses rolled up to their ICD-10 category

CATEGORY	DESCRIPTION	PATIENTS
Z00	Encounter for general examination without complaint, suspected or reported diagnosis	250,354
Z23	Encounter for immunization	244,617
Z13	Encounter for screening for other diseases and disorders	188,758
J06	Acute upper respiratory infections of multiple and unspecified sites	134,368
H66	Suppurative and unspecified otitis media	132,693
J02	Acute pharyngitis	131,194
R50	Fever of other and unknown origin	119,263
R05	Cough	115,563
Z71	Persons encountering health services for other counseling and medical advice, not elsewhere classified	99,130
Z20	Contact with and (suspected) exposure to communicable diseases	78,217
R63	Symptoms and signs concerning food and fluid intake	76,097
B34	Viral infection of unspecified site	74,947
H10	Conjunctivitis	74,858
Z68	Body mass index [BMI]	64,527
R21	Rash and other nonspecific skin eruption	63,738
P92	Feeding problems of newborn	62,842
Z29	Encounter for other prophylactic measures	61,612

CATEGORY	DESCRIPTION	PATIENTS
K59	Other functional intestinal disorders	57,481
H65	Nonsuppurative otitis media	57,167
L20	Atopic dermatitis	52,206
R09	Other symptoms and signs involving the circulatory and respiratory system	51,549
J30	Vasomotor and allergic rhinitis	48,747
R06	Abnormalities of breathing	48,551
Z28	Immunization not carried out and underimmunization status	47,456
IMO	[not in the ICD-10 lookup]	45,678

The 25 most frequent of 1,327 distinct values; 1,302 more are not shown.

Implications for analysis. Encounter diagnoses and problem-list entries answer different questions and should not be pooled without saying why: the first is what was coded at a contact, the second is what the chart asserts about the child, including resolved history. Neither is an adjudicated clinical truth, and a code's absence is not evidence a condition was absent.

5.2 Laboratory results

17,230,681 resulted components across 6,578,838 lab orders for 247,271 patients — 2.6 components per order. The grain is the component, not the order, which is the single most common source of double counting in this resource.

Most frequently ordered lab procedures

PROCEDURE	ROWS	PATIENTS
CBC	2,742,117	65,688
CBC DIFFERENTIAL	1,660,900	65,105
CE EXTERNAL LAB	1,455,867	152,867
URINALYSIS	1,326,746	46,480
POCT URINALYSIS DIPSTICK	1,079,426	54,102
COMPREHENSIVE METABOLIC PANEL	475,461	28,764
POCT COVID-19 NUCLEIC ACID (AMPLIFIED PROBE)	432,267	80,837
LEAD, BLOOD	394,009	93,596
COVID-19 (CORONAVIRUS 2019) PCR	392,834	97,191
POCT STREP A NUCLEIC ACID (AMPLIFIED PROBE)	314,977	75,754
POCT CBC WITH DIFF	303,486	15,309
POCT INFLUENZA A/B NUCLEIC ACID (AMPLIFIED PROBE)	272,419	48,609
POCT RAPID STREP A IMMUNOASSAY	268,011	69,548
POCT COVID-19, INFLUENZA, AND RSV NUCLEIC ACID (AMPLIFIED PROBE)	267,208	22,586

PROCEDURE	ROWS	PATIENTS
URINE CULTURE	214,567	43,877
POCT HEMOGLOBIN	193,586	86,845
RAPID STREP A, IMMUNOASSAY	182,893	45,620
POCT INFLUENZA A/B IMMUNOASSAY	172,419	35,207
LIPID PANEL	148,974	23,764
STREP A CULTURE	147,291	37,984
HEMOGLOBIN	140,801	60,227
THROAT CULTURE	131,125	37,007
URINALYSIS WITH MICROSCOPIC	130,811	7,635
STREP A NUCLEIC ACID DETECTION	119,448	28,102
INFLUENZA A/B NUCLEIC ACID	112,796	22,382

The 25 most frequent of 3,742 distinct values, covering 75.9% of rows; the remaining 3,717 values hold the rest. Every count here is a recorded frequency within a selected cohort. Patients carrying any code that occurred fewer than 11 times were removed before delivery (1.4), so rare entries are absent by construction and nothing in this table is a population rate.

2,494,261 rows (14.5%) carry no result value at all, and 2,283,186 orders (34.7%) have no resulted component on any line. Both are expected rather than broken: the extract includes externally sourced labs that arrive without results. 3.6 covers how the values that do exist are shaped.

Implications for analysis. Count orders when you mean tests and rows when you mean components, and never mix them in a rate. An order-with-no-result is a documented ordering event, not a missing result to impute.

5.3 Medications

3,823,049 medication records for 236,323 patients. A record is an order placed by a practice clinician or a documentation of an outside or historical medication, and the two behave differently.

Record type and date completeness

RECORD TYPE	RECORDS	PATIENTS	START AGE PRESENT	END AGE PRESENT
Internal	3,250,374	229,099	98.7%	91.6%
External	572,675	158,974	58.1%	69.1%

Most frequently recorded medications

GENERIC NAME	RECORDS	PATIENTS
Amoxicillin	351,609	136,002
Albuterol Sulfate	312,748	65,240

GENERIC NAME	RECORDS	PATIENTS
Methylphenidate HCl	219,731	13,658
Dexmethylphenidate HCl	124,415	8,083
Amphetamine-Dextroamphetamine	106,250	6,889
Acetaminophen	83,235	45,771
Cefdinir	81,224	41,608
Sodium Fluoride	78,612	31,439
Fluticasone Propionate HFA	77,412	19,545
Amoxicillin-Pot Clavulanate	76,894	47,580
EPINEPHrine	76,835	18,903
Ibuprofen	75,866	44,933
Mupirocin	74,769	51,626
Hydrocortisone	72,204	40,563
Cetirizine HCl	72,065	34,707
Azithromycin	69,883	46,509
Cephalexin	64,757	43,345
Triamcinolone Acetonide	60,074	31,238
Nystatin	58,782	34,023
FLUoxetine HCl	56,361	6,637
Cholecalciferol	55,797	40,031
Erythromycin	53,467	39,136
Sertraline HCl	49,233	6,289
guanFACINE HCl	48,115	6,701
Spacer/Aero-Holding Chambers	46,362	28,820


The 25 most frequent of 1,073 distinct values, covering 64.0% of records; the remaining 1,048 values hold the rest. Every count here is a recorded frequency within a selected cohort. Patients carrying any code that occurred fewer than 11 times were removed before delivery (1.4), so rare entries are absent by construction and nothing in this table is a population rate.

Three documented fields were never delivered. The data dictionary describes 3 medication classification columns — `med_therapeutic_class`, `med_pharmaceutical_class`, `med_pharmaceutical_subclass` — and none is present in the extract. Any analysis by drug class has to map `med_simple_generic_name` itself.

Implications for analysis. A record is not an administration and not evidence the child took the drug. Externally documented records carry a documentation date in the order-date column and approximate start dates, so exposure windows built from them are unreliable; 3.3 measures how often the dates contradict each other.

5.4 Referrals

349,827 referral orders for 138,071 patients. A referral is a recorded action, not an outcome: it says a clinician placed an order, not that the child was seen.



Referrals by age at order

Most frequently requested specialties

SPECIALTY	REFERRALS	PATIENTS	MEDIAN AGE
Otolaryngology	35,723	29,567	3.99 y
Ophthalmology	24,298	20,605	4.58 y
Orthopedic Surgery	22,887	19,521	9.97 y
Allergy	21,761	18,258	5.03 y
Behavioral Health	21,748	16,442	9.29 y
Dermatology	20,652	17,851	7.59 y
Audiology	15,972	13,616	2.24 y
Gastroenterology	14,344	12,467	5.57 y
Cardiology	13,610	12,134	6.83 y
Neurology	11,275	9,744	6.92 y
Nutrition	11,035	8,910	9.39 y
Urology	10,697	9,140	4.02 y
Speech Pathology	9,879	7,634	3.84 y
Physical Therapy	9,861	8,249	11.25 y
Early Intervention	9,302	8,511	1.41 y
Developmental Medicine	8,532	6,794	3.91 y
Endocrinology	6,641	5,583	9.37 y
Occupational Therapy	6,573	5,002	5.14 y
General Surgery	4,206	3,889	3.78 y
Psychology	4,134	3,537	9.03 y
Pulmonary Disease	3,926	3,539	4.80 y

SPECIALTY	REFERRALS	PATIENTS	MEDIAN AGE
Plastic Surgery	3,454	3,269	0.52 y
Psychiatry	2,885	2,458	10.59 y
Podiatry	2,863	2,608	11.48 y
Neurosurgery	2,415	2,302	0.40 y

The 25 most frequent of 119 distinct values, covering 85.4% of referrals; the remaining 94 values hold the rest. Every count here is a recorded frequency within a selected cohort. Patients carrying any code that occurred fewer than 11 times were removed before delivery (1.4), so rare entries are absent by construction and nothing in this table is a population rate.

27,452 referrals (7.85%) carry no requested specialty and 26,601 (7.6%) no requested visit count. The data dictionary also warns that referrals are not always documented in the source system, so absence of a referral is not evidence none was made.

Implications for analysis. This resource is positive-unlabelled: recorded referrals are real, but unrecorded ones are indistinguishable from referrals that never happened. Combined with the partial visit link measured in 3.2, a referral rate computed here is a documentation rate. Treat it as such and say so.

5.5 Recorded identity and patient-level observation

Identity fields are recorded categories, not attributes of the children. Non-response is shown separately from every substantive category, because blank, unknown, and declined are not clinically equivalent to a recorded value but are all missing for the purpose of a subgroup comparison.

Recorded sex

CATEGORY	PATIENTS	SHARE
M	127,699	51.0%
F	122,883	49.0%
U	6	0.0%

Recorded ethnicity

CATEGORY	PATIENTS	SHARE
Not Hispanic or Latino	170,594	68.1%
Hispanic or Latino	28,549	11.4%
Choose not to Answer	24,566	9.8%
Unknown	20,834	8.3%
[blank]	5,464	2.2%
Unable to collect	450	0.2%
Patient does not know	131	0.1%

First recorded race

CATEGORY	PATIENTS	SHARE
White	155,375	62.0%
Unknown	23,085	9.2%
Choose not to answer	17,534	7.0%
Another Race	15,950	6.4%
Asian	15,661	6.2%
Black or African American	12,162	4.9%
[blank]	8,818	3.5%
American Indian or Alaska Native	625	0.2%
Middle Eastern or Northern African	512	0.2%
Unable to collect	492	0.2%
Native Hawaiian or Other Pacific Islander	248	0.1%
Patient does not know	126	0.1%

Race is a multi-select of up to eight slots; only the first is shown. 13,191 patients (5.3%) have a second race recorded, so this table understates multiracial identity.

Observation per patient is dense, as the cohort rule in 1.4 requires. The median patient has 23 visits (quartiles 15 and 34, 95th percentile 56, maximum 244), spanning a median of 7.0 years (quartiles 3.3 and 10.9). The median patient's last recorded visit is at age 8.3 years.

Implications for analysis. Identity non-response is large enough to change a subgroup contrast on its own, so report it as its own category rather than dropping it. And because entry to this cohort required both a measurement history and a recent visit, the visit distribution describes the selection as much as the care; it is a feasibility figure, not an estimate of pediatric utilisation.


5.6 Patient-level derived flags and summaries

The augmented patient layer carries seven boolean flags and a block of per-patient z-score summaries. They are conveniences computed from the visit layer, not independent observations, and each inherits whatever the channel it summarises does — the height-z flags inherit the truncation of 4.6, the BMI flags inherit the age-2 floor of 1.3.

Patient-level flags

FLAG	SET WHEN THE PATIENT	PATIENTS	SHARE OF COHORT
healthy_flag	carries none of the tracked conditions	24,471	9.8%
chronic_dx_flag	any chronic diagnosis	203,935	81.4%
growth_dx_flag	any of the tracked growth-relevant diagnoses	35,907	14.3%
ever_stunting_flag	height z below the stunting threshold at any visit	17,889	7.1%
ever_wasting_flag	weight-for-length or -stature below the wasting threshold	66,704	26.6%

FLAG	SET WHEN THE PATIENT	PATIENTS	SHARE OF COHORT
ever_underweight_flag	BMI below the underweight threshold at any visit	33,608	13.4%
ever_obesity_flag	BMI at or above the obesity threshold at any visit	49,998	20.0%



Patients carrying each derived flag

`growth_dx_flag` marks 35,907 patients. Where an age at diagnosis is observed (35,890 patients) its median is 0.027 years, and 25,208 of those (70.2%) are assigned their code within the first month of life. That is a statement about when the code was recorded, not about when a condition began.

Per-patient z-score summaries, averaged over patients with more than one value

CHANNEL	PATIENTS	MEAN OF PATIENT MEANS	MEAN OF PATIENT SDS
height	248,172	0.2980	0.5147
weight	249,595	0.1275	0.5371
BMI	199,693	0.2927	0.5309

Implications for analysis. A flag is a recorded derivation, not an adjudicated clinical state, and the concentration of growth-diagnosis ages in the first month shows why: much of what the flag marks is perinatal coding rather than a growth trajectory that was observed and interpreted over years. Use the flags to describe the derived layer or to stratify descriptively; recompute from the visit layer against a stated rule if a flag is doing analytic work.

5.7 The extract's growth orientation: tracked codes and referral pathways

This extract was assembled around growth. Cohort entry required a growth-measurement history (1.4), and the augmentation layer records, for each patient, the age at which any of 33 specific diagnosis codes was first recorded. That panel is a design choice made upstream, and knowing which codes are in it is the difference between using the derived columns and guessing at them.

Because ICD-10 is a hierarchy (3.9), each code is counted here twice: as a literal string, and as a subtree including every descendant. The gap between the two columns is what a flat query would miss.

The tracked growth-relevant diagnosis codes

ICD-10	DESCRIPTION	DERIVED COLUMN	PATIENTS, LITERAL CODE	PATIENTS, CODE AND DESCENDANTS	MISSED BY A FLAT COUNT
P92.6	Failure to thrive in newborn	14,428	14,428	14,428	0
P07	Disorders of newborn related to short gestation and low birth weight, not elsewhere classified	11,014	0	11,029	11,029
P05	Disorders of newborn related to slow fetal growth and fetal malnutrition	4,069	0	4,074	4,074
E30.1	Precocious puberty	3,405	3,406	3,406	0
P70	Transitory disorders of carbohydrate metabolism specific to newborn	3,353	0	3,354	3,354
K90.0	Celiac disease	898	898	898	0
E10	Type 1 diabetes mellitus	491	0	491	491
E34.3	Short stature due to endocrine disorder	447	0	447	447
E30.0	Delayed puberty	419	419	419	0
E03.9	Hypothyroidism, unspecified	309	309	309	0
Q90	Down syndrome	205	0	205	205
E23.0	Hypopituitarism	150	150	150	0
K50	Crohn's disease [regional enteritis]	113	0	113	113
E34.4	Constitutional tall stature	83	83	83	0
N18	Chronic kidney disease (CKD)	70	0	70	70
K51	Ulcerative colitis	62	0	62	62
Q87.1	Congenital malformation syndromes predominantly associated with short stature	58	0	58	58
P04.3	Newborn affected by maternal use of alcohol	53	53	53	0
Q87.3	Congenital malformation syndromes involving early overgrowth	46	46	46	0
Q98.4	Klinefelter syndrome, unspecified	42	42	42	0
Q96	Turner's syndrome	36	0	36	36
Q87.2	Congenital malformation syndromes predominantly involving limbs	32	32	32	0
E23.6	Other disorders of pituitary gland	31	31	31	0
Q98.0	Klinefelter syndrome karyotype 47, XXY	26	26	26	0
Q87.4	Marfan syndrome	17	0	17	17
Q98.5	Karyotype 47, XYY	17	17	17	0

ICD-10	DESCRIPTION	DERIVED COLUMN	PATIENTS, LITERAL CODE	PATIENTS, CODE AND DESCENDANTS	MISSED BY A FLAT COUNT
Q77	Osteochondrodysplasia with defects of growth of tubular bones and spine	15	0	15	15
Q78.0	Osteogenesis imperfecta	10	10	10	0

Codes carried by fewer patients than the suppression threshold are omitted. Counts are recorded frequencies inside a cohort that excluded every patient with a code seen fewer than 11 times (1.4), so this panel cannot be read as prevalence.

The upstream derivation is hierarchical, and the two count columns verify it. 14 of the 33 tracked codes have descendants in this extract; the other 19 have none, so both readings coincide and they cannot distinguish the two rules. Of the 14 that can, **0 match the literal count** — in every case the derived column follows the subtree. The evidence is starkest because **all 14 of those codes never appear as a literal string at all**: an exact-match query returns zero patients for `E10` , `P07` , `K50` and the rest, while the derived column correctly reports hundreds or thousands.

4 codes (`P07` , `P05` , `E30.1` , `P70`) sit slightly below their subtree count. The shortfall is explained rather than unexplained: those patients carry the code only on a problem-list entry with no noted date, so no age could be determined. The derived column therefore means *the patient carries the code or one of its descendants **and** an age for it can be established* — not simply that the patient carries it.

The referral resource shows the same orientation from the action side. Grouping requested specialties into the families a growth question would reach for accounts for 36,182 of 349,827 referrals (10.3%).

Referrals by growth-relevant specialty family

SPECIALTY FAMILY	REFERRALS	SHARE OF ALL REFERRALS	PATIENTS	MEDIAN AGE
Endocrinology	6,916	1.98%	5,790	9.40 y
Gastroenterology	14,715	4.21%	12,764	5.58 y
Nutrition and dietetics	11,038	3.16%	8,912	9.39 y
Nephrology	1,087	0.31%	937	6.03 y
Genetics	2,426	0.69%	2,116	3.20 y
all other specialties	313,645	89.66%	—	—

Implications for analysis. Use the derived columns when you want an age at first record and are content with the panel upstream chose; go to the raw diagnosis resources for anything else, and match by prefix when you do. These tables describe what the pipeline tracks, not what is clinically relevant to growth in general: a code absent from the panel may still be present in 5.1, and a specialty family here is a string match on a free-text field rather than a clinical taxonomy.

6. Field index

Every column, with its population, range, and the findings that govern it.

6.1 Every column in the extract

All 176 distinct columns across the 8 resources, with how much of each is populated and how many values it takes. Repeated families — the 33 encounter-diagnosis slots and the 8 race slots — appear once each, summarised on their first member.

Field index

RESOURCE	FIELD	TYPE	POPULATED	MISSING	DISTINCT VALUES
patients	patient_id	VARCHAR	250,588	0.0%	250,588
patients	sex	VARCHAR	250,588	0.0%	3
patients	ethnicity	VARCHAR	245,124	2.2%	6
patients	race_1..8	VARCHAR	241,770	3.5%	11
patients_augmented	patient_id	VARCHAR	250,588	0.0%	250,588
patients_augmented	sex	VARCHAR	250,588	0.0%	3
patients_augmented	ethnicity	VARCHAR	199,143	20.5%	2
patients_augmented	race_1..8	VARCHAR	200,533	20.0%	7
patients_augmented	healthy_flag	BIGINT	250,588	0.0%	2
patients_augmented	chronic_dx_flag	BIGINT	250,588	0.0%	2
patients_augmented	growth_dx_flag	BIGINT	250,588	0.0%	2
patients_augmented	ever_stunting_flag	BIGINT	250,588	0.0%	2
patients_augmented	ever_wasting_flag	BIGINT	250,588	0.0%	2
patients_augmented	ever_underweight_flag	BIGINT	250,588	0.0%	2
patients_augmented	ever_obesity_flag	BIGINT	250,588	0.0%	2
patients_augmented	visits_count	BIGINT	250,588	0.0%	173
patients_augmented	visits_count_pre_dx	BIGINT	250,588	0.0%	170
patients_augmented	min_visit_age_days	BIGINT	250,588	0.0%	4,793
patients_augmented	max_visit_age_days	BIGINT	250,588	0.0%	6,559
patients_augmented	visits_span_days	BIGINT	250,588	0.0%	5,635
patients_augmented	dx_age_years	DOUBLE	35,890	85.7%	3,758
patients_augmented	dx_age_years_e03_9	DOUBLE	309	99.9%	283
patients_augmented	dx_age_years_e10	DOUBLE	491	99.8%	468
patients_augmented	dx_age_years_e22_0	DOUBLE	3	100.0%	3

RESOURCE	FIELD	TYPE	POPULATED	MISSING	DISTINCT VALUES
patients_augmented	dx_age_years_e23_0	DOUBLE	150	99.9%	148
patients_augmented	dx_age_years_e23_6	DOUBLE	31	100.0%	31
patients_augmented	dx_age_years_e24	DOUBLE	1	100.0%	1
patients_augmented	dx_age_years_e30_0	DOUBLE	419	99.8%	365
patients_augmented	dx_age_years_e30_1	DOUBLE	3,405	98.6%	1,969
patients_augmented	dx_age_years_e34_3	DOUBLE	447	99.8%	418
patients_augmented	dx_age_years_e34_4	DOUBLE	83	100.0%	82
patients_augmented	dx_age_years_e72_11	DOUBLE	1	100.0%	1
patients_augmented	dx_age_years_k50	DOUBLE	113	100.0%	113
patients_augmented	dx_age_years_k51	DOUBLE	62	100.0%	60
patients_augmented	dx_age_years_k90_0	DOUBLE	898	99.6%	799
patients_augmented	dx_age_years_n18	DOUBLE	70	100.0%	69
patients_augmented	dx_age_years_n25_0	DOUBLE	1	100.0%	1
patients_augmented	dx_age_years_p04_3	DOUBLE	53	100.0%	41
patients_augmented	dx_age_years_p05	DOUBLE	4,069	98.4%	384
patients_augmented	dx_age_years_p07	DOUBLE	11,014	95.6%	973
patients_augmented	dx_age_years_p70	DOUBLE	3,353	98.7%	96
patients_augmented	dx_age_years_p92_6	DOUBLE	14,428	94.2%	222
patients_augmented	dx_age_years_q77	DOUBLE	15	100.0%	14
patients_augmented	dx_age_years_q78_0	DOUBLE	10	100.0%	10
patients_augmented	dx_age_years_q78_1	DOUBLE	2	100.0%	2
patients_augmented	dx_age_years_q87_1	DOUBLE	58	100.0%	57
patients_augmented	dx_age_years_q87_2	DOUBLE	32	100.0%	31
patients_augmented	dx_age_years_q87_3	DOUBLE	46	100.0%	45
patients_augmented	dx_age_years_q87_4	DOUBLE	17	100.0%	17
patients_augmented	dx_age_years_q90	DOUBLE	205	99.9%	109
patients_augmented	dx_age_years_q96	DOUBLE	36	100.0%	27
patients_augmented	dx_age_years_q98_0	DOUBLE	26	100.0%	17
patients_augmented	dx_age_years_q98_4	DOUBLE	42	100.0%	29
patients_augmented	dx_age_years_q98_5	DOUBLE	17	100.0%	13
patients_augmented	count_weight_z_score	BIGINT	250,588	0.0%	174
patients_augmented	mean_weight_z_score	DOUBLE	250,577	0.0%	45,164
patients_augmented	std_weight_z_score	DOUBLE	249,595	0.4%	15,604
patients_augmented	min_weight_z_score	DOUBLE	250,577	0.0%	47,190
patients_augmented	max_weight_z_score	DOUBLE	250,577	0.0%	47,420
patients_augmented	count_height_z_score	BIGINT	250,588	0.0%	97
patients_augmented	mean_height_z_score	DOUBLE	250,261	0.1%	41,024

RESOURCE	FIELD	TYPE	POPULATED	MISSING	DISTINCT VALUES
patients_augmented	std_height_z_score	DOUBLE	248,172	1.0%	14,332
patients_augmented	min_height_z_score	DOUBLE	250,261	0.1%	42,098
patients_augmented	max_height_z_score	DOUBLE	250,261	0.1%	38,259
patients_augmented	count_bmi_z_score	BIGINT	250,588	0.0%	92
patients_augmented	mean_bmi_z_score	DOUBLE	213,053	15.0%	46,476
patients_augmented	std_bmi_z_score	DOUBLE	199,693	20.3%	15,234
patients_augmented	min_bmi_z_score	DOUBLE	213,053	15.0%	50,618
patients_augmented	max_bmi_z_score	DOUBLE	213,053	15.0%	47,254
patients_augmented	count_head_circ_z_score	BIGINT	250,588	0.0%	30
patients_augmented	mean_head_circ_z_score	DOUBLE	200,584	20.0%	46,033
patients_augmented	std_head_circ_z_score	DOUBLE	188,063	25.0%	23,188
patients_augmented	min_head_circ_z_score	DOUBLE	200,584	20.0%	31,399
patients_augmented	max_head_circ_z_score	DOUBLE	200,584	20.0%	36,528
patients_augmented	count_weight_for_length_z_score	BIGINT	250,588	0.0%	61
patients_augmented	mean_weight_for_length_z_score	DOUBLE	220,449	12.0%	42,565
patients_augmented	std_weight_for_length_z_score	DOUBLE	207,992	17.0%	18,775
patients_augmented	min_weight_for_length_z_score	DOUBLE	220,449	12.0%	42,518
patients_augmented	max_weight_for_length_z_score	DOUBLE	220,449	12.0%	38,600
patients_augmented	count_weight_for_stature_z_score	BIGINT	250,588	0.0%	63
patients_augmented	mean_weight_for_stature_z_score	DOUBLE	215,154	14.1%	45,654
patients_augmented	std_weight_for_stature_z_score	DOUBLE	203,194	18.9%	15,593
patients_augmented	min_weight_for_stature_z_score	DOUBLE	215,154	14.1%	39,544
patients_augmented	max_weight_for_stature_z_score	DOUBLE	215,154	14.1%	38,957
visits	patient_id	VARCHAR	6,494,473	0.0%	250,588
visits	visit_id	VARCHAR	6,494,473	0.0%	6,494,473
visits	age_in_days	BIGINT	6,494,473	0.0%	6,563
visits	encounter_type	VARCHAR	6,494,473	0.0%	45
visits	orig_enc_source_Epic_yn	VARCHAR	6,494,473	0.0%	2
visits	weight_oz	DOUBLE	6,488,028	0.1%	24,694
visits	height_in	DOUBLE	3,509,633	46.0%	4,836
visits	head_circ_cm	DOUBLE	1,635,690	74.8%	2,394
visits	BMI	DOUBLE	3,658,303	43.7%	8,216
visits	bmi_percentile	DOUBLE	2,961,185	54.4%	10,001
visits	enc_diag_1..33	VARCHAR	6,154,801	5.2%	6,892
visits_augmented	patient_id	VARCHAR	6,494,473	0.0%	250,588
visits_augmented	visit_id	VARCHAR	6,494,473	0.0%	6,494,473
visits_augmented	sex	VARCHAR	6,494,473	0.0%	3

RESOURCE	FIELD	TYPE	POPULATED	MISSING	DISTINCT VALUES
visits_augmented	ethnicity	VARCHAR	5,401,217	16.8%	2
visits_augmented	race_1..8	VARCHAR	5,423,318	16.5%	7
visits_augmented	age_in_days	BIGINT	6,494,473	0.0%	6,563
visits_augmented	age_in_months	DOUBLE	6,494,473	0.0%	6,563
visits_augmented	age_in_years	DOUBLE	6,494,473	0.0%	6,563
visits_augmented	weight_oz	DOUBLE	6,488,028	0.1%	24,694
visits_augmented	weight_kg	DOUBLE	6,483,007	0.2%	20,665
visits_augmented	weight_outlier_flag	BIGINT	6,488,028	0.1%	2
visits_augmented	delta_weight_kg	DOUBLE	5,754,032	11.4%	5,530
visits_augmented	delta_age_in_days_weight	BIGINT	5,754,032	11.4%	3,424
visits_augmented	weight_velocity	DOUBLE	5,754,032	11.4%	6,449
visits_augmented	weight_z_score	DOUBLE	6,482,932	0.2%	81,802
visits_augmented	weight_percentile	DOUBLE	6,482,932	0.2%	10,001
visits_augmented	weight_for_length_z_score	DOUBLE	2,027,317	68.8%	70,036
visits_augmented	weight_for_length_percentile	DOUBLE	2,027,317	68.8%	10,001
visits_augmented	weight_for_stature_z_score	DOUBLE	1,371,347	78.9%	62,664
visits_augmented	weight_for_stature_percentile	DOUBLE	1,371,347	78.9%	10,001
visits_augmented	wasting_flag	BIGINT	6,494,473	0.0%	2
visits_augmented	height_in	DOUBLE	3,509,633	46.0%	4,836
visits_augmented	height_cm	DOUBLE	3,491,662	46.2%	4,568
visits_augmented	height_outlier_flag	BIGINT	3,509,633	46.0%	2
visits_augmented	delta_height_cm	DOUBLE	2,786,770	57.1%	2,683
visits_augmented	delta_age_in_days_height	BIGINT	2,786,770	57.1%	3,135
visits_augmented	height_velocity	DOUBLE	2,786,770	57.1%	6,653
visits_augmented	height_velocity_z_score	DOUBLE	1,127,289	82.6%	78,625
visits_augmented	height_velocity_z_score_ep	DOUBLE	977,101	85.0%	90,660
visits_augmented	height_velocity_z_score_ap	DOUBLE	960,949	85.2%	78,394
visits_augmented	height_velocity_z_score_lp	DOUBLE	961,074	85.2%	91,509
visits_augmented	height_velocity_percentile	DOUBLE	1,127,289	82.6%	10,001
visits_augmented	height_velocity_percentile_ep	DOUBLE	977,101	85.0%	10,001
visits_augmented	height_velocity_percentile_ap	DOUBLE	960,949	85.2%	10,001
visits_augmented	height_velocity_percentile_lp	DOUBLE	961,074	85.2%	10,001
visits_augmented	height_z_score	DOUBLE	3,491,616	46.2%	62,757
visits_augmented	height_percentile	DOUBLE	3,491,616	46.2%	9,988
visits_augmented	stunting_flag	BIGINT	6,494,473	0.0%	2
visits_augmented	head_circ_cm	DOUBLE	1,635,690	74.8%	2,394
visits_augmented	head_circ_z_score	DOUBLE	1,635,640	74.8%	66,115

RESOURCE	FIELD	TYPE	POPULATED	MISSING	DISTINCT VALUES
visits_augmented	head_circ_percentile	DOUBLE	1,635,640	74.8%	10,001
visits_augmented	bmi	DOUBLE	1,955,339	69.9%	380,344
visits_augmented	bmi_z_score	DOUBLE	1,955,337	69.9%	71,202
visits_augmented	bmi_percentile	DOUBLE	1,955,337	69.9%	10,001
visits_augmented	bmi_category	VARCHAR	1,955,337	69.9%	4
visits_augmented	underweight_flag	BIGINT	6,494,473	0.0%	2
visits_augmented	obesity_flag	BIGINT	6,494,473	0.0%	2
visits_augmented	encounter_type	VARCHAR	6,494,473	0.0%	45
visits_augmented	orig_enc_source_Epic_yn	VARCHAR	6,494,473	0.0%	2
visits_augmented	enc_diag_1..33	VARCHAR	6,154,801	5.2%	6,892
labs	patient_id	VARCHAR	17,230,681	0.0%	247,271
labs	visit_id	VARCHAR	17,229,876	0.0%	2,859,084
labs	lab_order_id	VARCHAR	17,230,681	0.0%	6,578,838
labs	result_line_num	BIGINT	14,947,495	13.3%	149
labs	lab_order_date_age_in_days	BIGINT	17,230,681	0.0%	6,578
labs	lab_procedure_name	VARCHAR	17,230,681	0.0%	3,742
labs	lab_procedure_description	VARCHAR	17,230,681	0.0%	18,834
labs	lab_result_date_age_in_days	BIGINT	14,947,495	13.3%	6,613
labs	result_component_name	VARCHAR	14,947,495	13.3%	12,902
labs	result_loinc_code	VARCHAR	1,350,102	92.2%	2,194
labs	result_value	VARCHAR	14,736,420	14.5%	104,312
labs	result_flag	VARCHAR	1,679,696	90.3%	35
medications	patient_id	VARCHAR	3,823,049	0.0%	236,323
medications	visit_id	VARCHAR	3,823,049	0.0%	2,757,560
medications	med_record_id	VARCHAR	3,823,049	0.0%	3,823,049
medications	med_order_date_age_in_days	BIGINT	3,823,049	0.0%	6,605
medications	med_start_date_age_in_days	BIGINT	3,539,983	7.4%	6,644
medications	med_end_date_age_in_days	BIGINT	3,372,709	11.8%	6,939
medications	med_record_type	VARCHAR	3,823,049	0.0%	2
medications	med_simple_generic_name	VARCHAR	3,823,049	0.0%	1,073
problem_list	patient_id	VARCHAR	1,709,584	0.0%	238,823
problem_list	problem_list_id	VARCHAR	1,709,584	0.0%	1,709,584
problem_list	noted_date_age_in_days	BIGINT	1,702,300	0.4%	6,930
problem_list	resolved_date_age_in_days	BIGINT	757,907	55.7%	6,565
problem_list	pl_diag	VARCHAR	1,709,584	0.0%	4,739
referrals	patient_id	VARCHAR	349,827	0.0%	138,071
referrals	visit_id	VARCHAR	324,997	7.1%	298,615

RESOURCE	FIELD	TYPE	POPULATED	MISSING	DISTINCT VALUES
referrals	referral_id	VARCHAR	349,827	0.0%	349,827
referrals	referral_date_age_in_days	BIGINT	349,827	0.0%	6,535
referrals	requested_specialty	VARCHAR	322,375	7.8%	119
referrals	referral_number_of_visits	BIGINT	323,226	7.6%	6

7. Artifact catalogue

One row per known artifact, with its scale and whether it can be repaired.

7.1 Every artifact this report measured

One row per artifact, gathered from the findings that measured them. The class says who produced the artifact, which decides whether it can be repaired: a derivation artifact can be recomputed without touching the clinical record, a capture artifact cannot, a selection artifact is outside the extract entirely. 15 artifacts across 4 classes (capture, derivation, linkage, selection).

Artifact catalogue

ARTIFACT	CLASS	SCALE IN THIS SNAPSHOT	RECOVERABLE?	SECTION
Raw and augmented BMI disagree on infants	derivation	1,703,005 visits carry a raw BMI the augmented layer withholds; 536 differ outright	Yes — pick the layer deliberately and state which	1.3
Cohort selected on growth-measurement density and code rarity	selection	437,996 registry members reduced to 250,588; 61% of diagnosis codes, 56% of medications and 72% of lab procedures removed with their patients	No — the excluded patients are not in this extract	1.4
A patient-day can carry more than one visit	capture	5,478 patient-days holding 11,040 visits	Partly — define an explicit tie rule before ordering by age	3.1
Populated visit_id that resolves to no visit	linkage	up to 42% of populated values in a resource	No — treat visit linkage as partial by design	3.2
Age fields that violate their own ordering	capture	lab result before order, and medication start before order	No — do not treat differences between them as durations	3.3
Laboratory results are semi-structured text	capture	487,168 comparator-prefixed values; only 44.2% of rows parse as a number	Yes — parse comparators explicitly rather than casting	3.6
Anthropometrics recorded on encounters with no physical contact	capture	weight present on 99% of 22,053 telephone encounters	Partly — restrict by encounter type before counting measurement occasions	3.7
Two measurements of one channel on one patient-day that disagree	capture	942 patient-days for height, median spread 3.17 cm	Partly — define an explicit tie rule before ordering by age	3.8
Diagnosis codes counted flat rather than as a hierarchy	capture	1,204 of 1,327 categories never appear as a bare three-character code	Yes — match on a prefix, or roll up before counting	3.9
Terminal-digit heaping on the imperial recording grid	capture	80.0% of heights fall on a quarter inch	No — it is the precision the measurement actually has	4.2
Wrong-unit and decimal-place entry in the typed measurement fields	capture	1,371 whole-foot heights, 143 centimetre values in the inch field, and a weight decimal artifact enriched 17-fold	Yes — bound and repair the raw imperial columns before converting	4.4

ARTIFACT	CLASS	SCALE IN THIS SNAPSHOT	RECOVERABLE?	SECTION
Apparent height loss from the recording grid on a flat trajectory	capture	0.66% of pairs over a year apart, falling to 0.083% at ages 2 to 10	Not a defect — do not filter it as an outlier	4.5
Height z-score truncated above at +3 while the lower tail runs to -5	derivation	21 visits at or above +3 where roughly 15,800 would be expected	Yes — recompute from the retained raw height	4.6
Head circumference passed through an inch-to-centimetre conversion a second time	derivation	13,467 visits, 90% of all out-of-range values	Yes — divide by 2.54 before applying a plausible range, rather than deleting	4.7
Velocity computed over an age-dependent minimum interval, not between adjacent visits	derivation	99.99% reproduced under the interval rule against 43.7% under a naive lag	Not a defect — carry the interval rule alongside the field	4.8

8. Methods and limitations

How these figures were computed and what would invalidate them.

8.1 Methods, determinism, and limitations

Computation. Every figure was computed with DuckDB against the typed bundle of `ppoc-pediatric-ehr` 1.0.0, snapshot `2026-08-24`, sha256 `425c6f873cefc149344570561a03b33c69a6a6af7fa18bc777c0429579507116`, opened read-only. The bundle is never copied into this repository and no row-level identifier is read into any output.

Privacy. Output is aggregate only. Cells backed by fewer than 10 records are suppressed centrally rather than probe by probe, so a new probe inherits the rule without having to remember it.

Reproducibility. The generator computes the finding set once and renders every output from it, so the HTML, the PDF, the Markdown mirror, and `findings.json` cannot disagree. Prose carries templates rather than literals: a number reaches an output only by way of the finding that measured it. Outputs are rewritten only when the finding set changes, so rebuilding an unchanged snapshot leaves the committed files untouched.

Limitations. Everything here is specific to this snapshot and would need recomputing for another extract. The report describes recording and derivation behaviour, not clinical truth: a value being implausible does not establish what the child actually measured, and a value being plausible does not establish that it was measured at all. Where a mechanism is inferred rather than observed the report says so and shows the evidence.