

CHARGE NUMBER: 2501
PROJECT TITLE: Nuclear and Radiochemistry of Smoke
PROJECT LEADER: Roger A. Comes
PERIOD COVERED: December 1-31, 1981
DATE OF REPORT: January 8, 1982

¹⁴C-Menthol Study¹⁻⁵

A. Section data indicate a leveling off of the migration of the ¹⁴C-menthol into the filters after approximately 20 weeks. Equilibrium has also apparently been reached for the spiked 2R1 cigarettes although the activity is not uniform down the rod. The filtered cigarettes had no activity applied to the filters but now show, after 25 weeks of sealed individual storage, filter activities of: Merit 85mm-73% of that activity applied to the rod; Va. Slim-65%; B&H Lights-66%; Merit Ultra Lights-77%. The section data will continue to be obtained for several more months to insure maximum migration.

Five total smoke distribution runs have been completed on the Merit 85mm ¹⁴C-menthol spiked cigarettes. Average values for the five runs are shown. A slight butt build-up (2-4%) during smoking is indicated.

85mm Merit-¹⁴C-Distribution

| | |
|-------------------------|-------|
| Sidestream Gas | 10.0% |
| Mainstream Gas | 0.3 |
| Sidestream TPM | 5.9 |
| Mainstream TPM | 2.8 |
| Butt (filter + tipping) | 81.0 |
| Ash | 0.1 |

GLRC data on the small mainstream gas phase contribution showed the bulk of the activity to be in the CO₂, ethene, propene and butene peaks. Of that activity found on the MSTPM pad, greater than 99.3% has been identified as ¹⁴C-menthol. These smoke runs, carried out on a single port smoking machine, verify the small butt build-up evidenced in the total smoke recovery machine data.

0001401757

B. Uniformly labelled ¹⁴C-menthol containing cigarettes from a previous study were investigated for menthol delivery on a puff-by-puff basis. Evaluation