

Example activity collection

Bart Snapp

This document was typeset on May 19, 2014.

Contents


Contents

1 First example

In this activity we see some examples.

To start we can have theorem environments:

Theorem 1 *Given a right triangle:*




We have that:

$$a^2 + b^2 = c^2$$

Exercise 2 *Given that $r(v) = -2v^2 - 4v - 4$, evaluate $r(-0.4)$. Express your answer in decimal notation.*

Question 3 *What is the worst kind of cat?*

Question 4 *In the plot below, is P a function of k ?*



Use the plot to compute $P(2)$.


2 Second example

In this activity we give a second example.

Here we have a multi-part question with free-response.

Question 1 Suppose you are standing on a bridge that is 60 meters above sea-level. You toss a ball up into the air with an initial velocity of 30 meters per second. If t is the time (in seconds) after we toss the ball, then the height at time t is approximately $f(t) = -5t^2 + 30t + 60$. What does $f(2)$ mean in our context? Now suppose t is such that $f(t) = 100$. What does this mean in our context? Finally, if h is a small positive value what is the meaning of $f(t + h)$? How does this compare to the meaning of $f(t) + h$?

Here is a picture of a llama:



If you like, check out this video¹.

Exploration 2 Write a Python script that will compute factorial for you.

¹YouTube link: <http://www.youtube.com/watch?v=0aQpLSu2fMs>