

**IX Международная конференция
«Лазерные, плазменные исследования и технологии» ЛаПлаз-2023
Секция "Ускорители заряженных частиц и радиационные технологии"**

**Спиновая Когерентность и Бетатронная Хроматичность
Дейтронного Пучка в Режиме Квазизамороженного Спина.**



Докладчик: Колокольчиков С. (kolokolchikov@inr.ru)

Со-авторы: Сеничев Ю., Аксентьев А., Мельников А.,

Институт Ядерных Исследований РАН, Москва, Россия



Москва, 29 марта 2023 г.



TABLE OF CONTENTS

- ❑ «Quasi-Frozen Spin»
- ❑ Spin Tune Decoherence Effects
 - *Equilibrium Level Energy Shift*
 - *Orbit Lengthening and Betatron Chromaticity*
- ❑ Sextupole Correction
 - *Betatron Chromaticity*
 - *Spin Coherence*
 - α_1/η_1 *Correction*
- ❑ Conclusions



«Quasi-Frozen Spin»

T-BMT Equations

$$\begin{aligned}\frac{d\vec{S}}{dt} &= \vec{S} \times (\vec{\Omega}_{MDM} + \vec{\Omega}_{EDM}), \\ \vec{\Omega}_{MDM} &= \frac{q}{m\gamma} \left\{ (\gamma G + 1) \vec{B}_{\perp} + (G + 1) \vec{B}_{\parallel} - \left(\gamma G + \frac{\gamma}{\gamma + 1} \right) \frac{\vec{\beta} \times \vec{E}}{c} \right\}, \\ \vec{\Omega}_{EDM} &= \frac{q\eta}{2m} \left(\vec{\beta} \times \vec{B} + \frac{\vec{E}}{c} \right), \quad G = \frac{g - 2}{2},\end{aligned}$$

Spin-tunes in E, B fields

$$\begin{aligned}v_s^B &= \gamma G \\ v_s^E &= \frac{G + 1}{\gamma} - G\gamma\end{aligned}$$

Optimization in ByPass NICA Storage Ring



Spin Tune Decoherence Effects

Equilibrium Level Energy Shift

$$\gamma_{eff} = \gamma_s + \beta_s^2 \gamma_s \Delta\delta_{eq}$$


$$\Delta\delta_{eq} = \frac{\gamma_s^2}{\gamma_s^2 \alpha_0 - 1} \left[\frac{\delta_0^2}{2} \left(\alpha_1 + \frac{3\beta_s^2}{2\gamma_s^2} - \frac{\alpha_0}{\gamma_s^2} + \frac{1}{\gamma_s^4} \right) + \left(\frac{\Delta L}{L} \right)_\beta \right], \quad \left(\frac{\Delta L}{L} \right)_\beta = -\frac{\pi}{L_0} [\varepsilon_x v_x + \varepsilon_y v_y]$$

Orbit Lengthening and Betatron Chromaticity




$$\Delta C_\Sigma = -\pi(\varepsilon_x \xi_x + \varepsilon_y \xi_y) + \delta_0(\alpha_0 + \alpha_1 \delta_0 + \dots)$$






Sextupole Correction






Betatron Chromaticity






- 2 sextupole families
- Zero betatron chromaticity
- No spin coherence






Spin Coherence



- 3 sextupole families
- Non-zero betatron chromaticity
- Spin coherence achieved



α_1/η_1 Correction



- 3 sextupole families
- Zero betatron chromaticity
- No spin coherence
- α_1/η_1 up to zero value

$$\gamma_{eff} = \gamma_s + \beta_s^2 \gamma_s \Delta\delta_{eq}$$

$$\Delta\delta_{eq} = \frac{\gamma_s^2}{\gamma_s^2 \alpha_0 - 1} \left[\frac{\delta_0^2}{2} \left(\alpha_1 + \frac{3\beta_s^2}{2\gamma_s^2} - \frac{\alpha_0}{\gamma_s^2} + \frac{1}{\gamma_s^4} \right) + \left(\frac{\Delta L}{L} \right)_\beta \right],$$

$$\left(\frac{\Delta L}{L} \right)_\beta = -\frac{\pi}{L_0} [\varepsilon_x \nu_x + \varepsilon_y \nu_y]$$

$$\Delta C_\Sigma = -\pi(\varepsilon_x \xi_x + \varepsilon_y \xi_y) + \delta_0(\alpha_0 + \alpha_1 \delta_0 + \dots)$$



Optimization Summary Table

Optimization	No optimization	Chromaticity	Spin Coherence	Chromaticity + α_1	Chromaticity + η_1
Tunes	-17/-17	0/0	-13/-18	0/0	0/0
α_1	0.2	-0.4	$-0.37 \cdot 10^{-2}$	$\sim -10^{-12}$	-0.85
quadK_x	$-0.16 \cdot 10^{-1}$	$0.55 \cdot 10^{-1}$	$0.27 \cdot 10^{-13}$	$0.55 \cdot 10^{-1}$	$0.56 \cdot 10^{-1}$
quadK_y	$0.51 \cdot 10^{-2}$	$0.76 \cdot 10^{-1}$	$-0.12 \cdot 10^{-12}$	$0.78 \cdot 10^{-1}$	$0.78 \cdot 10^{-1}$
quadK_z	$-0.43 \cdot 10^{-1}$	$0.20 \cdot 10^{-1}$	$0.13 \cdot 10^{-12}$	$0.13 \cdot 10^{-1}$	$1.6 \cdot 10^{-1}$
Sextupole families	No sextupoles	2	3	3	3
Max. sextupole coefficient, m⁻³	—	2.7	19.4	4.9	104.2



Conclusions

- Considered the phenomenon of spin decoherence simultaneously with betatron chromaticity in ByPass NICA Storage Ring.
- Different cases of sextupole optimizations.
- 3 sextupole families in regular structure can't compensate both betatron chromaticities and get spin coherence.
- It can be possible to modulate dispersion function in such way to get now 3 linear independent sextupole families.
- One of the possible problem decisions is using cooled beam at the level of $dp/p \sim 10^{-5}$. This can help to minimize γ -effective.
- Maximum value of sextupole coefficient not satisfactory and can cause non-linear instabilities.

