


Metodología E — ventana2_pronostico

BOGOTA D.C. — ARROZ PARA SECO (Arroz de primera) | predicción dinámica/recursiva | banda: IC 95%

$t=-0.0207$ | $\beta.=-0.112$ ($t=-1.81$) | $\beta.=-0.162$ ($t=-0.78$) | $p_{\text{sym}}=0.793$

EG — Tipo 1 (sin cte/tend): $p=0.100$ (no coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)




Metodología E — ventana2_pronostico

BOGOTA D.C. — CEBOLLA CABEZONA (Cebolla cabeza blanca) | predicción dinámica/recursiva | banda: IC 95%

$t=-0.1450$ | $\beta.=-0.195$ ($t=-2.81$) | $\beta.=-0.407$ ($t=-2.11$) | $p_{\text{sym}}=0.250$

EG — Tipo 1 (sin cte/tend): $p=0.010$ (coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)




Metodología E — ventana2_pronostico

BOGOTA D.C. — PAPA (Papa R-12 negra) | predicción dinámica/recursiva | banda: IC 95%

$t=-0.0923$ | $\beta.=-0.174$ ($t=-2.81$) | $\beta.=-0.288$ ($t=-1.79$) | $p_{\text{sym}}=0.452$

EG — Tipo 1 (sin cte/tend): $p=0.010$ (coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)




Metodología E — ventana2_pronostico

BOGOTA D.C. — PAPA (Papa parda pastusa) | predicción dinámica/recursiva | banda: IC 95%

$t=-0.0879$ | $\beta.=-0.171$ ($t=-2.81$) | $\beta.=-0.300$ ($t=-1.92$) | $p_{\text{sym}}=0.387$

EG — Tipo 1 (sin cte/tend): $p=0.010$ (coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)




Metodología E — ventana2_pronostico

BOGOTA D.C. — PAPA (Papa suprema) | predicción dinámica/recursiva | banda: IC 95%

$t=-0.0915$ | $\beta.=-0.170$ ($t=-2.84$) | $\beta.=-0.297$ ($t=-1.91$) | $p_{\text{sym}}=0.393$

EG — Tipo 1 (sin cte/tend): $p=0.010$ (coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)




Metodología E — ventana2_pronostico

BOGOTA D.C. — PAPA (Papa única) | predicción dinámica/recursiva | banda: IC 95%

$t=-0.0936$ | $\beta.=-0.164$ ($t=-2.74$) | $\beta.=-0.289$ ($t=-1.90$) | $p_{\text{sym}}=0.393$

EG — Tipo 1 (sin cte/tend): $p=0.010$ (coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)




Metodología E — ventana2_pronostico

BOGOTA D.C. — TOMATE (Tomate chonto) | predicción dinámica/recursiva | banda: IC 95%

$t=-0.0412$ | $\beta.=-0.334$ ($t=-1.96$) | $\beta.=-0.664$ ($t=-3.69$) | $p_{\text{sym}}=0.142$

EG — Tipo 1 (sin cte/tend): $p=0.010$ (coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)




Metodología E — ventana2_pronostico

BOGOTA D.C. — TOMATE (Tomate larga vida) | predicción dinámica/recursiva | banda: IC 95%

$t=-0.0593$ | $\beta.=-0.369$ ($t=-2.20$) | $\beta.=-0.752$ ($t=-4.00$) | $p_{\text{sym}}=0.084$

EG — Tipo 1 (sin cte/tend): $p=0.010$ (coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)



Metodología E — ventana2_pronostico

BOGOTA D.C. — ZANAHORIA (Zanahoria) | predicción dinámica/recursiva | banda: IC 95%

$t=0.1609$ | $\beta.=-0.605$ ($t=-2.96$) | $\beta.=-0.280$ ($t=-3.01$) | $p_{\text{sym}}=0.126$

EG — Tipo 1 (sin cte/tend): $p=0.016$ (coint.) | Tipo 2 (con cte): $p=0.100$ (no coint.) |

Tipo 3 (cte+tend): $p=0.100$ (no coint.)

