

# **MATE 6540: Topology Qualifying Exam**

Due on 19 de mayo

*Prof. Iván Cardona* , C41, 19 de mayo

**Sergio Rodríguez**

### Problem 0

A topological space  $(X, \mathcal{T}_X)$  is pseudocompact  $\iff$  every continuous function  $f : (X, \mathcal{T}_X) \rightarrow (\mathbb{R}, \mathcal{T}_{\varepsilon^1})$  is bounded. Here  $\mathcal{T}_{\varepsilon^1}$  is the usual topology over  $\mathbb{R}$ .

- (i) Show that pseudocompactness is a continuous invariant. Explain.
- (ii) Show that if  $(Y, \mathcal{T}_Y)$  is compact, then  $(Y, \mathcal{T}_Y)$  is pseudocompact, but that the converse does not hold.

### Problem 1

(Kuratowski's closure operator) Let  $X$  be a set,  $\mathcal{P}(X)$  be its powerset, and  $c : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$  be a function that satisfies:

- (i)  $c(\emptyset) = \emptyset$
- (ii)  $A \subseteq c(A), \quad \forall A \in \mathcal{P}(X)$
- (iii)  $c(c(A)) = c(A), \quad \forall A \in \mathcal{P}(X)$
- (iv)  $c(A \cup B) = c(A) \cup c(B), \quad \forall A, B \in \mathcal{P}(X)$

Show that the collection  $\mathcal{T} = \{X \setminus c(A) \mid A \in \mathcal{P}(X)\}$  is a topology over  $X$ , and that in this topology  $\overline{A} = c(A), \quad \forall A \in \mathcal{P}(X)$ . Here  $\overline{A}$  is the closure of  $A$  in  $(X, \mathcal{T})$ .

### Problem 2

Let  $A$  be a subset of a topological space  $(X, \mathcal{T}_X)$ . Show that the following are equivalent:

- (i)  $\text{int}(\overline{A}) = \emptyset$ .
- (ii)  $X \setminus \overline{A}$  is dense in  $X$ .
- (iii)  $X \setminus \overline{(X \setminus \overline{A})} = \emptyset$ .
- (iv)  $A \subseteq \overline{(X \setminus \overline{A})}$ .

### Problem 3

A subset of a topological space is a  $G_\delta$ -set if it is the intersection of countably many open sets. On the other hand, a subset of a topological space is an  $F_\sigma$ -set if it is the union of countably many closed sets.

- (i) Let  $A$  be an  $F_\sigma$ -set of a topological space  $(X, \mathcal{T}_X)$ . Show that there is a nested sequence of closed sets  $C_1 \subseteq C_2 \subseteq C_3 \subseteq \dots$  such that  $A = \bigcup_{i=1}^{\infty} C_i$ .
- (ii) Show that every closed set in a metric space  $(X, d)$  is a  $G_\delta$ -set.

### Problem 4

Let  $A, B$  be two non-empty subsets of  $\mathbb{R}$  with the usual topology. Define:

$$C := \{x + y \mid x \in A \wedge y \in B\}. \quad (1)$$

(a) Show that, if  $A$  or  $B$  is open, then  $C$  is open.

(b) Show that, if  $A$  and  $B$  are compact, then  $C$  is compact.

### Problem 5

(Intermediate value theorem) Let  $f : (X, \mathcal{T}_X) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathbb{R}})$  be a continuous function, where  $(X, \mathcal{T}_X)$  is connected. Show that if  $a, b$  are two points in  $X$  and if  $r$  is a real number lying between  $f(a)$  and  $f(b)$ , then there is a  $c \in X$  such that  $f(c) = r$ .

### Problem 6

Let  $(X_1, \mathcal{T}_1)$  and  $(X_2, \mathcal{T}_2)$  be topological spaces and suppose  $X_1 \times X_2$  has the product topology. For each  $i = 1, 2$ , let  $A_i \subseteq X_i$ . Prove that:

(i)  $\overline{A_1 \times A_2} = \overline{A_1} \times \overline{A_2}$ .

(ii)  $\text{int}(A_1 \times A_2) = \text{int}(A_1) \times \text{int}(A_2)$ .