

Project: Image Completion and Inpainting

Week 1: Image Inpainting

Coordinator: Juan Fco. Garamendi

Lecturers: J.F. Garamendi, Vadim Fedorov

Universitat Pompeu Fabra

<jf.garamendi, vadim.fedorov>@upf.edu

Optimization and Inference Techniques for Computer Vision



Master in
Computer Vision
Barcelona

Oct. 6th. 2016

What is Inpainting?



Detail of "Cornelia, Mother of the Gracchi" by J. Suvee (Louvre). Courtesy of Emile-Male "The Restorer's Handbook of easel painting".

What is Inpainting?



What is Inpainting?



What is Inpainting?



What is not Inpainting?



Goal

How is education supposed to make me feel smarter? Besides, every time I learn something new, it pushes some old stuff out of my brain. Remember when I took that home winemaking course, and I forgot how to drive?



Inpainting using Laplace's Equation

Let $f : \Omega \rightarrow \mathbb{R}$ be a given grayscale image and let $D \subset \Omega$ be an open set representing the region to be inpainted.

It is supposed that f is known in $\Omega \setminus D := \{x \in \Omega : x \notin D\}$

The inpainting solution u can be found as the minimum of

$$\left\{ \begin{array}{l} \arg \min_{u \in W^{1,2}(\Omega)} \int_D |\nabla u(x)|^2 dx, \\ u|_{\partial D} = f \end{array} \right.$$

and satisfies the Laplace's equation

$$\left\{ \begin{array}{ll} \Delta u &= 0 \quad \text{in } D, \\ u &= f \quad \text{in } \partial D \end{array} \right.$$

where Δ denotes the Laplacian and $u = f$ in $\Omega \setminus D$

The equation is completed with homogeneous Neumann boundary conditions at the boundary of the image.

Inpainting using Laplace's Equation

Let $f : \Omega \rightarrow \mathbb{R}$ be a given grayscale image and let $D \subset \Omega$ be an open set representing the region to be inpainted.

It is supposed that f is known in $\Omega \setminus D := \{x \in \Omega : x \notin D\}$

The inpainting solution u can be found as the minimum of

$$\left\{ \begin{array}{l} \arg \min_{u \in W^{1,2}(\Omega)} \int_D |\nabla u(x)|^2 dx, \\ u|_{\partial D} = f \end{array} \right.$$

and satisfies the Laplace's equation

$$\left\{ \begin{array}{ll} \Delta u &= 0 \quad \text{in } D, \\ u &= f \quad \text{in } \partial D \end{array} \right.$$

where Δ denotes the Laplacian and $u = f$ in $\Omega \setminus D$

The equation is completed with homogeneous Neumann boundary conditions at the boundary of the image.

Inpainting using Laplace's Equation

Let $f : \Omega \rightarrow \mathbb{R}$ be a given grayscale image and let $D \subset \Omega$ be an open set representing the region to be inpainted.

It is supposed that f is known in $\Omega \setminus D := \{x \in \Omega : x \notin D\}$

The inpainting solution u can be found as the minimum of

$$\left\{ \begin{array}{l} \arg \min_{u \in W^{1,2}(\Omega)} \int_D |\nabla u(x)|^2 dx, \\ u|_{\partial D} = f \end{array} \right.$$

and satisfies the Laplace's equation

$$\left\{ \begin{array}{ll} \Delta u &= 0 & \text{in } D, \\ u &= f & \text{in } \partial D \end{array} \right.$$

where Δ denotes the Laplacian and $u = f$ in $\Omega \setminus D$

The equation is completed with homogeneous Neumann boundary conditions at the boundary of the image.

Inpainting using Laplace's Equation

Let $f : \Omega \rightarrow \mathbb{R}$ be a given grayscale image and let $D \subset \Omega$ be an open set representing the region to be inpainted.

It is supposed that f is known in $\Omega \setminus D := \{x \in \Omega : x \notin D\}$

The inpainting solution u can be found as the minimum of

$$\left\{ \begin{array}{l} \arg \min_{u \in W^{1,2}(\Omega)} \int_D |\nabla u(x)|^2 dx, \\ u|_{\partial D} = f \end{array} \right.$$

and satisfies the Laplace's equation

$$\left\{ \begin{array}{ll} \Delta u &= 0 \quad \text{in } D, \\ u &= f \quad \text{in } \partial D \end{array} \right.$$

where Δ denotes the Laplacian and $u = f$ in $\Omega \setminus D$

The equation is completed with homogeneous Neumann boundary conditions at the boundary of the image.

Inpainting using Laplace's Equation

Laplacian of u

$$\Delta u = \operatorname{div}(\nabla u)$$

Spatial Gradient

Forward differences

$$(\nabla u)_{i,j} = (u_{x_{i,j}}, u_{y_{i,j}})$$

with

$$u_{x_{i,j}} = \frac{u_{i+1,j} - u_{i,j}}{h_i}$$

$$u_{y_{i,j}} = \frac{u_{i,j+1} - u_{i,j}}{h_j}$$

Divergence

Backward Differences. Let $\bar{v} = (v_x, v_y)$ be a vector

$$\operatorname{div}(\bar{v})_{i,j} = \frac{(v_{x_{i,j}} - v_{x_{i-1,j}})}{h_i} + \frac{(v_{y_{i,j}} - v_{y_{i,j-1}})}{h_j}$$

Inpainting using Laplace's Equation

Laplacian of u

$$\Delta u = \operatorname{div}(\nabla u)$$

Spatial Gradient

Forward differences

$$(\nabla u)_{i,j} = (u_{x_{i,j}}, \ u_{y_{i,j}})$$

with

$$u_{x_{i,j}} = \frac{u_{i+1,j} - u_{i,j}}{h_i} \quad u_{y_{i,j}} = \frac{u_{i,j+1} - u_{i,j}}{h_j}$$

Inpainting using Laplace's Equation

Laplacian of u

$$\Delta u = \operatorname{div}(\nabla u)$$

Spatial Gradient

Forward differences

$$(\nabla u)_{i,j} = (u_{x_{i,j}}, u_{y_{i,j}})$$

with

$$u_{x_{i,j}} = \frac{u_{i+1,j} - u_{i,j}}{h_i} \quad u_{y_{i,j}} = \frac{u_{i,j+1} - u_{i,j}}{h_j}$$

Divergence

Backward Differences. Let $\bar{v} = (v_x, v_y)$ be a vector

$$\operatorname{div}(\bar{v})_{i,j} = \frac{(v_{x_{i,j}} - v_{x_{i-1,j}})}{h_i} + \frac{(v_{y_{i,j}} - v_{y_{i,j-1}})}{h_j}$$

Mandatory

- Build the matrix A and vector b for the algebraic system of equations obtained after the given discretization of the operators.
- Test with the given images. There is one image (image6.tif) that missed the 99% of information



- Proof that the solution of the Laplace's equation is a solution of the problem

$$\begin{cases} \arg \min_{u \in W^{1,2}(\Omega)} \int_D |\nabla u(x)|^2 dx, \\ u|_{\partial D} = f \end{cases}$$

Mandatory

- Build the matrix A and vector b for the algebraic system of equations obtained after the given discretization of the operators.
- Test with the given images. There is one image (image6.tif) that missed the 99% of information



- Proof that the solution of the Laplace's equation is a solution of the problem

$$\begin{cases} \arg \min_{u \in W^{1,2}(\Omega)} \int_D |\nabla u(x)|^2 dx, \\ u|_{\partial D} = f \end{cases}$$

Mandatory

- Build the matrix A and vector b for the algebraic system of equations obtained after the given discretization of the operators.
- Test with the given images. There is one image (image6.tif) that missed the 99% of information

- Proof that the solution of the Laplace's equation is a solution of the problem

$$\begin{cases} \arg \min_{u \in W^{1,2}(\Omega)} \int_D |\nabla u(x)|^2 dx, \\ u|_{\partial D} = f \end{cases}$$