

1 Compute the values for

1.  $\sum_{i=-1}^4 3$

$$\begin{aligned}\sum_{i=-1}^4 3 &= 3 \cdot (4 + 1 + 1) \\ &= 3 \cdot 6 \\ &= 18\end{aligned}$$

2.  $\sum_{i=1}^5 (\frac{1}{3})^i$

$$\begin{aligned}\sum_{i=1}^5 (\frac{1}{3})^i &= \sum_{i=0}^5 (\frac{1}{3})^i - \sum_{i=0}^0 (\frac{1}{3})^i \\ &= \frac{(\frac{1}{3})^6 - 1}{\frac{1}{3} - 1} - 1\end{aligned}$$

3.  $\sum_{i=1}^n 3$

$$\begin{aligned}\sum_{i=1}^n 3 &= 3 \cdot \sum_{i=1}^n 1 \\ &= 3 \cdot n\end{aligned}$$

4.  $\sum_{i=-3}^n 3$

$$\sum_{i=-3}^n 3 = 3 \cdot (n + 4)$$

5.  $\sum_{k=0}^n 2^k + \sum_{k=5}^n 2^k$

$$\begin{aligned}
\sum_{k=0}^n 2^k + \sum_{k=5}^n 2^k &= \sum_{k=0}^n 2^k + \sum_{k=0}^n 2^k - \sum_{k=0}^4 2^k \\
&= 2 \sum_{k=0}^n 2^k - \sum_{k=0}^4 2^k \\
&= 2 \cdot \frac{2^{n+1} - 1}{2 - 1} - \frac{2^5 - 1}{2 - 1}
\end{aligned}$$

6.  $\sum_{i=0}^n (\frac{2}{3})^i + \sum_{i=-4}^n (\frac{2}{3})^i$

$$\begin{aligned}
\sum_{i=0}^n (\frac{2}{3})^i + \sum_{i=-4}^n (\frac{2}{3})^i &= \frac{(\frac{2}{3})^{n+1} - 1}{\frac{2}{3} - 1} + \sum_{i=-4}^{-1} (\frac{2}{3})^i + \sum_{i=0}^n (\frac{2}{3})^i \\
&= \frac{(\frac{2}{3})^{n+1} - 1}{\frac{2}{3} - 1} + \sum_{i=-4}^{-1} (\frac{2}{3})^i + \frac{(\frac{2}{3})^{n+1} - 1}{\frac{2}{3} - 1} \\
&= \frac{(\frac{2}{3})^{n+1} - 1}{\frac{2}{3} - 1} + \sum_{i=1}^4 (\frac{3}{2})^i + \frac{(\frac{2}{3})^{n+1} - 1}{\frac{2}{3} - 1} \\
&= \frac{(\frac{2}{3})^{n+1} - 1}{\frac{2}{3} - 1} + \frac{(\frac{3}{2})^5 - 1}{(\frac{3}{2}) - 1} + \frac{(\frac{2}{3})^{n+1} - 1}{\frac{2}{3} - 1}
\end{aligned}$$

7.  $\sum_{i=1}^n (i^3 + 2 \cdot i^2 - i + 1)$

$$\begin{aligned}
\sum_{i=1}^n (i^3 + 2 \cdot i^2 - i + 1) &= \sum_{i=1}^n i^3 + 2 \cdot \sum_{i=1}^n i^2 - \sum_{i=1}^n i + \sum_{i=1}^n 1 \\
&= (\frac{n \cdot (n+1)}{2})^2 + 2 \cdot \frac{(n) \cdot (n+1) \cdot (2n+1)}{6} - \frac{n \cdot (n+1)}{2} + n
\end{aligned}$$

8.  $\sum_{i=5}^n (-4 \cdot i + \frac{i}{5})$

$$\begin{aligned}
\sum_{i=5}^n (-4 \cdot i + \frac{i}{5}) &= -4 \sum_{i=5}^n i + \frac{1}{5} \sum_{i=5}^n i \\
&= -4 \sum_{i=0}^n i + \frac{1}{5} \sum_{i=0}^n i + 4 \sum_{i=0}^4 i - \frac{1}{5} \sum_{i=0}^4 i \\
&= -4 \cdot \frac{n(n+1)}{2} + \frac{1}{5} \cdot \frac{n(n+1)}{2} + 4 \cdot \frac{4 \cdot 5}{2} - \frac{1}{5} \cdot \frac{4 \cdot 5}{2}
\end{aligned}$$

$$9. \sum_{j=0}^k \sum_{i=1}^j (i - j^2 - 2)$$

$$\begin{aligned}
\sum_{j=0}^k \sum_{i=1}^j (i - j^2 - 2) &= \sum_{j=0}^k \sum_{i=1}^j i - \sum_{j=0}^k \sum_{i=1}^j j^2 - \sum_{j=0}^k \sum_{i=1}^j 2 \\
&= \sum_{j=0}^k \frac{j(j+1)}{2} - \sum_{j=0}^k j^3 - \sum_{j=0}^k 2 \cdot j \\
&= \sum_{j=0}^k \frac{j^2}{2} + \sum_{j=0}^k \frac{j}{2} - (\frac{k \cdot (k+1)}{2})^2 - 2 \cdot \frac{k \cdot (k+1)}{2} \\
&= \frac{1}{2} \cdot \frac{(k) \cdot (k+1) \cdot (2k+1)}{6} + \frac{1}{2} \cdot \frac{k \cdot (k+1)}{2} - (\frac{k \cdot (k+1)}{2})^2 - 2 \cdot \frac{k \cdot (k+1)}{2}
\end{aligned}$$

$$10. \sum_{j=1}^m \sum_{k=1}^j (3 \cdot C + k - 3 \cdot j + i)$$

$$\begin{aligned}
\sum_{j=1}^m \sum_{k=1}^j (3 \cdot C + k - 3 \cdot j + i) &= \sum_{j=1}^m \sum_{k=1}^j (3 \cdot C + k - 3 \cdot j + i) \\
&= \sum_{j=1}^m \sum_{k=1}^j 3 \cdot C + \sum_{j=1}^m \sum_{k=1}^j k - 3 \sum_{j=1}^m \sum_{k=1}^j j + \sum_{j=1}^m \sum_{k=1}^j i \\
&= \sum_{j=1}^m 3 \cdot C \cdot j + \sum_{j=1}^m \frac{j(j+1)}{2} - 3 \sum_{j=1}^m j^2 + \sum_{j=1}^m i \cdot j \\
&= 3 \cdot C \cdot \frac{m \cdot (m+1)}{2} + \sum_{j=1}^m \frac{j^2}{2} + \sum_{j=1}^m \frac{j}{2} - 3 \cdot \frac{(m)(m+1)(2m+1)}{6} \\
&\quad + i \cdot \frac{(m)(m+1)}{2} \\
&= 3 \cdot C \frac{m \cdot (m+1)}{2} + \frac{1}{2} \cdot \frac{m(m+1)(2m+1)}{6} + \frac{1}{2} \cdot \frac{(m)(m+1)}{2} \\
&\quad - 3 \cdot \frac{(m)(m+1)(2m+1)}{6} + i \cdot \frac{(m)(m+1)}{2}
\end{aligned}$$

11.  $\sum_{l=-4}^n \sum_{j=1}^k \sum_{i=1}^j (i-4)$

$$\begin{aligned}
\sum_{l=-4}^n \sum_{j=1}^k \sum_{i=1}^j (i-4) &= \sum_{l=-4}^n \sum_{j=1}^k \sum_{i=1}^j i - 4 \sum_{l=-4}^n \sum_{j=1}^k \sum_{i=1}^j 1 \\
&= \sum_{l=-4}^n \sum_{j=1}^k \frac{j(j+1)}{2} - \sum_{l=-4}^n \sum_{j=1}^k j \\
&= \sum_{l=-4}^n \sum_{j=1}^k \frac{j^2}{2} + \sum_{l=-4}^n \sum_{j=1}^k \frac{j}{2} - \sum_{l=-4}^n \frac{k(k+1)}{2} \\
&= \sum_{l=-4}^n \frac{(k)(k+1)(2k+1)}{12} + \sum_{l=-4}^n \frac{(k)(k+1)}{4} - \sum_{l=-4}^n \frac{k(k+1)}{2} \\
&= (n+5) \frac{(k)(k+1)(2k+1)}{12} + (n+5) \frac{(k)(k+1)}{4} - (n+5) \frac{(k)(k+1)}{2}
\end{aligned}$$

2. Calculate the answer

1.  $\log_4 x = 5 \rightarrow x = ?$

$$\begin{aligned}
\log_4 x &= 5 \\
x &= 4^5
\end{aligned}$$

$$2. \log_3 y = 4 \rightarrow y = ?$$

$$\begin{aligned} \log_3 y &= 4 \\ y &= 3^4 \end{aligned}$$

$$3. x = 7^2 \rightarrow \log_7 x = ?$$

$$\begin{aligned} x &= 7^2 \\ \log_7 x &= \log_7 7^2 \\ \log_7 x &= 2 \cdot \log_7 7 \\ \log_7 x &= 2 \end{aligned}$$

$$4. x = 32 \rightarrow \log_2 x = ?$$

$$\begin{aligned} x &= 32 \\ \log_2 x &= \log_2 32 \\ \log_2 x &= 5 \end{aligned}$$

$$5. 2^{\log_2 5} + 4^{\log_2 6} - 27^{\log_3 5}$$

$$\begin{aligned} 2^{\log_2 5} + 4^{\log_2 6} - 27^{\log_3 5} &= 5 + 2^{2^{\log_2 6}} - 3^{3^{\log_3 5}} \\ &= 5 + 2^{\log_2 6^2} - 3^{\log_3 5^3} \\ &= 5 + 6^2 - 5^3 \\ &= 5 + 36 - 125 \\ &= -84 \end{aligned}$$

$$6. 9^{\log_3 2} - 25^{\log_5 4} - 36^{\log_6 7} + 8^{\log_8 6}$$

$$\begin{aligned} 9^{\log_3 2} - 25^{\log_5 4} - 36^{\log_6 7} + 8^{\log_8 6} &= 3^{2^{\log_3 2}} - 5^{2^{\log_5 4}} - 6^{2^{\log_6 7}} + 6 \\ &= 3^{\log_3 2^2} - 5^{\log_5 4^2} - 6^{\log_6 7^2} + 6 \\ &= \log_3(2^2) - 5^4 - 49 + 6 \\ &= 2 \cdot \log_3(2) - 5^4 - 49 + 6 \\ &= 2 \cdot \frac{\log_2(2)}{\log_2(3)} - 5^4 - 49 + 6 \\ &= \frac{2}{1.5} - 5^4 - 49 + 6 \end{aligned}$$

$$7. \log(4^5 \times 8^3) - \log(16 - 8) + \log\left(\frac{2^{10}}{4 \times 3^2}\right)$$

$$\begin{aligned} \log_2(4^5 \times 8^3) - \log_2(16 - 8) + \log_2\left(\frac{2^{10}}{4 \times 3^2}\right) &= \log_2(4^5) + \log_2(8^3) - \log_2(8) + \log_2(2^{10}) - \log_2(4 \times 3^2) \\ &= 5 \cdot \log_2(4) + 3 \cdot \log_2(8) - \log_2(8) + 10 \cdot \log_2(2) - \log_2(4) \\ &= 5 \cdot 2 + 3 \cdot 3 - 3 + 10 - 2 - 2 \cdot 1.5 \\ &= 21 \end{aligned}$$

$$8. \log(3^2 \times 64^3) - \log\left(\frac{2^{10} \times 128^3}{9 \times 8^2}\right)$$

$$\begin{aligned} \log_2(3^2 \times 64^3) - \log_2\left(\frac{2^{10} \times 128^3}{9 \times 8^2}\right) &= \log_2(3^2) + \log_2(64^3) - \log_2(2^{10} \times 128^3) + \log_2(9) + \log_2 8^2 \\ &= 2 \cdot \log_2(3) + 3 \cdot \log_2(64) - 10 \cdot \log_2 2 - 3 \cdot \log_2 128 + 2 \cdot \log_2 3 + 2 \cdot \\ &= 2 \cdot 1.5 + 3 \cdot 6 - 10 - 3 \cdot 7 + 2 \cdot 1.5 + 2 \cdot 3 \\ &= -1 \end{aligned}$$

$$9. \log \log 16$$

$$\begin{aligned} \log_2 \log_2 16 &= \log_2 4 \\ &= 2 \end{aligned}$$

$$10. \log 16 \times \log 16$$

$$\begin{aligned} \log_2 16 \times \log_2 16 &= 4 \times 4 \\ &= 16 \end{aligned}$$

$$11. \log^2 16$$

$$\begin{aligned} \log_2^2 16 &= 4^2 \\ &= 16 \end{aligned}$$

$$12. \log_2 \log_5 625 - \log_3 \log_4 2^{3^9} + \log^4 2^5 - \frac{\log^2(4^3 \times 3^5)}{\log_5 125}$$

$$\begin{aligned}
\log_2 \log_5 625 - \log_3 \log_4 2^{3^9} + \log_2^4 2^5 - \frac{\log^2(4^3 \times 3^5)}{\log_5 125} &= \log_2 4 - \log_3(3^9 \cdot \log_4 2) + \log_2^4 2^5 - \frac{\log_2^2(4^3 \times 3^5)}{\log_5 125} \\
&= 2 - 9 \log_3\left(\frac{3}{2}\right) + (5 \cdot \log_2(2))^4 - \frac{(\log_2(4^3 \times 3^5))^2}{3} \\
&= 2 - 9 + 9 \cdot \log_3\left(\frac{1}{2}\right) + 5^4 - \frac{(3 \cdot 2 + 5 \cdot \log_2(3))^2}{3} \\
&= 2 - 9 + 9 \cdot \frac{-1}{1.5} + 5^4 - \frac{(3 \cdot 2 + 5 \cdot (1.5)^2)^2}{3}
\end{aligned}$$

13.  $\log \log_8 \log 256 + \log^5(3^2) \times 4^{\log 7}$

$$\begin{aligned}
\log_2 \log_8 \log_2 256 + \log_2^5(3^2) \cdot 4^{\log_2 7} &= \log_2 \log_8 8 + \log_2^5(3^2) \cdot 4^{\log_2 7} \\
&= \log_2 1 + \log_2^5(3^2) \cdot 4^{\log_2 7} \\
&= \log_2^5(3^2) \cdot 4^{\log_2 7} \\
&= (2 \cdot 1.5)^5 \cdot 2^{\log_2 49} \\
&= (2 \cdot 1.5)^5 \cdot 49 \\
&= 11907
\end{aligned}$$

14.  $\log_6 x = 5 \rightarrow \log_x 6 = ?$

$$\begin{aligned}
\log_6 x &= 5 \\
x &= 6^5 \\
\log_x x &= 5 \cdot \log_x 6 \\
1 &= 5 \cdot \log_x 6 \\
\frac{1}{5} &= \log_x 6
\end{aligned}$$

15.  $\log_y x = 10 \rightarrow \log_x y = ?$

$$\begin{aligned}
\log_y x &= 10 \\
x &= y^{10} \\
\log_x x &= 10 \cdot \log_x y \\
\frac{1}{10} &= \log_x y
\end{aligned}$$

$$16. \log_4 32 - \log_8^2 4$$

$$\begin{aligned} \log_4 32 - \log_8^2 4 &= \log_4 2 \cdot 16 - \log_8^2 4 \\ &= \log_4 2 + \log_4 16 - \log_8^2 4 \\ &= \frac{1}{2} + 2 - \left(\frac{2}{3}\right)^2 \\ &= \frac{1}{2} + 2 - \frac{4}{9} \end{aligned}$$

$$17. \log_4 8 + \log_9 27 - \log_{25}^2 125 - \log_8^3 16 + \log_4 \log_2 256$$

$$\begin{aligned} \log_4 8 + \log_9 27 - \log_{25}^2 125 - \log_8^3 16 + \log_4 \log_2 256 &= \frac{3}{2} + \frac{3}{2} - \left(\frac{3}{2}\right)^2 - \left(\frac{4}{3}\right)^3 + \log_4(8) \\ &= \frac{3}{2} + \frac{3}{2} - \left(\frac{9}{4}\right) - \left(\frac{64}{27}\right) + \frac{3}{2} \end{aligned}$$

3. Compute the derivative of

$$1. -5 \cdot x^3 + 2 \cdot x - 1$$

$$\frac{d}{dx}(-5 \cdot x^3 + 2 \cdot x - 1) = -15 \cdot x^2 + 2$$

$$2. 3 \cdot x^4 - 2\sqrt{x} + x^{\frac{1}{2}} - 6x^{-\frac{2}{3}} - 5$$

$$\begin{aligned} \frac{d}{dx}(3 \cdot x^4 - 2\sqrt{x} + x^{\frac{1}{2}} - 6x^{-\frac{2}{3}} - 5) &= 12 \cdot x^3 - x^{-\frac{1}{2}} + \frac{1}{2} \cdot x^{-\frac{1}{2}} + 4 \cdot x^{-\frac{5}{3}} \\ &= 12 \cdot x^3 - \frac{1}{2}x^{-\frac{1}{2}} + 4 \cdot x^{-\frac{5}{3}} \end{aligned}$$

$$3. x \cdot \sqrt{x} + \sqrt{\sqrt{x}}$$

$$\frac{d}{dx}(x^{\frac{3}{2}} + x^{\frac{1}{4}}) = \frac{3}{2}x^{\frac{1}{2}} + \frac{1}{4} \cdot x^{-\frac{3}{4}}$$



4.  $\log x - x^2 \ln x + \ln x^4$

$$\frac{d}{dx}(\log_2 x - x^2 \cdot \ln x + \ln x^4) = \frac{1}{x \cdot \ln 2} - 2 \cdot x \cdot \ln x - x + \frac{4}{x}$$

5.  $\ln^3(x\sqrt{2x-3}) + \sqrt{\ln x^2}$

$$\begin{aligned} \frac{d}{dx}(\ln^3(x \cdot \sqrt{2x-3}) + \sqrt{\ln x^2}) &= \frac{3 \cdot (\ln(x\sqrt{2x-3}))^2 \cdot (\sqrt{2x-3} + x \cdot (2 \cdot x - 3)^{-\frac{1}{2}})}{x\sqrt{2x-3}} \\ &+ \frac{1}{x \cdot \sqrt{\ln(x^2)}} \end{aligned}$$

6.  $\frac{\sqrt[4]{x+5} - \ln x}{(x-1)^3}$

$$\frac{d}{dx}\left(\frac{(x+5)^{\frac{1}{4}} - \ln x}{(x-1)^3}\right) = \frac{(x-1)^3 \cdot \left(\frac{1}{4} \cdot (x+5)^{-\frac{3}{4}} - \frac{1}{x}\right) - 3 \cdot ((x+5)^{\frac{1}{4}} - \ln x) \cdot (x-1)^2}{(x-1)^6}$$

4. Determine the limit of

1.  $\lim_{x \rightarrow \infty} \frac{3x+2}{-5x-6}$

$$\lim_{x \rightarrow \infty} \frac{3x+2}{-5x-6} = -\frac{3}{5}$$

2.  $\lim_{x \rightarrow \infty} \left(\frac{1}{x} + 3\right)$

$$\begin{aligned} \lim_{x \rightarrow \infty} \left(\frac{1}{x} + 3\right) &= \lim_{x \rightarrow \infty} \frac{1}{x} + \lim_{x \rightarrow \infty} 3 \\ &= \lim_{x \rightarrow \infty} 3 \\ &= 3 \end{aligned}$$

3.  $\lim_{x \rightarrow \infty} \frac{x^3 + x - \sqrt{3x}}{\sqrt{x}}$

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x^3 + x - \sqrt{3x}}{\sqrt{x}} &= \lim_{x \rightarrow \infty} \frac{x^3}{\sqrt{x}} + \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x}} - \lim_{x \rightarrow \infty} \frac{\sqrt{3x}}{\sqrt{x}} \\ &= \infty \end{aligned}$$

$$4. \lim_{x \rightarrow \infty} \frac{x^3 + x - \sqrt{3x}}{5 \cdot x^{2.25} \cdot \sqrt{\sqrt{x}}}$$

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x^3 + x - \sqrt{3x}}{5 \cdot x^{2.25} x^{.25}} &= \lim_{x \rightarrow \infty} \frac{x^3 + x - \sqrt{x}}{5 \cdot x^3} \\ &= \lim_{x \rightarrow \infty} \frac{x^3}{5 \cdot x^3} + \lim_{x \rightarrow \infty} \frac{x}{5 \cdot x^3} - \lim_{x \rightarrow \infty} \frac{\sqrt{x}}{5 \cdot x^3} \\ &= \frac{1}{5} \end{aligned}$$

$$5. \lim_{x \rightarrow \infty} \frac{x^{0.1} - \sqrt{3}}{\sqrt{\sqrt{x}}}$$

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x^{0.1} - \sqrt{3}}{x^{0.25}} &= \lim_{x \rightarrow \infty} \frac{x^{0.1}}{x^{0.25}} - \frac{\sqrt{3}}{x^{0.25}} \\ &= \lim_{x \rightarrow \infty} \frac{1}{x^{0.15}} - \frac{\sqrt{3}}{x^{0.25}} \\ &= 0 \end{aligned}$$

$$6. \lim_{x \rightarrow \infty} \frac{x^x}{2^x}$$

$$\begin{aligned} \ln(\lim_{x \rightarrow \infty} \frac{x^x}{2^x}) &= \lim_{x \rightarrow \infty} \ln(\frac{x^x}{2^x}) \\ &= \lim_{x \rightarrow \infty} \ln(x^x) - \lim_{x \rightarrow \infty} \ln(2^x) \\ &= \lim_{x \rightarrow \infty} x \ln(x) - \lim_{x \rightarrow \infty} x \ln(2) \\ &= \infty \\ e^{\ln(\lim_{x \rightarrow \infty} \frac{x^x}{2^x})} &= e^\infty \\ &= \infty \end{aligned}$$

$$7. \lim_{x \rightarrow \infty} \frac{x^x}{x(2^x)}$$

$$\begin{aligned} \ln(\lim_{x \rightarrow \infty} \frac{x^x}{x(2^x)}) &= \lim_{x \rightarrow \infty} \ln(\frac{x^x}{x \cdot 2^x}) \\ &= \lim_{x \rightarrow \infty} \ln(x^x) - \lim_{x \rightarrow \infty} \ln(x \cdot 2^x) \\ &= \lim_{x \rightarrow \infty} x \ln(x) - \lim_{x \rightarrow \infty} \ln(x) - \lim_{x \rightarrow \infty} \ln(2^x) \\ &= \lim_{x \rightarrow \infty} x \ln(x) - \lim_{x \rightarrow \infty} \ln(x) - \lim_{x \rightarrow \infty} x \ln(2) \\ &= \infty \\ e^{\ln(\lim_{x \rightarrow \infty} \frac{x^x}{x(2^x)})} &= e^\infty \\ &= \infty \end{aligned}$$

$$8. \lim_{x \rightarrow \infty} \frac{\sqrt{2}^{\log^4 x^3}}{\log(2 \cdot x + 7)}$$

$$9. \lim_{x \rightarrow \infty} \frac{x+1}{\frac{3 \cdot x \ln x}{2x^2}}$$

$$10. \lim_{x \rightarrow \infty} \frac{\sqrt{2}^{\log x^3}}{\log \ln x(2x)}$$

5. Compute the exact values for

$$1. \int_1^n (2 \cdot x^4 + 5\sqrt{x}) \cdot dx$$

$$\begin{aligned} \int_1^n (2 \cdot x^4 + 5\sqrt{x}) \cdot dx &= 2 \int_1^n x^4 dx + 5 \int_1^n \sqrt{x} dx \\ &= 2 \cdot \left( \frac{x^5}{5} \Big|_1^n \right) + 5 \left( \frac{2}{3} x^{\frac{3}{2}} \Big|_1^n \right) \\ &= 2 \cdot \left( \frac{n^5}{5} - \frac{1}{5} \right) + 5 \left( \frac{2}{3} n^{\frac{3}{2}} - \frac{2}{3} \right) \\ &= \frac{2 \cdot n^5}{5} - \frac{2}{5} + \frac{10}{3} n^{\frac{3}{2}} - \frac{10}{3} \end{aligned}$$

$$2. \int_1^n (x^4 - 3 \cdot x^2 + \frac{1}{x} - \frac{1}{x^2}) dx$$

$$\begin{aligned} \int_1^n (x^4 - 3 \cdot x^2 + \frac{1}{x} - \frac{1}{x^2}) dx &= \int_1^n x^4 dx - 3 \int_1^n x^2 dx + \int_1^n \frac{1}{x} dx - \int_1^n \frac{1}{x^2} dx \\ &= \frac{x^5}{5} \Big|_1^n - 3 \cdot \frac{x^3}{3} \Big|_1^n + \ln x \Big|_1^n + \frac{1}{x} \Big|_1^n \\ &= \left( \frac{n^5}{5} - \frac{1}{5} \right) - n^3 + 1 + \ln n + \frac{1}{n} - 1 \\ &= \frac{n^5}{5} - n^3 + \ln n + \frac{1}{n} - \frac{1}{5} \end{aligned}$$

$$3. \int_1^n \left( \frac{3}{\sqrt{x}} + \ln x + e^x \right) dx$$

$$\begin{aligned} \int_1^n \frac{3}{\sqrt{x}} + \ln x + e^x dx &= \int_1^n \frac{3}{\sqrt{x}} dx + \int_1^n \ln x dx + \int_1^n e^x dx \\ &= 6\sqrt{x} \Big|_1^n + (x \ln x - x) \Big|_1^n + e^n - e \\ &= 6\sqrt{n} - 6 + (n \ln n - n + 1) + e^n - e \\ &= 6\sqrt{n} + n \ln n - n - 5 + e^n - e \end{aligned}$$

$$4. \int_1^n x \cdot \sin x dx$$

$$\begin{aligned} \int_1^n x \cdot \sin x dx &= -x \cos x \Big|_1^n + \int_1^n \cos x dx \\ &= -n \cos n + \cos(1) + \sin(n) - \sin(1) \end{aligned}$$

6. Use mathematical induction to prove that

$$1 + 2 + \dots + n = \frac{n(n+1)}{2}$$

*Proof.*  $1 + 2 + \dots + n = \frac{n(n+1)}{2}$

Base case  $n = 1$ : If  $n = 1$  then the left hand side and the right hand size is  $1 = 1 = \frac{1(2)}{2}$ .

Inductive hypothesis: Suppose the theorem holds for all values of  $n$  up to some  $k, k \geq 1$ .

Inductive step: let  $n = k + 1$  then our left hand side is

$$\begin{aligned} \sum_{i=1}^{k+1} i &= \sum_{i=1}^k i + (k+1) \\ &= \frac{k(k+1)}{2} + (k+1) \\ &= \frac{k(k+1) + 2 \cdot k + 2}{2} \\ &= \frac{(k+1) \cdot (k+2)}{2} \end{aligned}$$

which is equal to our right hand side. By the principle of mathematical induction, the theorem holds for all integers  $n \geq 1$ .  $\square$

7. Use mathematical induction to prove that

$$1 + 2^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$1 + 2 + \dots + n = \frac{n(n+1)}{2}$$

*Proof.* Base case  $n = 1$ : If  $n = 1$  then the left hand side and the right hand size is  $1^2 = 1 = \frac{1(2)(3)}{6}$ .

Inductive hypothesis: Suppose the theorem holds for all values of  $n$  up to some  $k, k \geq 1$ .

Inductive step: let  $n = k + 1$  then our left hand side is

$$\begin{aligned}
\sum_{i=1}^{k+1} i^2 &= \sum_{i=1}^k i^2 + (k+1)^2 \\
&= \frac{k \cdot (k+1) \cdot (2 \cdot k + 1)}{6} + (k+1)^2 \\
&= \frac{k \cdot (k+1) \cdot (2 \cdot k + 1) + 6 \cdot (k+1)^2}{6} \\
&= \frac{(6 \cdot (k+1) + k \cdot (2 \cdot k + 1)) \cdot (k+1)}{6} \\
&= \frac{(6 \cdot k + 6 + 2 \cdot k^2 + k) \cdot (k+1)}{6} \\
&= \frac{(2 \cdot k^2 + 7 \cdot k + 6) \cdot (k+1)}{6} \\
&= \frac{(2 \cdot k^2 + 4 \cdot k + 3 \cdot k + 6) \cdot (k+1)}{6} \\
&= \frac{(2 \cdot k \cdot (k+2) + 3 \cdot (k+2)) \cdot (k+1)}{6} \\
&= \frac{(2 \cdot k + 3) \cdot (k+2) \cdot (k+1)}{6}
\end{aligned}$$

which is equal to our right hand side. By the principle of mathematical induction, the theorem holds for all integers  $n \geq 1$ .  $\square$