

1 Compute the values for

1. $\sum_{i=-1}^4 3$

$$\begin{aligned}\sum_{i=-1}^4 3 &= 3 + 3 + 3 + 3 + 3 + 3 \\ &= 3 \cdot 6 \\ &= 18\end{aligned}$$

2. $\sum_{i=1}^5 (\frac{1}{3})^i$

$$\begin{aligned}\sum_{i=1}^5 (\frac{1}{3})^i &= (\frac{1}{3})^1 + (\frac{1}{3})^2 + (\frac{1}{3})^3 + (\frac{1}{3})^4 + (\frac{1}{3})^5 \\ &= \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \frac{1}{81} + \frac{1}{243} \\ &= \frac{121}{243}\end{aligned}$$

3. $\sum_{i=1}^n 3$

$$\begin{aligned}\sum_{i=1}^n 3 &= 3 \cdot \sum_{i=1}^n 1 \\ &= \frac{3 \cdot n \cdot (n+1)}{2}\end{aligned}$$

4. $\sum_{i=-3}^n 3$

$$\begin{aligned}\sum_{i=-3}^n 3 &= \sum_{i=-3}^0 3 + \sum_{i=1}^n 3 \\ &= 3 \cdot 4 + \frac{3 \cdot n \cdot (n+1)}{2} \\ &= \frac{3 \cdot n^2 + 3 \cdot n + 24}{2}\end{aligned}$$

5. $\sum_{k=0}^n 2^k + \sum_{k=5}^n 2^k$

$$\begin{aligned}
\sum_{k=0}^n 2^k + \sum_{k=5}^n 2^k &= 2^{n+1} - 1 + \sum_{k=5}^n 2^k \\
&= 2^{n+1} - 1 + \sum_{k=0}^n 2^k - \sum_{k=0}^4 2^k \\
&= 2^{n+1} - 1 + 2^{n+1} - 1 - (2^5 - 1) \\
&= 2 \cdot 2^{n+1} - 2 - (31) \\
&= 2 \cdot 2^{n+1} - 33 \\
&= 2^{n+2} - 33
\end{aligned}$$

$$6. \sum_{i=0}^n \left(\frac{2}{3}\right)^i + \sum_{i=-4}^n \left(\frac{2}{3}\right)^i$$

$$\begin{aligned}
\sum_{i=0}^n \left(\frac{2}{3}\right)^i + \sum_{i=-4}^n \left(\frac{2}{3}\right)^i &= \sum_{i=0}^n \left(\frac{2}{3}\right)^i + \sum_{i=-4}^{-1} \left(\frac{2}{3}\right)^i + \sum_{i=0}^n \left(\frac{2}{3}\right)^i \\
&= 2 \cdot \sum_{i=0}^n \left(\frac{2}{3}\right)^i + \sum_{i=-4}^{-1} \left(\frac{2}{3}\right)^i \\
&= 2 \cdot \frac{1}{1 - \frac{2}{3}} + \sum_{i=-4}^{-1} \left(\frac{2}{3}\right)^i \\
&= \frac{2}{\frac{1}{3}} + \sum_{i=-4}^{-1} \left(\frac{2}{3}\right)^i \\
&= 6 + \left(\frac{2}{3}\right)^{-4} + \left(\frac{2}{3}\right)^{-3} + \left(\frac{2}{3}\right)^{-2} + \left(\frac{2}{3}\right)^{-1} \\
&= \frac{291}{16}
\end{aligned}$$

$$7. \sum_{i=1}^n (i^3 + 2 \cdot i^2 - i + 1)$$

$$\begin{aligned}
\sum_{i=1}^n (i^3 + 2 \cdot i^2 - i + 1) &= \sum_{i=1}^n i^3 + \sum_{i=1}^n 2 \cdot i^2 - \sum_{i=1}^n i + \sum_{i=1}^n 1 \\
&= \left(\frac{n(n+1)}{2}\right)^2 + 2 \cdot \frac{n(n+1)(2n+1)}{6} - \frac{n(n+1)}{2} + n \\
&= \left(\frac{n(n+1)}{2}\right)^2 + \frac{n(n+1)(2n+1)}{3} - \frac{n(n+1)}{2} + n
\end{aligned}$$

$$8. \sum_{i=5}^n (-4 \cdot i + \frac{i}{5})$$

$$\begin{aligned}
\sum_{i=5}^n (-4 \cdot i + \frac{i}{5}) &= \sum_{i=5}^n -4 \cdot i + \sum_{i=5}^n \frac{i}{5} \\
&= -4 \cdot \sum_{i=5}^n i + \frac{1}{5} \cdot \sum_{i=5}^n i \\
&= -4 \cdot \sum_{i=1}^n i + \frac{1}{5} \cdot \sum_{i=1}^n i + 4 \cdot \sum_{i=1}^4 i - \frac{1}{5} \cdot \sum_{i=1}^4 i \\
&= -4 \cdot \frac{n(n+1)}{2} + \frac{1}{5} \cdot \frac{n(n+1)}{2} + 4 \cdot \frac{4(5)}{2} - \frac{1}{5} \cdot \frac{4(5)}{2} \\
&= -4 \cdot \frac{n(n+1)}{2} + \frac{n(n+1)}{10} + 40 - 2 \\
&= -4 \cdot \frac{n(n+1)}{2} + \frac{n(n+1)}{10} + 38
\end{aligned}$$

$$9. \sum_{j=0}^k \sum_{i=1}^j (i - j^2 - 2)$$

$$\begin{aligned}
\sum_{j=0}^k \sum_{i=1}^j (i - j^2 - 2) &= \sum_{j=0}^k \sum_{i=1}^j i - \sum_{j=0}^k \sum_{i=1}^j j^2 - \sum_{j=0}^k \sum_{i=1}^j 2 \\
&= \sum_{j=0}^k \frac{j(j+1)}{2} - \sum_{j=0}^k j^3 - \sum_{j=0}^k 2 \cdot j \\
&= \sum_{j=0}^k \frac{j^2 + j}{2} - \sum_{j=0}^k j^3 - \sum_{j=0}^k 2 \cdot j \\
&= \sum_{j=0}^k \frac{j^2}{2} + \sum_{j=0}^k \frac{j}{2} - \sum_{j=0}^k j^3 - \sum_{j=0}^k 2 \cdot j \\
&= \frac{1}{2} \cdot \sum_{j=0}^k j^2 + \frac{1}{2} \cdot \sum_{j=0}^k j - \sum_{j=0}^k j^3 - 2 \cdot \sum_{j=0}^k j \\
&= \frac{n(n+1)(2n+1)}{12} + \frac{n(n+1)}{4} - (\frac{n(n+1)}{2})^2 - n(n+1)
\end{aligned}$$

$$10. \sum_{j=1}^m \sum_{k=1}^j (3 \cdot C + k - 3 \cdot j + i)$$

$$\begin{aligned}
\sum_{j=1}^m \sum_{k=1}^j (3 \cdot C + k - 3 \cdot j + i) &= \sum_{j=1}^m \sum_{k=1}^j 3 \cdot C + \sum_{j=1}^m \sum_{k=1}^j k - \sum_{j=1}^m \sum_{k=1}^j 3 \cdot j + \sum_{j=1}^m \sum_{k=1}^j i \\
&= \sum_{j=1}^m 3 \cdot C \cdot j + \sum_{j=1}^m \frac{j \cdot (j+1)}{2} - \sum_{j=1}^m 3 \cdot j^2 + \sum_{j=1}^m i \cdot j \\
&= 3 \cdot C \frac{m \cdot (m+1)}{2} + \frac{1}{2} \sum_{j=1}^m j^2 + \frac{1}{2} \sum_{j=1}^m j - 3 \cdot \sum_{j=1}^m j^2 + i \cdot \sum_{j=1}^m j \\
&= 3 \cdot C \cdot \frac{m \cdot (m+1)}{2} + \frac{1}{2} \cdot \frac{m \cdot (m+1)(2m+1)}{6} + \frac{1}{2} \cdot \frac{m \cdot (m+1)}{2} - \\
&\quad 3 \cdot \frac{m \cdot (m+1) \cdot (2m+1)}{6} + i \cdot \frac{m(m+1)}{2}
\end{aligned}$$

$$11. \sum_{l=-4}^n \sum_{j=1}^k \sum_{i=1}^j (i-4)$$

$$\begin{aligned}
\sum_{l=-4}^n \sum_{j=1}^k \sum_{i=1}^j (i-4) &= \sum_{l=-4}^n \sum_{j=1}^k \sum_{i=1}^j i - 4 \sum_{l=-4}^n \sum_{j=1}^k \sum_{i=1}^j 1 \\
&= \sum_{l=-4}^n \sum_{j=1}^k \frac{j \cdot (j+1)}{2} - 4 \cdot \sum_{l=-4}^n \cdot \sum_{j=1}^k j \\
&= \frac{1}{2} \cdot \sum_{l=-4}^n \sum_{j=1}^k j^2 + \frac{1}{2} \cdot \sum_{l=-4}^n \sum_{j=1}^k j - 4 \cdot \sum_{l=-4}^n \sum_{j=1}^k j \\
&= \frac{1}{2} \cdot \sum_{l=-4}^n \frac{k(k+1)(2k+1)}{6} + \frac{1}{2} \cdot \sum_{l=-4}^n \frac{k(k+1)}{2} - 4 \cdot \sum_{l=-4}^n \frac{k(k+1)}{2} \\
&= \frac{1}{2} \cdot (n+5) \cdot \frac{k \cdot (k+1) \cdot (2k+1)}{6} + \frac{1}{2} \cdot (n+5) \cdot \frac{k \cdot (k+1)}{2} \\
&\quad - 4 \cdot (n+5) \cdot \frac{k \cdot (k+1)}{2}
\end{aligned}$$

2. Calculate the answer

$$1. \log_4 x = 5 \rightarrow x = ?$$

$$\begin{aligned}
\log_4 x &= 5 \\
x &= 4^5
\end{aligned}$$

$$2. \log_3 y = 4 \rightarrow y = ?$$

$$\begin{aligned}
\log_3 y &= 4 \\
y &= 3^4
\end{aligned}$$

3. $x = 7^2 \rightarrow \log_7 x = ?$

$$\begin{aligned} x &= 7^2 \\ \log_7 x &= \log_7 7^2 \\ \log_7 x &= 2 \cdot \log_7 7 \\ \log_7 x &= 2 \end{aligned}$$

4. $x = 32 \rightarrow \log_2 x = ?$

$$\begin{aligned} x &= 32 \\ \log_2 x &= \log_2 32 \\ \log_2 x &= 5 \end{aligned}$$

5. $2^{\log 5} + 4^{\log 6} - 27^{\log 3 5}$

6. $9^{\log_3 2} - 25^{\log_5 4} - 36^{\log_6 7} + 8^{\log_8 6}$

7. $\log(4^5 \times 8^3) - \log(16 - 8) + \log\left(\frac{2^{10}}{4 \times 3^2}\right)$

8. $\log(3^2 \times 64^3) - \log\left(\frac{2^{10} \times 128^3}{9 \times 8^2}\right)$

9. $\log \log 16$

10. $\log 16 \times \log 16$

11. $\log^2 16$

12. $\log_2 \log_5 625 - \log_3 \log_4 2^{3^9} + \log^4 2^5 - \frac{\log^2(4^3 \times 3^5)}{\log_5 125}$

13. $\log \log_8 \log 256 + \log^5(3^2) \times 4^{\log 7}$

14. $\log_6 x = 5 \rightarrow \log_x 6 = ?$

15. $\log_y x = 10 \rightarrow \log_x y = ?$

16. $\log_4 32 - \log_8^2 4$

17. $\log_4 8 + \log_9 27 - \log_{25}^2 125 - \log_8^3 16 + \log_4 \log 256$

3. Compute the derivative of

1. $-5 \cdot x^3 + 2 \cdot x - 1$

2. $3 \cdot x^4 - 2\sqrt{x} + x^{\frac{1}{2}} - 6x^{-\frac{2}{3}} - 5$

3. $x \cdot \sqrt{x} + \sqrt{\sqrt{x}}$

4. $\log x - x^2 \ln x + \ln x^4$
5. $\ln^3(x\sqrt{2x-3}) + \sqrt{\ln x^2}$
6. $\frac{\sqrt[4]{x+5} - \ln x}{(x-1)^3}$

4. Determine the limit of

1. $\lim_{x \rightarrow \infty} \frac{3x+2}{-5x-6}$
2. $\lim_{x \rightarrow \infty} (\frac{1}{x} + 3)$
3. $\lim_{x \rightarrow \infty} \frac{x^3+x-\sqrt{3x}}{\sqrt{x}}$
4. $\lim_{x \rightarrow \infty} \frac{x^3+x-\sqrt{3x}}{5 \cdot x^{2.25} \cdot \sqrt{\sqrt{x}}}$
5. $\lim_{x \rightarrow \infty} \frac{x^{0.1} - \sqrt{3}}{\sqrt{\sqrt{x}}}$
6. $\lim_{x \rightarrow \infty} \frac{x^x}{2^x}$
7. $\lim_{x \rightarrow \infty} \frac{x^x}{x(2^x)}$
8. $\lim_{x \rightarrow \infty} \frac{\sqrt{2}^{\log^4 x^3}}{\log(2 \cdot x + 7)}$
9. $\lim_{x \rightarrow \infty} \frac{\frac{x+1}{3 \cdot x \ln x}}{\frac{2x^2}{2x^2}}$
10. $\lim_{x \rightarrow \infty} \frac{\sqrt{2}^{\log x^3}}{\log^{\ln x}(2x)}$

5. Compute the exact values for

1. $\int_1^n (2 \cdot x^4 + 5\sqrt{x})$
2. $\int_1^n (x^4 - 3 \cdot x^2 + \frac{1}{x} - \frac{1}{x^2}) dx$
3. $\int_1^n (\frac{3}{\sqrt{x}} + \ln x + e^x) dx$
4. $\int_1^n x \cdot \sin x dx$

6. Use mathematical induction to prove that

$$1 + 2 + \dots + n = \frac{n(n+1)}{2}$$

Proof. $1 + 2 + \dots + n = \frac{n(n+1)}{2}$

Base case $n = 1$: If $n = 1$ then the left hand side and the right hand size is $1 = 1 = \frac{1(2)}{2}$.

Inductive hypothesis: Suppose the theorem holds for all values of n up to some $k, k \geq 1$.

Inductive step: let $n = k + 1$ then our left hand side is

$$\begin{aligned}\sum_{i=1}^{k+1} i &= \sum_{i=1}^k i + (k+1) \\ &= \frac{k(k+1)}{2} + (k+1) \\ &= \frac{k(k+1) + 2 \cdot k + 2}{2} \\ &= \frac{(k+1) \cdot (k+2)}{2}\end{aligned}$$

which is equal to our right hand side. By the principle of mathematical induction, the theorem holds for all integers $n \geq 1$. \square

7. Use mathematical induction to prove that

$$1 + 2^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$1 + 2 + \dots + n = \frac{n(n+1)}{2}$$

Proof. Base case $n = 1$: If $n = 1$ then the left hand side and the right hand size is $1^2 = 1 = \frac{1(2)(3)}{6}$.

Inductive hypothesis: Suppose the theorem holds for all values of n up to some $k, k \geq 1$.

Inductive step: let $n = k + 1$ then our left hand side is

$$\begin{aligned}
\sum_{i=1}^{k+1} i^2 &= \sum_{i=1}^k i^2 + (k+1)^2 \\
&= \frac{k \cdot (k+1) \cdot (2 \cdot k + 1)}{6} + (k+1)^2 \\
&= \frac{k \cdot (k+1) \cdot (2 \cdot k + 1) + 6 \cdot (k+1)^2}{6} \\
&= \frac{(6 \cdot (k+1) + k \cdot (2 \cdot k + 1)) \cdot (k+1)}{6} \\
&= \frac{(6 \cdot k + 6 + 2 \cdot k^2 + k) \cdot (k+1)}{6} \\
&= \frac{(2 \cdot k^2 + 7 \cdot k + 6) \cdot (k+1)}{6} \\
&= \frac{(2 \cdot k^2 + 4 \cdot k + 3 \cdot k + 6) \cdot (k+1)}{6} \\
&= \frac{(2 \cdot k \cdot (k+2) + 3 \cdot (k+2)) \cdot (k+1)}{6} \\
&= \frac{(2 \cdot k + 3) \cdot (k+2) \cdot (k+1)}{6}
\end{aligned}$$

which is equal to our right hand side. By the principle of mathematical induction, the theorem holds for all integers $n \geq 1$. \square