- 1) Consider the perceptron in two dimensions: $h(x) = sign(w^T x)$ where $w = \begin{bmatrix} w_0, w_1, w_2 \end{bmatrix}^T$ and $x = \begin{bmatrix} 1, x_1, x_2 \end{bmatrix}^T$. Technically, x has three coordinates, but we call this perceptron two-dimensional because the first coordinate is fixed at 1.
- (a) Show that the regions on the plane where h(x)=+1 and h(x)=-1 are separated by a line. If we express this line by the equation $x_2=ax_1+b$, what are the slope a and intercept b in terms of w_0,w_1,w_2
- (b) Draw a picture for the cases $w = \begin{bmatrix} 1,\,2,\,3 \end{bmatrix}^T$ and $w = -\begin{bmatrix} 1,\,2,\,3 \end{bmatrix}^T$.

In more than two dimensions, the +1 and -1 regions are separated by a hyperplane, the generalization of a line.

2) Given $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, show that the rank of a matrix xy^T is one.

$$xy^{T} = \begin{bmatrix} x_{1} \\ x_{2} \\ \vdots \\ x_{m} \end{bmatrix} \begin{bmatrix} y_{1}^{T} & y_{2}^{T} & \dots & y_{n}^{T} \end{bmatrix}$$

$$= \begin{bmatrix} x_{1} \cdot y_{1}^{T} & x_{1} \cdot y_{2}^{T} & \dots & x_{1} \cdot y_{n}^{T} \\ x_{2} \cdot y_{1}^{T} & x_{2} \cdot y_{2}^{T} & \dots & x_{2} \cdot y_{n}^{T} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m} \cdot y_{1}^{T} & x_{m} \cdot y_{2}^{T} & \dots & x_{m} \cdot y_{n}^{T} \end{bmatrix}$$

Columns linear combinations of x thus we conclude the rank is one

3) Given $X = [x_1, x_2, \dots, x_n] \in \mathbb{R}^{m \times n}$ where $x_i \in \mathbb{R}^m$ for all i, and $Y^T = [y^1, y^2, \dots, y^n] \in \mathbb{R}^{p \times n}$ where $y^i \in \mathbb{R}^p$ for all i. Show that

$$XY = \sum_{i=1}^{N} x_i(y^i)^T$$

$$\sum_{i=1}^{N} x_i(y^i)^T =$$

4) Given $X \in \mathbb{R}^{m \times n}$, show that the matrix $X^T X$ is symmetric and positive semi-definite. When is it positive definite?

We first show that X^TX is symmetric

$$(X^T X)^T = X^T X^{TT}$$

= $X^T X$

Thus X^TX is symmetric

Next we show X^TX is positive-semidefinite

$$v^{T}(X^{T}X)v = \|Xv\|^{2}$$
$$\|Xv\|^{2} \geq 0 \forall v \in \mathbb{R}^{n}$$

Which shows that the matrix is positive-semidefinite

 X^TX is positive definite iff $\mathcal{N}(X)=\{\hat{0}\}$. Thus the matrix X must be full-column rank

- 5) Given $g(x, y) = e^x + e^{y^2} + e^{3xy}$, compute $\frac{\partial g}{\partial y}$. $\frac{\partial g}{\partial y}g(x,y) = 2ye^{y^2} + 3xe^{3xy}$
- 6) Consider the matrix

$$A = \begin{pmatrix} 2 & 1 & 3 \\ 1 & 1 & 2 \\ 3 & 2 & 5 \end{pmatrix}$$

- (a) Compute the eigenvalues and corresponding eigenvectors of A. You are allowed to use Matlab to compute the eigenvectors
- (b) What is the eigen-decomposition of A?
- (c) What is the rank of A?
- (d) Is A positive definite? Is A positive semi-definite?
- (e) Is A singular?