

Tema lab 8

1. $h(t) = e^{-4|t|}$

$$\begin{aligned} H(j\omega_0) &= \int_{-\infty}^{\infty} h(\tau) \cdot e^{-jk\omega_0\tau} d\tau = \int_{-\infty}^{\infty} e^{-4|\tau|} \cdot e^{-jk\omega_0\tau} d\tau = \\ &= \int_{-\infty}^0 e^{+4\tau} \cdot e^{-jk\omega_0\tau} d\tau + \int_0^{\infty} e^{-4\tau} \cdot e^{-jk\omega_0\tau} d\tau = \\ &= \frac{1}{4-jk\omega_0} \cdot e^{(4-jk\omega_0)\tau} \Big|_{-\infty}^0 + \frac{1}{-4-jk\omega_0} \cdot e^{(4-jk\omega_0)\tau} \Big|_0^{\infty} = \\ &= \frac{1}{4-jk\omega_0} + \frac{1}{4+jk\omega_0} = \frac{4+jk\omega_0-4+jk\omega_0}{16+(k\omega_0)^2} = \frac{2jk\omega_0}{16+(k\omega_0)^2} \end{aligned}$$

Fourier series repr. of the output $y(t)$ for these input signals

a) $x(t) = \cos(2\pi t)$

$\Rightarrow \omega_0 = 2\pi$; $x(t) = \frac{1}{2} (e^{2\pi jt} + e^{-2\pi jt}) = \sum_{k=-1}^1 a_k \cdot e^{jk2\pi t}$

where $a_0 = 0$ and $a_{-1} = a_1 = \frac{1}{2}$

$y(t) = \sum_{k=-1}^1 b_k \cdot e^{jk2\pi t}$, unde $b_0 = a_0 \cdot H(0) = 0$

$b_1 = a_1 \cdot H(\omega_0) = \frac{1}{2} H(2\pi) = \frac{1}{2} \cdot \frac{4\pi j}{16+4\pi^2} = \frac{\pi j}{8(1+\frac{1}{2}\pi^2)}$

$b_{-1} = a_{-1} \cdot H(-\omega_0) = \frac{1}{2} \cdot H(-2\pi) = \frac{1}{2} \cdot \frac{4\pi j}{16+4\pi^2} = \frac{\pi j}{8(1-\frac{1}{2}\pi^2)}$

b) $x(t) = \sum_{n=-\infty}^0 \delta(t-n) \Rightarrow T_0 = 1, \omega_0 = 2\pi$

$a_k = \frac{1}{T_0} \int_{T_0} x(t) \cdot e^{-jk\omega_0 t} dt = \frac{1}{T_0} \int_{-\frac{T_0}{2}}^{\frac{T_0}{2}} \delta(t) \cdot e^{-jk\omega_0 t} dt = \frac{1}{T_0} \cdot e^{-jk\omega_0 t} \Big|_{t=0} = \frac{1}{T_0} = 1$

$x(t) = \sum_{k=-\infty}^{\infty} a_k \cdot e^{jk\omega_0 t} = \frac{1}{T_0} \sum_{k=-\infty}^{\infty} e^{jk\omega_0 t} = \sum_{k=-\infty}^{\infty} e^{jk2\pi t}$

$b_k = a_k \cdot H(2\pi k) = \frac{1}{T_0} H(2\pi k) = \frac{1}{T_0} \cdot \frac{4jk\pi}{16+4\pi^2 k^2} = \frac{4\pi k j}{4+\pi^2 k^2}$

$\Rightarrow y(t) = \sum_{k=-\infty}^{\infty} \frac{\pi k j}{4+\pi^2 k^2} \cdot e^{jk2\pi t}$

$$c) x(t) = \sum_{n=-\infty}^{\infty} (-1)^n \delta(t-n) \Rightarrow T_0 = 2, \omega_0 = \pi$$

$$a_k = \frac{1}{T_0} \int_{T_0} x(t) \cdot e^{-jk\omega_0 t} dt = \frac{1}{T_0} \left(\int_{-\frac{T_0}{2}}^{\frac{T_0}{2}} \delta(t) \cdot e^{-jk\omega_0 t} dt + \int_{\frac{T_0}{2}}^{\frac{3T_0}{2}} -\delta(t - \frac{T_0}{2}) \cdot e^{-jk\omega_0 t} dt \right)$$

$$= \frac{1}{T_0} \left(e^{-jk\omega_0 t} \Big|_{t=0} - e^{-jk\omega_0 t} \Big|_{t=\frac{T_0}{2}} \right) = \frac{1}{T_0} (1 - e^{-jk\omega_0 \cdot \frac{T_0}{2}}) = \frac{1}{T_0} (1 - e^{-jk\pi})$$

$$= \frac{1}{T_0} (1 - (-1)^k) = \frac{1}{2} (1 - (-1)^k)$$

$$x(t) = \sum_{n=-\infty}^{\infty} a_k \cdot e^{jk\omega_0 t} = \frac{1}{2} \sum_{n=-\infty}^{\infty} (1 - (-1)^k) \cdot e^{jk\pi t}$$


$$b_k = a_k \cdot H(k\omega_0) = \frac{1 - (-1)^k}{T_0} \cdot H(k\pi) = \frac{1 - (-1)^k}{T_0} \cdot \frac{2k\pi j}{16 + (k\pi)^2}$$

$$\Rightarrow b_k = \begin{cases} \frac{2k\pi j}{16 + (k\pi)^2}, & \text{for } k\text{-odd} \\ 0, & \text{for } k\text{-even} \end{cases}$$

$$y(t) = \sum_{n=-\infty}^{\infty} b_k \cdot e^{jk\pi t}$$

```
clear variables
```


```
T0 = 1; w0 = 2*pi/T0;  
Ts = 0.0005; t = -1:Ts:2;  
x = t - 0.1 - round(t);  
figure; plot(t,x);
```



```
N = 8;  
ak = fsAnalysis(x, t, T0, Ts, N);
```


```
%Pentru Ts = 0.01:
```

```
T0 = 1; w0 = 2*pi/T0;  
Ts1 = 0.01; t = -1:Ts1:2;  
x1 = t - 0.1 - round(t);  
figure; plot(t,x1);
```



```
N = 3;
ak = fsAnalysis(x1, t, T0, Ts1, N);
```

```
%Pentru Ts = 0.001:
T0 = 1; w0 = 2*pi/T0;
Ts2 = 0.001; t = -1:Ts2:2;
x2 = t - 0.1 - round(t);
figure; plot(t,x2);
```



```

N = 10;
ak = fsAnalysis(x2, t, T0, Ts2, N);
a0 = ak(N+1);
ksi0 = 0;
AK = abs(ak(N+2:end));
ksi = angle(ak(N+2:end));

x_est = a0*ones(size(t));
for k = 1:N
x_est = x_est + 2*AK(k)*cos(w0*k*t+ksi(k));
end

figure; plot(t,x,'b',t,x_est,'r');

```

Error using plot
 Vectors must be the same length.

```

xlabel('t');legend('x(t)','x_{est}(t)');

n0 = 0:N;
figure, subplot(3,1,1);stem(n0,[a0, AK],'.');
legend('Amplitude spectrum')
subplot(3,1,2),stem(n0,[ksi0, ksi],'.g')
legend('Phase spectrum')
subplot(3,1,3),stem(n0,[a0^2, (AK.^2)/2],'.r')
label('n'), legend('Power spectrum');

```

```

function ak = fsAnalysis(x, t, T0, Ts, N)
% function for estimating the first N coefficients of the Fourier series for signal x
% T0 - fundamental period of x; Ts - sampling period

% extraction of one period from x
    t = t(1:floor(T0/Ts));
    x = x(1:length(t));

% estimation of the coefficients using the trapezoidal method
    w0 = 2*pi/T0;
    ak = [];
    for k = -N:N
        ak = [ak, (1/T0)*trapz(t, x.*exp(-j*k*w0*t))];
    end
end

```