

Acknowledgements

Publications

The presented thesis is a compendium of the following research articles:

- **Sergi Valverde**, Eloy Roura, Arnau Oliver, and Xavier Lladó. MSSEG: a Multiple Sclerosis brain tissue SEGmentation method . *Submitted to Medical Image analysis*. 2016.
- **Sergi Valverde**, Arnau Oliver, Eloy Roura, Deborah Pareto, Joan Carles Vilanova, Lluís Ramió-Torrentà, Jaume Sastre-Garriga, Xavier Montalban, Àlex Rovira and Xavier Lladó. Quantifying brain tissue volume in multiple sclerosis with automated lesion segmentation and filling SPM8 toolboxes. *NeuroImage Clinical*. 2016. Elsevier. [JCR N IF 2.526, Q2(6/14)].
- **Sergi Valverde**, Arnau Oliver, Yago Díez, Mariano Cabezas, Joan Carles Vilanova, Lluís Ramió-Torrentà, Àlex Rovira, and Xavier Lladó. Evaluating the effects of white matter multiple sclerosis lesions on the volume estimation of six brain tissue segmentation methods. *American Journal of Neuroradiology*. Vol. 36(6), pp. 1109-1115. 2015. American Society of Neuroradiology. [JCR RNMMI IF 3.589, Q1(19/125)].
- **Sergi Valverde**, Arnau Oliver, Mariano Cabezas, Eloy Roura and Xavier Lladó. Comparison of ten brain tissue segmentation methods using revisited IBSR annotations. *Journal of Magnetic Resonance Imaging*. Vol. 41, Issue 1, pp. 93-101. January 2015. DOI: 10.1002/jmri.24517. Wiley. [JCR RNMMI IF: 3.210 Q1(23/125)].
- **Sergi Valverde**, Arnau Oliver, and Xavier Lladó. A white matter lesion-filling approach to improve brain tissue volume measurements. *NeuroImage: Clinical*. Vol. 6, pp 86-92. 2014. Elsevier. [JCR N IF 2.526, Q2(6/14)].

The rest of publications derived from this PhD thesis are the following:

Journals

- Eloy Roura, Nicolae Sarbu, Arnau Oliver, **Sergi Valverde**, Sandra Gonzàlez, Ricard Cervera, Nuria Bargalló and Xavier Lladó. Automated detection of Lupus white matter lesions in MRI images. *Submitted to Frontiers in Human Neuroscience*. 2016. Frontiers. [JCR PS IF 3.626, Q1(13/76)].
- Eloy Roura, Arnau Oliver, Mariano Cabezas, **Sergi Valverde**, Deborah Pareto, Joan Carles Vilanova, Lluís Ramió-Torrentà, Àlex Rovira and Xavier Lladó. A toolbox for multiple sclerosis lesion segmentation. *Neuroradiology on October 2015, Vol. 57, Issue 10, pp. 1031-1043*. DOI: 10.1007/s00234-015-1552-2. Springer. [JCR RNMMI IF 2.485, Q2(41/125)].
- Mariano Cabezas, Arnau Oliver, **Sergi Valverde**, Brigitte Beltrán, Joan Carles Vilanova, Lluís Ramió-Torrentà, Àlex Rovira and Xavier Lladó. BOOST: a supervised approach for multiple sclerosis lesion segmentation. *Journal of Neuroscience Methods*. Vol. 237, pp 108-117, 2014. Elsevier. [JCR N IF 2.025, Q3(174/252)].
- Yago Díez, Arnau Oliver, Mariano Cabezas, **Sergi Valverde**, Robert Martí, Joan Carles Vilanova, Lluís Ramió-Torrentà, Àlex Rovira and Xavier Lladó. Intensity based methods for brain MRI longitudinal registration. A study on multiple sclerosis patients. *Neuroinformatics*. Vol 12(3), pp 365-379, 2014. Springer. [JCR CSTM IF 2.825, Q1(13/102)]

Conferences

- Eloy Roura, Arnau Oliver, Mariano Cabezas, **Sergi Valverde**, Deborah Pareto, Joan Carles Vilanova, Lluís Ramió-Torrentà, Àlex Rovira and Xavier Lladó. An SPM12 extension for multiple sclerosis lesion segmentation. *SPIE Medical Imaging 2016*. February 2016, San Diego, USA.
- **Sergi Valverde**, Arnau Oliver, Eloy Roura, Deborah Pareto, Joan Carles Vilanova, Lluís Ramió-Torrentà, Jaume Sastre-Garriga, Xavier Montalban, Àlex Rovira and Xavier Lladó. Evaluation of two automated lesion segmentation and filling pipelines for brain tissue segmentation of multiple sclerosis patients. *ECTRIMS 2015. Multiple Sclerosis*. October 2015, Barcelona, Spain. [JCR CN IF:4.472 Q1(25/191)].
- Eloy Roura, Arnau Oliver, Mariano Cabezas, **Sergi Valverde**, Deborah Pareto, Joan Carles Vilanova, Lluís Ramió-Torrentà, Àlex Rovira and Xavier Lladó.

A toolbox for segmenting multiple sclerosis lesions using T1w and FLAIR images. *ECTRIMS 2015. Multiple Sclerosis*. October 2015, Barcelona, Spain. [JCR CN IF:4.472 Q1(25/191)].

- **Sergi Valverde**, Arnau Oliver, Deborah Pareto, Joan Carles Vilanova, Àlex Rovira, Lluís Ramió-Torrentà and Xavier Lladó. SLF: a MS white matter lesion filling toolbox for the SPM software. *ECTRIMS 2014. Multiple Sclerosis*. September 2014, Boston, USA. [JCR CN IF:4.822 Q1(22/192)].
- Ester Quintana, Brigitte Beltrán, **Sergi Valverde**, René Robles-Cedeno, Hector Perkal, Xavier Lladó, José Manuel Fernández-Real and Lluís Ramió-Torrentà. Expression of miRNAs in multiple sclerosis cerebrospinal fluid and their relation to MR activity. *ECTRIMS 2014. Multiple Sclerosis*. September 2014, Boston, USA. [JCR CN IF:4.822 Q1(22/192)].
- Ester Quintana, Brigitte Beltrán, **Sergi Valverde**, René Robles-Cedeno, Hector Perkal, Xavier Lladó, José Manuel Fernández-Real and Lluís Ramió-Torrentà. Relación entre la expresión de mirnas en LCR y variables de RM. *Neurología*, vol 29, pp 66-67. 2014. [JCR CN IF:1.322 Q3(142/191)].
- **Sergi Valverde**, Arnau Oliver, Mariano Cabezas, Yago Díez, Jordi Freixenet, Xavier Lladó, Joan Carles Vilanova, Àlex Rovira and Lluís Ramió-Torrentà. A quantitative study of the effects of White Matter MS lesions on tissue segmentation methods. *ECTRIMS 2013. Multiple Sclerosis*. October 2013, Copenhagen, Denmark. [JCR CN IF:4.472 Q1(25/191)].

Acronyms

BET Brain Extraction Tool

BSE Brain Surface Extractor

CSF Cerebrospinal Fluid

EDSS Extended Disability Status Scale

GM Gray Matter

CNS Central Nervous System

MARGA Multispectral Adaptive Region Growing Algorithm

MRI Magnetic Resonance Image

MS Multiple Sclerosis

SPM Statistical Parametric Mapping

T1w T1-weighted

T2w T2-weighted

WM White Matter

MICCAI Medical Image Computing and Computer Assisted Interventions

CHB Childrens Hospital Boston

UNC University of North Carolina

Contents

Acknowledgements	i
Publications	iii
Acronyms	vii
1 Introduction	1
Introduction	1
1.1 Research context	1
1.1.1 Multiple Sclerosis	1
1.1.2 Magnetic resonance imaging in MS	2
1.1.3 Image analysis in MS	3
1.2 Research background	5
1.3 Objectives	7
1.3.1 Document structure	7

Chapter 1

Introduction

In this first chapter, we introduce the reader to the research context and background of the thesis. Afterwards, we describe the proposed objectives, and we situate this work within the research line of our group.

1.1 Research context

1.1.1 Multiple Sclerosis

The human nervous system can be divided into the central nervous system (CNS) consisting on the brain and the spinal chord, and the peripheral nervous system, which connects the CNS with the sense organs [7]. CNS is mainly constituted by two tissue components: gray matter (GM), which consists of neuronal cell bodies; and white matter tissue (WM), which is mainly composed of myelinated axon tracts [63]. In the case of the brain, it is mostly composed by GM and WM, both evolved by the Cerebro-spinal fluid (CSF), which provides basic mechanical and immunological protection to the brain inside the skull [63].

Multiple sclerosis (MS) is the most common chronic immune-mediated disabling neurological disease of the CNS [64], in which the insulating covers of the nerve cells in the spinal chord and brain are damaged [18]. Nowadays, MS is the most frequent non-traumatic neurological disease that causes more disability in young adults. It follows a similar behavior also seen in other putative autoimmune diseases, and affects twice as many women as it does in men [19]. It has a low incidence in childhood, but it increases rapidly in adulthood reaching a peak between 25 and 35 years, and then slowly declines, becoming rare at 50 and older [8]. So far, the world estimate for the disease is between 1.3 to 2.5 million cases, being relatively common

in Europe, the United States, Canada, New Zealand, and parts of Australia, but rare in Asia, and in the tropics and subtropics of all continents [8].

MS is characterized by areas of inflammation, demyelination, axonal loss, and gliosis scattered throughout the CNS, often causing motor, sensorial, vision, coordination, deambulation, and cognitive impairment [17]. Demyelination is the process of progressive damage to the protective covering (myelin sheath) around the axon of the neurons. Demyelinated axons conduct impulses at reduced or spontaneous velocity causing impairment in sensation, movement and cognition [18]. The different clinical courses of the disease are generally grouped in four subtype forms [43]. The *Relapsing/Remitting* (RRMS) form of the disease is characterized by exacerbation times where symptoms are present. These periods are followed by periods of remission, where the patient recovers partial or totally from the disease symptoms. The *Secondary Progressive* (SPMS) form is characterized by a gradual intensification of symptoms between affection relapses. The *Progressive remitting* (PRMS) form is typified by an increase in the relapsing times with significant recovery but with worsening symptoms in new relapsing intervals. Lastly, the *Primary Progressive* (PPMS) form is characterized by a severe decrease of remitting times with special localization in the brain. In general, 50% of RRMS patients after 10 years develop the SPMS form of the disease. After 25 years, the 90% of RRMS patients would develop the SPMS form.

1.1.2 Magnetic resonance imaging in MS

Magnetic Resonance Imaging (MRI) is a noninvasive medical imaging technique that is used in radiology to generate image representations of different internal anatomical organs and physiological processes of the body. Over the last 40 years, MRI have evolved as a clinical modality [33], and in particular as an essential tool for the diagnosis and evaluation of central nervous system disorders such as MS [24]. On MRI, MS plaques are well-delimited regions with hypo-intense signal intensity with respect to GM on T1-weighted (T1-w), while hyper-intense with respect to GM on T2-weighted (T2), Proton Density (PD) and Fluid Attenuated Inverse Recovery (FLAIR) modalities (see Figure 1.1).

In this aspect, new criteria for MS diagnosis and monitoring has been revised in the last years [48], due to the MRI sensitivity to reveal focal white matter (WM) lesion plaques and disease activity in time and space [26]. Additionally, various studies have also analyzed the correlation between MRI brain tissue atrophy measures and MS disability status, showing that tissue loss is an important component of the disease progression [14, 25, 27, 56]. Tissue loss seems to increase through the course of MS with a similar rate between 0.3% and 0.5% per year, and independently of the MS subtype [21, 56]. In general, GM atrophy is more associated with disability changes than WM atrophy [28], and not only in the RRMS and SPMS MS subtypes [27, 56], but also in CIS patients where several studies have shown a significantly

greater ventricular cavities and an associated GM loss on MRI scans of CIS patients that will develop MS compared to those who not [13, 25].

1.1.3 Image analysis in MS

Manual analysis of brain images is unfeasible in practice, given the large number of three-dimensional MRI slices for each patient and the possible intra/inter observer variability between experts [8]. This has led to the development since the early nineties of a wide number of lesion and tissue segmentation methods, with the aim to reduce the execution time and the inherent variability of manual annotations [16, 32, 39].

Pre-processing of MRI images

Acquired brain MRI volumes incorporate non brain tissue parts of the head such as eyes, fat, spinal cord or brain skull. Brain tissue extraction from non-brain tissue is commonly referred in the literature as skull-stripping. Skull-stripping has a direct effect on the performance of automated methods, as differences in skull stripping would lead into unexpected results in the tissue classification if skull or eyes are included as brain tissue [1, 49]. Among the different proposed methods for skull-stripping [1, 40, 54], the Brain Extraction Tool (BET) [62], and the Brain Surface Extractor (BSE) [59] are the most commonly used methods by the neuro-imaging community.

Furthermore, inherent characteristics of the MRI acquisition process such as differences in the magnetic field, bandwidth filtering of the data or eddy currents driven by field gradients usually derive in image artifacts that may also have a negative impact on the performance of methods [60]. In these cases, intensity correction of MRI images is either performed before lesion/tissue segmentation, or as an integrated part of the tissue segmentation pipeline. Among the former available strategies proposed [3, 37], the N3 [61], and N4 [68] methods are currently the de-facto standard tools used for intensity correction.

Automated lesion segmentation

MRI based diagnostic criteria for MS has led to an increasing need to analyze quantitatively focal MS lesions in individual and temporal studies [9, 48]. Different sequences such as T2-w, PD and FLAIR are often used in lesion classification, as MS lesions appear brighter than GM and WM on them. However, WM lesions often present a similar signal intensity profile to CSF on T2-w. In contrast, FLAIR

sequences suppress fluids from the image, restraining the CSF tissue effects on the acquired image, although some severe T2-w hyper-intense lesions appear similar to CSF in FLAIR [36].

A wide number of automated lesion segmentation techniques have been proposed during the last years [30, 42]. In these methods, classification is based either in supervised or unsupervised learning. Supervised learning methods employ a training set of correctly-identified observations that are used as prior information to learn the lesion characteristics. Newer proposed strategies integrate spatial decision forest [31], statistical methods [65], patch-based models [35] or adaptive dictionary learning strategies [22]. In contrast, unsupervised learning methods do not use any prior information in the classification task, which involves grouping data into categories based on some measure of inherent similarity or distance characteristic of the input images. Among these, most recent methods include probabilistic models which separate WM lesions from normal-appearing tissue by considering lesions as an outlier class [36, 38, 67], or techniques that make use of the signal intensity of lesions on FLAIR to apply several thresholding methods with post-processing steps to automatically segment lesions [53, 57].

Automated brain tissue segmentation in MS

The existent correlation between brain tissue atrophy measures and MS disability status [25, 27], has increased the necessity to develop robust automated brain tissue segmentation methods capable to perform accurate brain tissue volume measurements [34]. However, automated segmentation of brain tissue is still a challenging problem due to the complexity of the images, existence of lesions, differences in tissue intensities, noise, intensity inhomogeneities and the absence of models of the anatomy that fully capture the possible deformations in each structure [8, 39].

A wide number of brain tissue segmentation methods have been proposed so far. General purpose intensity based methods usually perform tissue segmentation on T1-w sequences, as this modality clearly separates gray matter from white matter. These include probabilistic strategies based on Bayesian inference [4, 46, 55, 59], Markov Random Fields models [6, 66, 72], or unsupervised clustering methods such as Zhang2001, *Fuzzy logic* [11, 47]. In contrast, supervised learning approaches also combines T1-w sequences with other modalities such as T2-w and PD using *K-Nearest-Neighbor* classifiers [20, 70], *Support Vector Machines* [2, 69], *Random Forests* [71, 45], or trained *Gaussian mixture models* [51].

However, different studies¹ have shown that tissue abnormalities found in MS image patients such as WM lesions reduce the accuracy of tissue segmentation methods

¹Em puc referenciar jo AJNR2015?

[5, 15]. Effectively, WM lesions on T1-w are hypo-intense with respect to normal-appearing WM, and therefore, lesion voxels that are classified as GM are distorting the overall GM volume. However, lesion voxels may also have an effect in the observed differences in normal-appearing tissue. WM lesions which are actually classified as WM decrease the mean overall signal intensity of the WM, causing that GM voxels with signal intensities similar to WM lesions may be also mis-classified as WM. In contrast, if WM lesions are classified as GM, normal-appearing WM voxels with signal intensities similar to lesions may be miss-classified as GM.

Lesion filling

In MS, hypo-intense WM lesions have to be pre-processed before tissue segmentation in order to reduce the effects of WM lesions on tissue segmentation. Historically, WM lesions have been masked-out of the T1-w before segmentation, and their volume have been added to WM afterwards [14]. Although this method effectively reduces the error in tissue volume, it has been show that this approach is not optimal [5, 15].

In this aspect, several strategies have proposed to in-paint lesions on the T1-w with signal intensities of the normal-appearing WM before tissue segmentation [5, 15, 44, 58], a process which is usually known in the literature as lesion filling. However, most of the available lesion filling methods require manual delineations of lesions, which may be tedious, challenging and time-consuming task depending of the characteristics of the image [42]. When available, lesion filling has demonstrated not only a significant reduction in the associated errors of WM lesions in tissue volume measurements [50], but also in image registration [12, 23, 58] and cortical thickness measurements [44].

1.2 Research background

This thesis is located within the framework of different research projects associated to the Computer Vision and Robotics research group (VICOROB), a research institute of the University of Girona². VICOROB has been working on several medical image analysis projects since 1996, mainly in segmentation and registration of mammography images. Lately in 2010, the research group started a fruitful collaboration with several medical MS research teams with the aim to develop new automated techniques capable to segment MS lesions and to perform atrophy measurements that can be transferred to experts for clinical use. In particular, our research in the MS field has been carried out within the following research projects:

²www.vicorob.udg.edu

1. [2009 – 2012] PI09/91918 “SALEM: Segmentación Automática de Lesiones de Esclerosis Múltiple en imágenes de resonancia magnética” awarded by the Instituto Carlos III.
2. [2010 – 2012] VALTEC09-1-0025 “Salem: toolkit para la segmentación automática de lesiones de Esclerosis Múltiple en resonancia magnética” awarded in 2009 by the Generalitat de Catalunya within the “Projectes de valorització VALTEC”.
3. [2015 – 2017] TIN2014-55710-R: “Herramientas de neuroimagen para mejorar el diagnosis y el seguimiento clínico de los pacientes con Esclerosis Múltiple” awarded in 2014 by the spanish call Retos de investigación 2014.
4. [2015 – 2019] BiomarkEM.cat: To develop, validate and transfer to clinical practice robust tools and totally automated for measuring new lesions and changes on the brain volume within MS patients. Awarded in 2015 by the Fundació Marató de TV3.

Since then, the research group has published original contributions in different fields such as image pre-processing [54], MS lesion segmentation [9, 10, 42, 53], temporal analysis [29, 41], image registration [23, 52], and tissue segmentation [8]. All this works have been published in partnership with different medical MS teams from:

- from the Hospital Vall d’Hebron: Dr. Rovira, who is the director of the “Unitat de Ressonància Magnètica-Centre Vall d’Hebron” (URMVH) and has participated in several research projects funded by public and private institutions in the last few years, as well as Dr. Pareto and technicians Huerga and Corral. This group is part of the MAGNIMS network, a European network of centres that share an interest in the MS study through MRI.
- from the Clínica Girona / Hospital Santa Caterina: Dr. Vilanova and Dr. Barceló are the codirectors of the “Unitat de Ressonància Magnètica” at the Clínica Girona and are members of several national and international radiology societies.
- from the Hospital Josep Trueta: Dr. Ramió-Torrentà, who is the current coordinator of the “Unitat de Neuroimmunologia i Esclerosi Múltiple”, as well as Drs. Robles and Beltrán, who work for the radiology unit.

1.3 Objectives

As part of the TIN2014-55710-R and BiomarkEM.cat research project frameworks, the main goal of this thesis is:

to develop a novel fully automated brain tissue segmentation method capable of computing accurate tissue volume measurements of MS patient images.

This objective refers to the brain tissue segmentation of MS patient images at a specific time but we do not consider the differences in tissue volume at different stages.

In order to fulfill this particular goal, different stages have to be covered. Each of them has been covered on different stages of the thesis.

This general goal can be actually divided into several sub-goals to focus at different time stages of this thesis.

1.3.1 Document structure

Bibliography

- [1] Julio Acosta-Cabronero, Guy B. Williams, João M S Pereira, George Pengas, and Peter J. Nestor. The impact of skull-stripping and radio-frequency bias correction on grey-matter segmentation for voxel-based morphometry. *NeuroImage*, 39(4):1654–1665, 2008.
- [2] Ayelet Akselrod-Ballin, Meirav Galun, Moshe John Gomori, Ronen Basri, and Achi Brandt. Atlas guided identification of brain structures by combining 3D segmentation and SVM classification. *Medical image computing and computer-assisted intervention : MICCAI ... International Conference on Medical Image Computing and Computer-Assisted Intervention*, 9(Pt 2):209–16, 2006.
- [3] J B Arnold, J S Liow, K a Schaper, J J Stern, J G Sled, D W Shattuck, a J Worth, M S Cohen, R M Leahy, J C Mazziotta, and D a Rottenberg. Qualitative and quantitative evaluation of six algorithms for correcting intensity nonuniformity effects. *NeuroImage*, 13(5):931–943, 2001.
- [4] John Ashburner and Karl J. Friston. Unified segmentation. *NeuroImage*, 26:839–851, 2005.
- [5] Marco Battaglini, Mark Jenkinson, Nicola De Stefano, and Nicola De Stefano. Evaluating and reducing the impact of white matter lesions on brain volume measurements. *Human Brain Mapping*, 33(9):2062–71, 2012.
- [6] S. Bricq, Ch. Collet, and J.P. Armspach. Unifying framework for multimodal brain mri segmentation based on hidden markov chains. *Medical Image Analysis*, 12(6):639 – 652, 2008. [Special issue on information processing in medical imaging 2007](#).
- [7] Per Brodal. *The central nervous system*. 2010.
- [8] Mariano Cabezas, Arnau Oliver, Xavier Lladó, Jordi Freixenet, and Meritxell Bach Cuadra. A review of atlas-based segmentation for magnetic resonance brain images. *Computer Methods and Programs in Biomedicine*, 104(3):e158–e177, 2011.

- [9] Mariano Cabezas, Arnau Oliver, Eloy Roura, Jordi Freixenet, Joan C. Vilanova, Lluís Rami-Torrent, Alex Rovira, and Xavier Lladó. Automatic multiple sclerosis lesion detection in brain MRI by FLAIR thresholding. *Computer Methods and Programs in Biomedicine*, 115(3):147–161, 2014.
- [10] Mariano Cabezas, Arnau Oliver, Sergi Valverde, Brigitte Beltran, Jordi Freixenet, Joan C. Vilanova, Lluís Rami-Torrent, Alex Rovira, and Xavier Lladó. BOOST: A supervised approach for multiple sclerosis lesion segmentation. *Journal of Neuroscience Methods*, 237:108–117, 2014.
- [11] Benoît Caldaïrou, Nicolas Passat, Piotr A Habas, Colin Studholme, and François Rousseau. A non-local fuzzy segmentation method: application to brain mri. *Pattern Recognition*, 44(9):1916–1927, 2011.
- [12] A Ceccarelli, J S Jackson, S Tauhid, A Arora, J Gorky, E Dell’Oglio, A Bakshi, T Chitnis, S J Khoury, H L Weiner, C R G Guttman, R Bakshi, and M Neema. The impact of lesion in-painting and registration methods on voxel-based morphometry in detecting regional cerebral gray matter atrophy in multiple sclerosis. *American Journal of Neuroradiology*, 33(8):1579–85, September 2012.
- [13] Antonia Ceccarelli, Maria a Rocca, Mohit Neema, Vittorio Martinelli, Ashish Arora, Shahamat Tauhid, Angelo Ghezzi, Giancarlo Comi, Rohit Bakshi, and Massimo Filippi. Deep gray matter T2 hypointensity is present in patients with clinically isolated syndromes suggestive of multiple sclerosis. *Multiple sclerosis (Houndmills, Basingstoke, England)*, 16(1):39–44, 2010.
- [14] D T Chard, G J M Parker, R Kapoor, A J Thompson, and D H Miller. Brain atrophy in clinically early relapsing–remitting multiple sclerosis. 2002.
- [15] Declan T. Chard, Jonathan S. Jackson, David H. Miller, and Claudia A M Wheeler-Kingshott. Reducing the impact of white matter lesions on automated measures of brain gray and white matter volumes. *Journal of Magnetic Resonance Imaging*, 32:223–228, 2010.
- [16] Harvey E. Cline, William E. Lorensen, Ron Kikinis, and Ferenc Jolesz. Three-Dimensional Segmentation of MR Images of the Head Using Probability and Connectivity. *Journal of Computer Assisted Tomography*, 14(6):1037–1045, 1990.
- [17] Alastair Compston and Alasdair Coles. Multiple sclerosis. *Lancet*, 359(9313):1221–31, 2002.
- [18] Alastair Compston and Alasdair Coles. Multiple sclerosis. *Lancet*, 372(9648):1502–17, 2008.

- [19] Devic M Confavreux C, Aimard G. Course and prognosis of multiple sclerosis assessed by the computerized data processing of 349 patients. *Brain*, 103(2):281–300, 1980.
- [20] Renske De Boer, Henri A Vrooman, Fedde Van Der Lijn, Meike W Vernooij, M Arfan Ikram, Aad Van Der Lugt, Monique M B Breteler, and Wiro Niessen. White matter lesion extension to automatic brain tissue segmentation on mri. *NeuroImage*, 45(4):1151–1161, 2009.
- [21] N. De Stefano, A. Giorgio, M. Battaglini, M. Rovaris, M. P. Sormani, F. Barkhof, T. Korteweg, C. Enzinger, F. Fazekas, M. Calabrese, D. Dinacci, G. Tedeschi, A. Gass, X. Montalban, A. Rovira, A. Thompson, G. Comi, D. H. Miller, and M. Filippi. Assessing brain atrophy rates in a large population of untreated multiple sclerosis subtypes. *Neurology*, 74(23):1868–1876, 2010.
- [22] Hrishikesh Deshpande, Pierre Maurel, and Christian Barillot. Classification of Multiple Sclerosis Lesions using Adaptive Dictionary Learning. *Computerized Medical Imaging and Graphics*, 2015.
- [23] Yago Diez, Arnau Oliver, Mariano Cabezas, Sergi Valverde, Robert Martí, Joan Carles Vilanova, Lluís Ramió-Torrentà, Àlex Rovira, and Xavier Lladó. Intensity based methods for brain MRI longitudinal registration. A study on multiple sclerosis patients. *Neuroinformatics*, 12(3):365–379, 2014.
- [24] Robert R Edelman and Steven Warach. Magnetic Resonance Imaging. *New England Journal of Medicine*, 328(10):708–716, 1993.
- [25] Massimo Filippi, Paolo Preziosa, Massimiliano Copetti, Gianna Riccitelli, Mark a Horsfield, Vittorio Martinelli, Giancarlo Comi, and Maria a Rocca. Gray matter damage predicts the accumulation of disability 13 years later in MS. *Neurology*, 81(20):1759–67, nov 2013.
- [26] Massimo Filippi and Maria A Rocca. MR imaging of multiple sclerosis. *Radiology*, 259(3):659–81, 2011.
- [27] Elizabeth Fisher, Jar-Chi Lee, Kunio Nakamura, and Richard A Rudick. Gray matter atrophy in multiple sclerosis: A longitudinal study. *Annals of Neurology*, 64(3):255–265, sep 2008.
- [28] Leonora K Fisniku, Declan T Chard, Jonathan S Jackson, Valerie M Anderson, Daniel R Altmann, Katherine A Miszkiel, Alan J Thompson, and David H Miller. Gray Matter Atrophy Is Related to Long- Term Disability in Multiple Sclerosis. *Ann Neurol*, 64:247–254, 2008.

- [29] Onur Ganiler, Arnau Oliver, Yago Diez, Jordi Freixenet, Joan C. Vilanova, Brigitte Beltran, Lluís Rami-Torrent, Alex Rovira, and Xavier Lladó. A subtraction pipeline for automatic detection of new appearing multiple sclerosis lesions in longitudinal studies. *Neuroradiology*, 56(5):363–374, 2014.
- [30] Daniel García-Lorenzo, Simon Francis, Sridar Narayanan, Douglas L Arnold, and D Louis Collins. Review of automatic segmentation methods of multiple sclerosis white matter lesions on conventional magnetic resonance imaging. *Medical Image Analysis*, 17(1):1–18, January 2013.
- [31] Ezequiel Geremia, Olivier Clatz, Bjoern H. Menze, Ender Konukoglu, Antonio Criminisi, and Nicholas Ayache. Spatial decision forests for MS lesion segmentation in multi-channel magnetic resonance images. *NeuroImage*, 57(2):378–390, 2011.
- [32] Guido Gerig, John Martin, Ron Kikinis, Olaf Kubler, Martha Shenton, and Ferenc A. Jolesz. Unsupervised tissue type segmentation of 3D dual-echo MR head data. *Image and Vision Computing*, 10(6):349–360, 1992.
- [33] Tal Geva. Magnetic resonance imaging: historical perspective. *Journal of Cardiovascular Magnetic Resonance*, 8(4):573–580, 2006.
- [34] Antonio Giorgio and Nicola De Stefano. Clinical use of brain volumetry. *Journal of Magnetic Resonance Imaging*, 37(1):1–14, 2013.
- [35] Nicolas Guizard, Pierrick Coupé, Vladimir S. Fonov, Jose V. Manjón, Douglas L. Arnold, and D. Louis Collins. Rotation-invariant multi-contrast non-local means for MS lesion segmentation. *NeuroImage: Clinical*, 8:376–389, 2015.
- [36] Rola Harmouche, Nagesh K Subbanna, D Louis Collins, Douglas L Arnold, and Tal Arbel. Based on Modeling Regional Intensity Variability and Local Neighborhood Information. 62(5):1281–1292, 2015.
- [37] Zujun Hou. A review on MR image intensity inhomogeneity correction, 2006.
- [38] Saurabh Jain, Diana M. Sima, Annemie Ribbens, Melissa Cambron, Anke Maertens, Wim Van Hecke, Johan De Mey, Frederik Barkhof, Martijn D. Steenwijk, Marita Daams, Frederik Maes, Sabine Van Huffel, Hugo Vrenken, and Dirk Smeets. Automatic segmentation and volumetry of multiple sclerosis brain lesions from MR images. *NeuroImage: Clinical*, 8:367–375, 2015.
- [39] T Kapur, W E Grimson, W M Wells, and R Kikinis. Segmentation of brain tissue from magnetic resonance images. *Medical image analysis*, 1(2):109–27, 1996.

- [40] Jong Min Lee, Uicheul Yoon, Sang Hee Nam, Jung Hyun Kim, In Young Kim, and Sun I. Kim. Evaluation of automated and semi-automated skull-stripping algorithms using similarity index and segmentation error. *Computers in Biology and Medicine*, 33(6):495–507, 2003.
- [41] Xavier Lladó, Onur Ganiler, Arnau Oliver, Robert Martí, Jordi Freixenet, Laia Valls, Joan C. Vilanova, Lluís Ramió-Torrentà, and Àlex Rovira. Automated detection of multiple sclerosis lesions in serial brain MRI. *Neuroradiology*, 54:787–807, 2012.
- [42] Xavier Lladó, Arnau Oliver, Mariano Cabezas, Jordi Freixenet, Joan C. Vilanova, Ana Quiles, Laia Valls, Lluís Ramió-Torrentà, and Àlex Rovira. Segmentation of multiple sclerosis lesions in brain MRI: A review of automated approaches. *Information Sciences*, 186(1):164–185, March 2012.
- [43] F. D. Lublin and S. C. Reingold. Defining the clinical course of multiple sclerosis: results of an international survey. National Multiple Sclerosis Society (USA) Advisory Committee on Clinical Trials of New Agents in Multiple Sclerosis. *Neurology*, 46(4):907–911, April 1996.
- [44] Stefano Magon, Laura Gaetano, M Mallar Chakravarty, Jason P Lerch, Yvonne Naegelin, Christoph Stippich, Ludwig Kappos, Ernst-Wilhelm Radue, and Till Sprenger. White matter lesion filling improves the accuracy of cortical thickness measurements in multiple sclerosis patients: a longitudinal study. *BMC Neuroscience.*, 15(1):106, January 2014.
- [45] D Mahapatra. Analyzing Training Information From Random Forests for Improved Image Segmentation. *Image Processing, IEEE Transactions on*, 23(4):1504–1512, 2014.
- [46] J.L. Marroquin, B.C. Vemuri, S. Botello, E. Calderon, and A. Fernandez-Bouzas. An accurate and efficient bayesian method for automatic segmentation of brain mri. *Medical Imaging, IEEE Transactions on*, 21(8):934 –945, aug. 2002.
- [47] Dzung L Pham. Spatial Models for Fuzzy Clustering. *Computer Vision and Image Understanding*, 297:285–297, 2001.
- [48] Chris H Polman, Stephen C Reingold, Brenda Banwell, Michel Clanet, Jeffrey a Cohen, Massimo Filippi, Kazuo Fujihara, Eva Havrdova, Michael Hutchinson, Ludwig Kappos, Fred D Lublin, Xavier Montalban, Paul O’Connor, Magnhild Sandberg-Wollheim, Alan J Thompson, Emmanuelle Waubant, Brian Weinshenker, and Jerry S Wolinsky. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Annals of neurology*, 69(2):292–302, 2011.

- [49] V. Popescu, M. Battaglini, W. S. Hoogstrate, S. C J Verfaillie, I. C. Sluimer, R. A. van Schijndel, B. W. van Dijk, K. S. Cover, D. L. Knol, M. Jenkinson, F. Barkhof, N. de Stefano, H. Vrenken, F. Barkhof, X. Montalban, F. Fazekas, M. Filippi, J. Frederiksen, L. Kappos, D. Miller, J. Palace, C. Polman, M. Rocca, A. Rovira, and T. Yousry. Optimizing parameter choice for FSL-Brain Extraction Tool (BET) on 3D T1 images in multiple sclerosis. *NeuroImage*, 61(4):1484–1494, 2012.
- [50] V. Popescu, N. C G Ran, F. Barkhof, D. T. Chard, C. A. Wheeler-Kingshott, and H. Vrenken. Accurate GM atrophy quantification in MS using lesion-filling with co-registered 2D lesion masks. *NeuroImage: Clinical*, 4:366–373, 2014.
- [51] Martin Rajchl, John S.H. Baxter, a. Jonathan McLeod, Jing Yuan, Wu Qiu, Terry M. Peters, and Ali R. Khan. Hierarchical max-flow segmentation framework for multi-atlas segmentation with Kohonen self-organizing map based Gaussian mixture modeling. *Medical Image Analysis*, 000:1–12, 2015.
- [52] E. Roura, T. Schneider, M. Modat, P. Daga, N. Muhlert, D. Chard, S. Ourselin, X. Lladó, and C. A. M. Wheeler-Kingshott. Multi-channel registration of fa and t1w images in the presence of atrophy: application to multiple sclerosis. *Functional Neurology*, 30(4), 2015.
- [53] Eloy Roura, Arnau Oliver, Mariano Cabezas, Sergi Valverde, Deborah Pareto, Joan C Vilanova, Lluís Ramió-Torrentà, Àlex Rovira, and Xavier Lladó. A toolbox for multiple sclerosis lesion segmentation. *Neuroradiology*, 57(10):1031–1043, 2015.
- [54] Eloy Roura, Arnau Oliver, Mariano Cabezas, Joan C. Vilanova, Àlex Rovira, Lluís Ramió-Torrentà, and Xavier Lladó. MARGA: Multispectral Adaptive Region Growing Algorithm for brain extraction on axial MRI. *Computer Methods and Programs in Biomedicine*, 113(2):655–673, 2014.
- [55] Snehashis Roy, Aaron Carass, Pierre-Louis Bazin, Susan Resnick, and Jerry L. Prince. Consistent segmentation using a rician classifier. *Medical Image Analysis*, 16(2):524 – 535, 2012.
- [56] Richard a Rudick, Jar-Chi Lee, Kunio Nakamura, and Elizabeth Fisher. Gray matter atrophy correlates with MS disability progression measured with MSFC but not EDSS. *Journal of the neurological sciences*, 282(1-2):106–11, jul 2009.
- [57] Paul Schmidt, Christian Gaser, Milan Arsic, Dorothea Buck, Annette Förchler, Achim Berthele, Muna Hoshi, Rüdiger Ilg, Volker J Schmid, Claus Zimmer, Bernhard Hemmer, and Mark Mühlau. An automated tool for detection of

- FLAIR-hyperintense white-matter lesions in Multiple Sclerosis. *NeuroImage*, 59(4):3774–3783, February 2012.
- [58] Michaël Sdika and Daniel Pelletier. Nonrigid registration of multiple sclerosis brain images using lesion inpainting for morphometry or lesion mapping. *Human Brain Mapping*, 30(4):1060–1067, 2009.
- [59] D W Shattuck, S R Sandor-Leahy, K a Schaper, D a Rottenberg, and R M Leahy. Magnetic resonance image tissue classification using a partial volume model. *NeuroImage*, 13(5):856–876, 2001.
- [60] A. Simmons, P. S. Tofts, G. J. Barker, and S. R. Arridge. Sources of intensity nonuniformity in spin echo images at 1.5 T. *Magnetic Resonance in Medicine*, 32(1):121–128, 1994.
- [61] J G Sled, A P Zijdenbos, and A C Evans. A nonparametric method for automatic correction of intensity nonuniformity in MRI data. *IEEE transactions on medical imaging*, 17(1):87–97, 1998.
- [62] Stephen M. Smith. Fast robust automated brain extraction. *Human Brain Mapping*, 17(3):143–155, 2002.
- [63] G H Sperber. Clinically Oriented Anatomy. *Journal of anatomy*, 208(3):393, 2006.
- [64] Lawrence Steinman, M.D. Multiple Sclerosis: A Coordinated Immunological Attack against Myelin in the Central Nervous System. *Cell*, 85(3):299–302, may 1996.
- [65] Elizabeth M Sweeney, Russell T Shinohara, Navid Shiee, Farrah J Mateen, Avni a Chudgar, Jennifer L Cuzzocreo, Peter a Calabresi, Dzung L Pham, Daniel S Reich, and Ciprian M Crainiceanu. OASIS is Automated Statistical Inference for Segmentation, with applications to multiple sclerosis lesion segmentation in MRI. *NeuroImage: Clinical*, 2:402–13, jan 2013.
- [66] Jussi Tohka, Ivo D. Dinov, David W. Shattuck, and Arthur W. Toga. Brain mri tissue classification based on local markov random fields. *Magnetic Resonance Imaging*, 28(4):557 – 573, 2010.
- [67] Xavier Tomas-Fernandez and Simon Keith Warfield. A Model of Population and Subject (MOPS) Intensities with Application to Multiple Sclerosis Lesion Segmentation. *IEEE transactions on medical imaging*, 0062(c):1–15, jan 2015.

-
- [68] Nicholas J. Tustison, Brian B. Avants, Philip A. Cook, Yuanjie Zheng, Alexander Egan, Paul A. Yushkevich, and James C. Gee. N4ITK: Improved N3 bias correction. *IEEE Transactions on Medical Imaging*, 29(6):1310–1320, 2010.
 - [69] Annegreet van Opbroek, Fedde van der Lijn, and Marleen de Bruijne. Automated brain-tissue segmentation by multi-feature svm classification. In *Proceedings of the MICCAI Workshops – The MICCAI Grand Challenge on MR Brain Image Segmentation (MRBrainS – 13)*, 2013.
 - [70] H Vrooman, Fedde van der Lijn, and W Niessen. Auto-knn: brain tissue segmentation using automatically trained knearest-neighbor classification. In *Proceedings of the MICCAI Workshops – The MICCAI Grand Challenge on MR Brain Image Segmentation (MRBrainS – 13)*, 2013.
 - [71] Zhao Yi, Antonio Criminisi, Jamie Shotton, and Andrew Blake. Discriminative, semantic segmentation of brain tissue in mr images. In Guang-Zhong Yang, David Hawkes, Daniel Rueckert, Alison Noble, and Chris Taylor, editors, *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2009*, volume 5762 of *Lecture Notes in Computer Science*, pages 558–565. Springer Berlin / Heidelberg, 2009.
 - [72] Y Zhang, M Brady, and S Smith. Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Transactions on Medical Imaging*, 20:45–57, 2001.