

Coherence in Chaos: Diffusion, Localization, and Decoherence in the Standard Map / Quantum Kicked Rotor Family

Author: Hadi Servat

Affiliation: Independent Researcher

Contact: publish@fractalresonance.com

Website: fractalresonance.com

© 2026 H. Servat, All Rights Reserved

Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0)

DOI: [10.5281/zenodo.21283579](https://doi.org/10.5281/zenodo.21283579) · **Version:** v2.3

Permanent record: <https://doi.org/10.5281/zenodo.21283579> · PDF: <https://zenodo.org/records/21283579>

Abstract

Version 2.3 of FRC 100.002 (v2.2 corrected: RP-match executed and retracted, §5.5) replaces the stadium-billiard fractal-dimension signature with a canonical chaos benchmark built on the Standard Map and Quantum Kicked Rotor, and states the basin claim in its corrected open-system form. The claim: chaotic dynamics are not structureless randomness; coherence structure inside chaos is measurable, and the transitions among classical diffusion, quantum dynamical localization, and decoherence-restored diffusion are organized by the entropy–coherence reciprocity ledger of the 566 line, read as an open-system balance with a measured — never posited — residual. The paper fixes

four anchors that exist independently of any simulation performed here: (1) the exact von Mises curve of FRC 566.030, against which a parameter-free classical structure functional is defined; (2) the analytic leading Ruelle–Pollicott resonance of area-preserving maps, as the forced target for the classical coherence decay rate; (3) the coherence-lifetime criterion, the operational form of FRC 787.787's flight condition, predicting an order-one delocalization crossover in $t^ \cdot \gamma_C$; (4) the cross-mechanism collapse target: diffusion reopening as a function of coherence degradation across distinct decoherence channels. Pilot simulations reported here support anchor (1), bracket anchor (3), and give a tensioned preliminary test of anchor (4); strict cross-mechanism collapse is not yet passed. Anchor (2) was executed (§5.5) and **retracted as a coherence-channel test**: the pilot decay rates are packet-spreading transients, the bare phase autocorrelation is analytically resonance-blind on the torus invariant measure, and the true leading transfer-operator resonance is a distinct, $\sim 5\times$ faster eigenmode (results-negative-kept). Two of this revision's own pre-registered instruments failed and are reported at equal volume. The stadium result is retained as a demoted geometric case study.*

1. Position in the basin series and lineage

The 100 series is the basin of the FRC corpus: each paper carries the current settled statement of one foundational claim, and is updated in place as the framework moves. This document is the updated statement of 100.002. It replaces the v1 stadium-billiard paper as the carrier of the chaos claim and retains the stadium result as a demoted case study in Appendix A with its corrected caveats.

Computational results — the pilot reported in §10 and the full simulation program of §12 — live in the answer appendix line FRC 100.002.001, so that the basin statement here remains stable while evidence accumulates.

Three corrections govern this version, stated up front:

1. **$D \approx 1.90$ is not a pillar.** The v1 fractal-dimension value is representative, not universal, and nothing below depends on it. The fractal content of the chaos claim is relocated to an object that is textbook-real and carries no fitted exponent: the KAM island hierarchy (§5.4).
2. **The ledger is stated in its open-system form, with the residual measured, not posited.** FRC 566.030 established that the system-only quantity $Q = S + k \cdot \ln C$ must not* be conserved across a transition in which a system gains or loses coherence; its change is the

reciprocity current, and a conserved system-only Q would refute the framework rather than confirm it. This paper defines its ledger current accordingly (§3) and defines nothing into an identity. In particular, no relation of the form $\lambda = -d_e S$ appears here: writing that identity would force system-only conservation and rebuild the exact closed-system error 566.030 was written to kill.

3. **The load-bearing arena is the Standard Map / Quantum Kicked Rotor family**, the most standardized system in classical and quantum chaos: one drive parameter, an exact quantum counterpart, dynamical localization, a mature atom-optics experimental record including controlled decoherence, and observables hostile readers already trust.

2. The claim

Chaos is not the absence of coherence; it is an arena in which coherence structure is measurable, and in which the entropy–coherence reciprocity of the 566 line organizes what standard treatments describe piecewise. Concretely: (i) in the classical mixed regime, residual coherence structure persists inside chaos and is quantified parameter-free against the exact maximum-entropy curve of 566.030; (ii) in the quantum regime, coherence built by the chaotic dynamics itself arrests the diffusion that same dynamics drives — dynamical localization read as an emergent coherence structure satisfying the operational flight criteria of 787.787 in momentum space; (iii) under decoherence, the target claim is that diffusion reopening is organized primarily by coherence-channel degradation, with strict mechanism-independent collapse still under test.

Scope of this version. This is a corrected preprint and pilot-supported statement of the 100.002 claim. The evidence is not yet final: Sigma must survive the full null suite, the $t^*\gamma_C$ order-one crossover needs parameter sweeps with a defined tolerance, and the cross-mechanism collapse must be checked out of sample against the experimental record. The gates (§8), controls (§9), and kill conditions (§11) define what would promote, revise, or remove the claim.

3. The ledger, in its correct frame

FRC 566.030 states the open-system balance:

$$dS_{\text{total}} + k_* \ln C = \sigma \geq 0,$$

with the system-only deficit $dQ = d(S_{\text{sys}} + k_* \ln C) = \sigma - dS_{\text{env}}$ as the exchange current. For a chaos experiment the environment is not modeled, so the operational object of this paper is the **system-only ledger current**

$$J_{\text{FRC}}(t) \equiv \frac{dS}{dt} + k_* \frac{d \ln C}{dt} \equiv \frac{dQ}{dt},$$

with S a coarse-grained entropy and C an operational coherence measure, both fixed before any outcome is examined (§6, §9). Three points of discipline:

- **The residual is measured.** J_{FRC} is whatever the system does; the predictions of this paper concern the *structure* of that residual (sign pattern, regime dependence, collapse across mechanisms), never its value by definition. Nothing here closes the ledger algebraically.
- **The residual's dynamical origin, where one is claimed, is the relaxation law of 787.787:** the coherence field lags its state-dependent target with finite relaxation time,
$$\tau_{\Lambda}(\partial_t \Lambda + u \cdot \nabla \Lambda) = D_{\Lambda} \nabla^2 \Lambda - \Lambda + \Lambda_{\mathrm{eq}}[\rho, \psi, E],$$
 and that lag — not a posited constant — is what generates nonzero current, hysteresis, and delayed collapse in driven systems.
- **k^* is a scale-covariant units bridge** (566.001, 566.030 §3): 1 nat in information units throughout this paper. A report of one fixed fitted k across domains would misread the 566 line; the invariant is the form and sign structure of the ledger, not a universal constant.

4. Symbol hygiene

Four symbols in this territory are shaped alike and must never be conflated:

Symbol	Meaning	Never confuse with
K_{drive}	classical/experimental kick strength; tunes the map	k^*
k^*	entropy–coherence units bridge of the 566 line (1 nat here)	K_{drive} , any fitted constant
λ / J_{FRC}	measured open-system ledger residual/ current	anything defined; d_e in particular
Λ	the coherence field of 787.787/100.007, $\Lambda = \Lambda_0 \ln C$, pinned by its relaxation law	the residual λ ; the quartic coupling λ_4 of 100.007

Additional working symbols: $\hbar_{\mathrm{eff}}$ (effective Planck scale of the quantum rotor), $C_{\mathrm{in}}(0,1]$ (operational coherence), S (coarse-grained entropy, nats), t^* (localization break time), γ_C (coherence decay rate), Σ (classical structure functional, §5.2).

5. Classical arena: the Standard Map and the structure functional

5.1 The map

$$p_{n+1} = p_n + K_{\mathrm{drive}} \sin x_n, \quad x_{n+1} = x_n + p_{n+1} \pmod{2\pi}.$$

As K_{drive} increases: regular motion, mixed phase space with KAM islands, resonance overlap near $K_c \approx 0.9716$, global chaos, momentum diffusion. The FRC reading replaces none of this; it adds an accounting layer with an exact, pre-existing null.

5.2 The structure functional (parameter-free, anchored on 566.030)

The rotor's position variable lives on a circle, so the corpus-standard coherence channel applies directly: $C = |\langle e^{ix} \rangle|$, the mean resultant length — the Kuramoto order parameter, the same object that is $C(\kappa) = I_1(\kappa)/I_0(\kappa)$ in 566.030 and the coherence readout of 840.101. For fixed C , the maximum-entropy distribution on the circle is von Mises, with entropy $S_{\mathrm{vM}}(C)$ known in closed form (566.030 §4). Define the **structure functional**

$$\Sigma(t) = S_{\mathrm{vM}}(C(t)) - S_{\mathrm{vM}}(C_{\mathrm{exact}}); \quad \Sigma(t) \geq 0$$

The inequality is a theorem (max-ent), not a finding — which is precisely its value: Σ is a properly signed, zero-free-parameter measure of *how much organization the phase marginal carries beyond what its coherence alone requires*. $\Sigma = 0$ means the ensemble sits on the exact Bessel curve published in 566.030 before this paper existed; $\Sigma > 0$ is coherence structure inside chaos, in nats. Equivalently $\Sigma = -\Delta Q$, the vertical distance below the exact $Q(C)$ curve, so the classical ledger of this paper is literally an overlay on the 566.030 figure.

5.3 Pre-registered readings

- **Globally chaotic regime** ($K_{\mathrm{drive}} \gtrsim 5$): $\Sigma(t)$ is transient — a filamentation spike as the initial structure is stretched, then decay to the max-ent curve as the marginal equilibrates.
- **Mixed regime** ($K_{\mathrm{drive}} \sim 0.5$ – 2.5): $\Sigma(\infty) > 0$ persists, consistent with KAM islands holding the stationary marginal off the max-ent curve. $\Sigma(\infty)$ should decrease monotonically with K_{drive} as the island hierarchy is destroyed; direct island-area tracking is appendix work.

5.4 The fractal, relocated honestly

The v1 intuition — that the fractal side of FRC lives in chaos — survives, but its carrier changes. The persistent Σ of the mixed regime is hypothesized to be anchored on the KAM island-around-island hierarchy: a self-similar object that is textbook-real, requires no box-counting fit, and

carries no $D \approx 1.90$ exposure. The pilot is consistent with that reading but does not directly measure island survival. The direct statement to be tested is: *coherence structure inside chaos sits on a fractal set*. That is the honest heir of the v1 claim, and the island-area comparison in FRC 100.002.001 decides how much of the claim survives. Criticality remains the deeper fusion point (fractal across scale, entanglement across space, joined at the transition), but that statement belongs to the 826 line and is only pointed to here.

5.5 The classical decay-rate anchor — executed, and retracted as a coherence-channel test (results-negative-kept, 2026-07-08)

This anchor was pre-registered as: the decay rate of the coherence channel $\langle e^{ix_n} e^{-ix_0} \rangle$ equals the leading Ruelle–Pollicott resonance of the transfer operator (Venegeroles 2008), with pilot empirical rates $\gamma \approx 0.152/\text{kick}$ at $K_{\text{drive}}=5$ and $\gamma \approx 0.418/\text{kick}$ at $K_{\text{drive}}=10$ as the targets. The test was run properly and **does not pass as stated; the pilot rates are not resonances**. Three checkable facts:

1. **The pilot rates are packet transients.** Reproduced (0.150/0.463) — but they measure the spreading of a concentrated non-equilibrium ensemble (the pilot starts $p \approx 0$ on a momentum line, x concentrated, and fits $n \in [1, 12]$), not the asymptotic correlation decay from the invariant measure.
1. **The bare phase autocorrelation is analytically resonance-blind on the torus.** The one-step correlation from the uniform invariant measure is exactly zero, $\langle e^{i(x_1 - x_0)} \rangle = \langle e^{ip_0} \rangle, \langle e^{iK \sin x_0} \rangle = 0 \cdot J_0(K) = 0$ (p_0 uniform, independent of x_0); numerically it drops to the finite- N floor in one step and is noise thereafter. The e^{ix} mode cannot carry the leading resonance.
1. **The true leading resonance is a different, faster object.** Computed from first principles as the leading eigenvalue of the Fourier–Bessel Koopman matrix — $U, e^{i(mx+np)} = \sum_l J_l((m+n)K) e^{i((m+l)x + (m+n)p)}$ — the leading Pollicott–Ruelle resonance is $|z_1| \approx 0.4$, $\gamma \approx 0.8$ – 0.95 at $K=5, 10$ (truncation-converged up to the residual dependence of a non-compact operator). It is ~ 5 times faster than the pilot transient and corresponds to a specific eigenmode combination, not the bare e^{ix} .

Disposition. The claim "the coherence-channel (phase-autocorrelation) decay rate is the leading resonance," and the specific values 0.152/0.418, are **retracted** — they were a finite-window artifact of a mis-specified observable, caught by executing the registered test. The transfer-operator leading resonance ($|z_1| \approx 0.4$) stands as a separate, first-principles characterization of the map. A corrected coherence-channel $\leftrightarrow$ resonance identity — projecting the coherence observable onto the leading Koopman eigenmode — is registered for the answer appendix (100.002.001); it is not claimed here. This retraction touches nothing else: the Σ structure functional (§5.2), the localization results, and the 566 ledger are independent of it. (Report and scripts: `output/probe/FRC-100-002-001-RP-2026-07-08/`.)

6. Quantum arena: the Kicked Rotor, localization as coherence structure

6.1 The system

$$H = \frac{p^2}{2I} + K_{\mathrm{drive}} \cos x \sum_n \delta(t - nT),$$

whose classical limit is the Standard Map and whose quantum evolution exhibits the decisive triplet: early classical-like diffusion, arrest by **dynamical localization** at break time t^* , and restoration of diffusion under decoherence. All three are experimentally realized in atom optics (Moore et al. 1995; Ammann et al. 1998; Klappauf et al. 1998, and the modern line through the Lille Anderson-transition experiments and interaction-driven breakdown, Cao et al. 2022).

6.2 Fixed channels and the anti-circularity control

The ledger requires two channels that are demonstrably not one function of each other. The Ising c-function gate (840.102 line) killed a prior FRC claim precisely because coherence and the flowing quantity closed on each other by definition; that failure mode is named here as a mandatory control.

- **Coherence channel $\mathcal{C}$** : ensemble purity $\mathrm{Tr}(\rho^2)$ of the noise-averaged density matrix (primary; operational, continuous with the existing pilot line), with nearest-off-diagonal momentum coherence $\sum_m |\rho_{m,m+1}|$ as the secondary interference channel.
- **Entropy channel $\mathcal{S}$** : Shannon entropy of the **coarse-grained diagonal** momentum distribution (bin width fixed in advance) — a population measure deliberately distinct from any functional of the full spectrum of ρ .
- **Identity check (named control)**: plot $\mathcal{S}$ against $-\ln \mathcal{C}$ across all runs. If the points collapse to a single curve, the ledger is algebra, not physics, and the quantum claim is void. Passing looks like what the pilot found (§10): at zero noise, $\mathcal{S}$ grows from 0 to ~ 1.6 nats while $\mathcal{C}$ stays pinned at 1 — a vertical branch no function can produce.

6.3 Localization read through 787.787

FRC 787.787 defines Lambda-Flight operationally: a structure is in flight if its internal coherence stays above threshold, its radius grows slower than the diffusive $\sqrt{2dDt}$, and its residual motion is predicted by the coherence gradient. The dynamically localized rotor satisfies the first two conditions verbatim, in momentum space: coherence holds near 1 while the momentum variance arrests against the classical diffusion it initially tracked. **Dynamical localization is an emergent coherence structure — order the chaotic dynamics itself builds through interference — and it is the first canonical, exactly simulable testbed for 787.787's flight criteria.** This paper therefore tests 787's predictions, not only 566's ledger.

6.4 The coherence-lifetime criterion (operational form of 787's flight/collapse condition)

787's collapse threshold says coherent transport fails when the coherence layer can no longer keep up with the structure. For the kicked rotor the operational translation is one dimensionless number:

$$\boxed{\text{decoherence crossover governed by } t^* \gamma_C = O(1).}$$

where t^* is the zero-noise break time and γ_C the early-window decay rate of the coherence channel under noise. Interference needs $\sim t^*$ kicks to build the localized state; if coherence dies faster than that, the structure never assembles and diffusion continues. The pilot brackets an order-one crossover: material reopening is already visible near $t^* \gamma_C \approx 0.5$, while the next resolved points sit near $2-3$. A sharp onset and tolerance band require the appendix sweep; this release does not claim them.

6.5 The cross-mechanism collapse

The ledger target is that diffusion reopens because the coherence channel is degraded. The strong version says the reopening magnitude is a **single function of coherence degradation, independent of the physical mechanism doing the degrading**: kick-phase noise, kick-amplitude noise, spontaneous emission, or interactions. Plotted against γ_C , all mechanisms would fall on one curve. This is the Gate-4 "many sweeps, one curve" target, not yet a passed result; the pilot reports both preliminary collapse and a low-noise mechanism split.

7. Junction: what 100.002 supplies to the corpus

Paper	Core claim	What 100.002 supplies
566 / 566.030	open-system entropy–coherence reciprocity; system-only Q moves, its motion is the current	a chaos-family ledger with an exact max-ent anchor; the structure functional as the overlay on 566.030's own curve
100.003	collapse is phase-locking into attractor basins	a concrete chaotic basin geometry (mixed phase space) where locking, basin structure, and decoherence are simulable
100.006	Born rule is the equilibrium of coherence dynamics	a mixing-plus-relaxation arena where equilibrium is approached, perturbed, and restored; 100.006's own coherence gauge is the von Mises family used here
100.009	local reciprocity dynamics; $\Phi_R = s + k_* \ln C$; activation from reciprocity defect	the global reading: $J_{\mathrm{FRC}} = d\Phi_R/dt$ with flux integrated out; one vocabulary across the local and global papers
787.787		

	Lambda-Flight: coherence transport, relaxation law, flight/collapse criteria, six failure conditions	the first canonical testbed: localization = flight in momentum space; the $t^*\gamma_C$ criterion; the hysteresis prediction (P4) testable by noise up/down sweeps
--	--	--

Boundary note (from 840.101, offered as framing only): classical chaotic mixing destroys information about the initial condition at the mixing rate — smearing in physical dynamics; the localized quantum state retains its initial momentum indefinitely — addressable structure. Smearing versus retention is one boundary in two substrates, and coherence is what buys retention. This sentence carries no evidential weight in this paper.

8. Gates for pillar status

Unchanged in structure from the working draft, tightened in content. Passing all four is what would promote this document from candidate to established pillar; none is yet claimed as passed:

1. **Canonical arena:** Standard Map / Quantum Kicked Rotor load-bearing; stadium demoted (Appendix A).
2. **Fixed observables before results:** Σ , C , J_{FRC} , Σ , γ_C , t^* all defined above, before outcomes; one primary coherence channel per regime, robustness channels secondary; no post-hoc switching.
3. **Cross-regime win:** the same ledger organizes classical diffusion (via Σ), localization (via the flight reading and $t^*\gamma_C$), and decoherence (via the cross-mechanism collapse).
4. **Better-than-baseline:** at least one of — $\Sigma(\infty)$ tracks directly measured KAM survival where entropy alone does not; the $t^*\gamma_C = O(1)$ crossover predicts the delocalization noise strength after a tolerance band is set; the reopening-vs- γ_C collapse holds across mechanisms including published experimental data; the classical decay rate matches the analytic Ruelle–Pollicott resonance. If J_{FRC} only redescribes diffusion or entropy growth, this paper is an application, not a pillar.

9. Controls and nulls

1. **Identity check** (§6.2) — mandatory; failure voids the quantum ledger. *[DONE in pilot as a zero-noise quantitative sanity check plus plotted channel scan: entropy span > 1 nat while purity remains pinned; full non-collapse statistic across all runs is appendix work.]*
2. **Shuffled/randomized-kick control:** deterministic structure destroyed while kick statistics preserved; Σ and the ledger structure must vanish. *[DONE in pilot as the matched stochastic null; passes. Note: temporal shuffles are the null for correlation/RP quantities, not for the one-time functional Σ .]*

3. **Matched stochastic diffusion null:** no deterministic structure; must not reproduce $\Sigma(\infty) > 0$ or the collapse. *[DONE in pilot; passes. Variance-matched variant registered for the appendix.]*
4. **Coarse-graining sweep:** entropy bins and windows varied; sign/threshold structure must be stable. *[DONE in pilot for Σ , $M = 36\text{--}144$; passes. Quantum bin sweep: appendix.]*
5. **$\hbar_{\text{eff}}$ sweep:** separate quantum interference from discretization artifact. *[Appendix.]*
6. **Finite-N / finite-R floors:** the classical overlay is read only above the $3/\sqrt{N}$ coherence floor; Σ bias calibrated on synthetic von Mises draws *[DONE]*; the quantum purity floor $1/R$ and the γ_C sampling bound are stated wherever they bind *[DONE for $R = 32$; larger R : appendix].*
7. **Pre-registration:** all thresholds (St^* definition, γ_C window, bin widths) fixed in the analysis scripts before sweeps are extended.

10. Pilot status — 2026-07-08

Reproducible scripts, seeds, JSON metrics, and figures accompany this preprint (`pilot_classical.py`, `pilot_quantum3.py`, `nulls_classical.py`). Pilot only; not publication-grade. Wins, kills, and open tensions are reported at equal volume.

Classical (Standard Map, $N = 4 \times 10^4$, 300 iterations, ensembles started on the max-ent curve):

- Exact-curve cross-check passes: the implementation reproduces the published 566.030 table ($Q = 0.9275$ at $C = 0.622$ vs published 0.928).
- **Estimator calibration (control a, done):** on exact synthetic von Mises samples, the Σ estimator bias is ≤ 0.0017 nats across $M = 36/72/144$ bins, matching the $(M-1)/2N$ histogram theory — an order of magnitude below the mixed-regime signals. Reported Σ values below are bias-corrected.
- Mixed regime: $\Sigma(\infty) > 0$ persists and decreases monotonically with drive — ~ 0.034 nats at $K = 0.6$, 0.011 at K_c , 0.0091 at $K = 1.4$, 0.0086 at $K = 2.0$, 0.0027 at $K = 2.5$ — consistent with the expected destruction of the island hierarchy. Direct KAM-survival tracking remains appendix work.
- **Matched stochastic diffusion null (control b, done):** with the kick's deterministic x -dependence destroyed (kick phase resampled uniformly per particle per step), the null loses all phase coherence ($C(\infty) \sim 0.004$, below the 0.015 finite-N floor) and carries no persistent Σ at any mixed-regime K where the true map holds $C(\infty) = 0.12\text{--}0.34$ with $\Sigma(\infty) > 0$. Caveat recorded: this null matches the kick statistics, not the diffusion coefficient (the true map's D is island-suppressed at these K); a variance-matched variant is registered for the appendix.
- **Coarse-graining sweep (control c, done):** $\Sigma(\infty)$ is stable across $M = 36/72/144$ (e.g., $K = K_c$: 0.0113 / 0.0114 / 0.0118 nats). Sign and ordering unchanged.

- Globally chaotic regime: Sigma is transient (filamentation peaks ~ 0.34 nats at $K = 2$, 0.10 at $K = 5$, 0.02 at $K = 10$, then decay to the curve within estimator noise).
- Empirical phase-correlation decay rates recorded ($0.152/\text{kick}$ at $K = 5$, $0.418/\text{kick}$ at $K = 10$).

The RP match was executed (§5.5, 2026-07-08) and these rates are retracted as resonances — they are packet-spreading transients; the bare phase autocorrelation is resonance-blind on the torus invariant measure, and the true leading transfer-operator resonance is $\gamma \approx 0.8 - 0.95$ ($\sim 5\times$ faster). Results-negative-kept.

Quantum (kicked rotor, $K_{\text{drive}} = 5$, $\hbar_{\text{eff}} = 1$, 512 momentum levels, $R = 32$ realizations, 200 kicks; uncertainties: split-half spread for γ_C , SEM over realizations for reopening):

- Zero noise: diffusion arrest at $t^* \sim 51$ kicks with purity pinned at 1 while coarse momentum entropy grows to ~ 1.6 nats — the identity check passes the stored zero-noise sanity criterion; localization meets 787's flight conditions 1-2 in momentum space.
- Coherence-lifetime criterion, with uncertainties: $t_{\gamma_C} = 0.48 \pm 0.01 \rightarrow \text{reopening } 1.88 \pm 0.06 \text{ (phase) and } 0.46 \pm 0.02 \rightarrow 1.66 \pm 0.04 \text{ (amplitude)}$; $t_{\gamma_C} = 2-3 \rightarrow \text{reopening } 4.5-4.9$; $t^*_{\gamma_C} \gg 1 \rightarrow 9-16x$. The pilot brackets an order-one crossover, but onset is unresolved between roughly 0.5 and 2-3; no tolerance band is claimed until the appendix sweep.
- Cross-mechanism collapse: preliminary and tensioned — consistent with a shared increasing curve in the resolved region, but with an **open tension reported at full volume**: at matched $\gamma_C \sim 0.009$ the reopenings differ by ~ 3 sigma (1.88 ± 0.06 phase vs 1.66 ± 0.04 amplitude). Either the collapse has a small mechanism-dependent residual at low noise, or the γ_C estimator (purity, early-window fit) is not yet the right abscissa there. The appendix must resolve this; the strict one-curve claim is NOT yet passed.
- Resolution limit recorded: γ_C saturates at the channel-sampling bound ($1/\text{EVERY} = 0.25/\text{kick}$) for $\sigma \geq 0.1$ (phase) / 0.2 (amplitude); the high-noise end of the collapse plot is bounded, not measured. Finer sampling registered for the appendix.

Kills recorded in this revision (this paper's own instruments, falsified by its own runs):

- **Kill A:** the l1-normalized off-diagonal coherence gauge is non-monotonic under localization-length growth — a badly designed instrument, dropped. Purity retained as primary.
- **Kill B:** the naive differential prediction "white amplitude noise does not reopen diffusion" is false — at matched sigma, per-kick amplitude jitter and phase jitter reopen diffusion almost identically. The experimental amplitude-noise robustness in the atom-optics record (quantum resonances, colored-noise regimes) does not transfer to white per-kick jitter, and this paper does not claim it. What survives — and is stronger — is the mechanism-collapse claim of §6.5, now itself under tension at low noise (above) and awaiting the out-of-sample test against the published record (Ammann 1998; Klappauf 1998; Steck dissertation ch. 4; White—

Ruddell–Hoogerland 2014; Cao 2022), mechanism by mechanism, with the coherence measure each experiment actually reports.




Figure 1. Classical Standard Map trajectories against the exact von Mises ledger curve, with structure-functional and empirical RP diagnostics.




Figure 2. Quantum kicked-rotor channels: diffusion reopening, purity decay, cross-noise collapse, lifetime criterion, and identity check.

10a. Data and code availability

All quoted values are produced by the accompanying scripts with fixed seeds; no value in this paper is hand-entered from an unscripted run. The release package contains the manuscript, HTML print source, PDF, rendering script, analysis scripts, generated figures, JSON metrics, a checksum manifest, and a separate reproduction report. Analysis scripts create a scratch `results/` directory when rerun. JSON metrics are compared with `pilot/compare_metrics.py`, which uses relative and absolute tolerances so near-zero floating-point values do not create false failures.

File	Role
FRC-100-002-v2.2.md	this paper
FRC-100-002-v2.2.html	HTML print source for the PDF
FRC-100-002-v2.2.pdf	rendered PDF preprint
render_frc_html_pdf.py	local HTML-to-PDF renderer
pilot/pilot_classical.py	Standard Map overlay, structure functional, empirical RP rates
pilot/pilot_quantum3.py	kicked rotor, channels, criterion, collapse, uncertainties (R = 32)
CHECKSUMS.txt	SHA-256 manifest — transfer integrity of the shipped package
pilot/compare_metrics.py	tolerance-aware reproduction check for regenerated JSON (default rtol 1e-6, atol 1e-12)
pilot/nulls_classical.py	estimator calibration, matched stochastic null, coarse-grain sweep
pilot/classical_metrics.json, pilot/quantum_metrics.json, pilot/classical_nulls.json	machine-readable outputs
fig1_classical_vonmises_overlay.png, fig2_quantum_channels.png	figures
REPRODUCTION-REPORT.md	release-regeneration commands, environment, and comparator results

11. Kill conditions

This chaos application fails if any of the following holds:

- the identity check fails (\$\$\$ and \$\ln C\$ collapse to one function under the fixed definitions);
- \$\Sigma(\infty)\$ in the mixed regime does not track KAM survival, or vanishes under reasonable coarse-graining, or is reproduced by the shuffled/matched-noise controls;
- the delocalization crossover is not controlled by order-one \$t^*\gamma_C\$ after a fine sweep, or the reopening-vs-\$\gamma_C\$ curve fails to collapse across mechanisms;
- ~the classical coherence decay rate does not match the analytic leading Ruelle–Pollicott resonance~~ (this kill item was mis-specified — the bare phase-autocorrelation observable is

analytically resonance-blind on the torus, §5.5; the corrected item is that a coherence observable *projected onto the leading Koopman eigenmode* fails to match $|z_1|$, registered for 100.002.001);

- everything above is fully explained by standard observables with no stable residual structure.

Failure here kills this chaos application. It does not kill the 566 reciprocity line, whose own kill conditions live in the 566 papers.

12. Answer appendix program (FRC 100.002.001)

1. Full $K_{\{\mathrm{drive}\}}$ sweep of $\Sigma(\infty)$ against direct KAM-measure estimates (island area fraction), with coarse-graining and N sweeps.
2. Ruelle–Pollicott match: numerical leading resonance of the transfer operator vs the analytic values vs the coherence-channel decay rate.
3. Quantum sweeps in $K_{\{\mathrm{drive}\}}$, $\hbar_{\mathrm{eff}}$, and noise; criterion tolerance band for $t^*\gamma_C$; mechanism set extended to simulated spontaneous emission and, from the literature, interactions.
4. Out-of-sample test of the cross-mechanism collapse against the published atom-optics record.
5. Hysteresis test of 787 Prediction 4: noise swept up then down at rate r ; lag vs τ_{Λ_r} .
6. Matched stochastic and shuffled-phase nulls throughout.
7. Negative results retained in the public record as results-negative-kept.

Appendix A — The stadium billiard, demoted

The v1 stadium result is retained as a geometric case study: a multiscale perturbation of the stadium billiard altering nodal geometry, with quantum scars as the established phenomenon of residual structure in chaotic eigenfunctions. Its correct role is illustrative; its incorrect roles — central proof, universal $D \approx 1.90$ signature, sole test of "chaos has structure" — are withdrawn. Any future use requires matched random-potential controls, parameter sweeps, and resolution-sensitivity analysis. The v1 record remains visible in the concept lineage with corrected caveats.

References

1. B. V. Chirikov, "A universal instability of many-dimensional oscillator systems," *Physics Reports* 52, 263–379 (1979).
2. G. Casati, B. V. Chirikov, F. M. Izrailev, J. Ford, "Stochastic behavior of a quantum pendulum under a periodic perturbation," *Lecture Notes in Physics* 93, 334–352 (1979).
3. S. Fishman, D. R. Grempel, R. E. Prange, "Chaos, Quantum Recurrences, and Anderson Localization," *PRL* 49, 509 (1982).

4. F. L. Moore, J. C. Robinson, C. Bharucha, B. Sundaram, M. G. Raizen, "Atom Optics Realization of the Quantum Delta-Kicked Rotor," *PRL* 75, 4598 (1995). DOI: 10.1103/PhysRevLett.75.4598.
5. H. Ammann, R. Gray, I. Shvarchuck, N. Christensen, "Quantum Delta-Kicked Rotor: Experimental Observation of Decoherence," *PRL* 80, 4111 (1998). DOI: 10.1103/PhysRevLett.80.4111.
6. B. G. Klappauf, W. H. Oskay, D. A. Steck, M. G. Raizen, "Observation of Noise and Dissipation Effects on Dynamical Localization," *PRL* 81, 1203 (1998). DOI: 10.1103/PhysRevLett.81.1203.
7. D. A. Steck, dissertation, ch. 4: Localization and Decoherence in the Kicked Rotor.
8. R. Venegeroles, "Leading Pollicott–Ruelle resonances for chaotic area-preserving maps," *Phys. Rev. E* 77, 027201 (2008). DOI: 10.1103/PhysRevE.77.027201.
9. D. H. White, S. K. Ruddell, M. D. Hoogerland, "Phase noise in the delta kicked rotor: from quantum to classical," *New J. Phys.* 16, 113039 (2014). DOI: 10.1088/1367-2630/16/11/113039.
10. A. Cao et al., "Interaction-driven breakdown of dynamical localization in a kicked quantum gas," *Nature Physics* 18, 1302 (2022). DOI: 10.1038/s41567-022-01724-7.
11. L. A. Bunimovich, "On the ergodic properties of nowhere dispersing billiards," *Commun. Math. Phys.* 65, 295 (1979).
12. E. J. Heller, "Bound-State Eigenfunctions of Classically Chaotic Hamiltonian Systems: Scars of Periodic Orbits," *PRL* 53, 1515 (1984).
13. H. Servat, *FRC 566.001 — Entropy–Coherence Reciprocity and UCC*. DOI: 10.5281/zenodo.17437759.
14. H. Servat, *FRC 566.030 — Reciprocity in Action: The Entropy–Coherence Current Across a Synchronization Transition*.
15. H. Servat, *FRC 100.003 — Resonant Collapse*. DOI: 10.5281/zenodo.15079820.
16. H. Servat, *FRC 100.005 — Thermodynamic Consistency*. DOI: 10.5281/zenodo.17438231.
17. H. Servat, *FRC 100.006 — Born Rule from Resonant Equilibrium*. DOI: 10.5281/zenodo.17438360.
18. H. Servat, *FRC 100.009 — Local Reciprocity Dynamics*. DOI: 10.5281/zenodo.20328893.
19. H. Servat, *FRC 787.787 — Lambda-Flight: Coherence Transport in Driven Open Physical Systems*.
20. H. Servat, *FRC 840.101 — The Phase–Attention Boundary*. DOI: 10.5281/zenodo.20416012.