

Strong Coupling Constant from the Photon Structure Function

Simon Albino and Michael Klasen*

*II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, D-22761 Hamburg, Germany*

Stefan Söldner-Rembold†

FNAL, P.O. Box 500, MS 357, Batavia, IL 60510, USA

(Dated: April 12, 2018)

We extract the value of the strong coupling constant α_s from a single-parameter pointlike fit to the photon structure function F_2^γ at large x and Q^2 and from a first five-parameter full (pointlike and hadronic) fit to the complete F_2^γ data set taken at PETRA, TRISTAN, and LEP. In next-to-leading order and the $\overline{\text{MS}}$ renormalization and factorization schemes, we obtain $\alpha_s(m_Z) = 0.1183 \pm 0.0050(\text{exp.})^{+0.0029}_{-0.0028}(\text{theor.})$ [pointlike] and $\alpha_s(m_Z) = 0.1198 \pm 0.0028(\text{exp.})^{+0.0034}_{-0.0046}(\text{theor.})$ [pointlike and hadronic]. We demonstrate that the data taken at LEP have reduced the experimental error by about a factor of two, so that a competitive determination of α_s from F_2^γ is now possible.

The theory of strong interactions, Quantum Chromodynamics (QCD), is one of the corner stones of the Standard Model of elementary particle physics. The precise determination of its fundamental parameter, the strong coupling constant α_s , bears important implications for the validity not only of QCD itself, but also of even more fundamental theories, since these have to contain the Standard Model as an effective field theory in the low-energy limit. A more fundamental theory, which might explain the size of the strong coupling constant, has yet to be established. Therefore, α_s must currently be extracted from experiment. Among the large variety of processes that have been used to this end, the most precise values have been obtained in Z -boson- and τ -decays at LEP, scaling violations in structure functions at HERA, and quarkonium decay branching fractions and lattice calculations of quarkonium mass splittings, leading – together with other, less precise measurements – to a current world average of $\alpha_s(m_Z) = 0.1172 \pm 0.0020$ at the mass of the Z -boson, $m_Z = 91.1876$ GeV [1].

When the photon structure function F_2^γ was first discussed in the context of QCD, a precise determination of α_s quickly emerged as one of its most interesting applications. Due to the pointlike coupling of the photon to quarks, the leading order (LO, $\mathcal{O}(\alpha/\alpha_s)$) [2] and next-to-leading order (NLO, $\mathcal{O}(\alpha)$) [3] contributions to $F_2^\gamma(x, Q^2)$ are calculable in QCD perturbation theory, if the virtuality Q in the deep-inelastic electron-photon scattering process is significantly larger than the asymptotic scale parameter Λ . Unfortunately, this pointlike contribution exhibits a power singularity at small Bjorken- x [4], which becomes rapidly stronger in higher orders [5]. The singularity can be regularized with a non-perturbative [6] or transverse-momentum [7] cut-off, but then the sensitivity to α_s is reduced and a dependence on the unphysical cut-off is introduced [8]. It is then necessary to fit both α_s and the cut-off to experimental data. Alternatively, the singularity can be canceled order by order in perturbation theory by retaining a hadronic boundary

condition at a low starting scale Q_0 [9]. In this case it is necessary to fit α_s and the hadronic input to experimental data. However, the evolution of the hadronic input to the physical scale Q is still predicted by perturbative QCD through inhomogeneous evolution equations [10], and the negligibility of the hadronic input can be tested *a posteriori*. Both methods have been applied in the past to PEP and PETRA data yielding $\Lambda_{\overline{\text{MS}}}^{(4)} = 180^{+100}_{-90}$ MeV [11] or $\alpha_s(m_Z) = 0.108^{+0.008}_{-0.010}$. This value contributed to the world average in the 1988 [12], 1990 [13], and 1992 [14] issues of the Review of Particle Properties, but was then abandoned on the grounds that there were “no new results and the data do not contribute significantly to the average” [15]. Since then it has been widely believed [16–20] that the sensitivity of F_2^γ to α_s is small.

In this Letter, we point out that over the last decade a wealth of new F_2^γ data has been collected at the e^+e^- -colliders TRISTAN and LEP, which extends to high average values of Q^2 , $\langle Q^2 \rangle \leq 780$ GeV². We demonstrate that the new data improve the sensitivity of F_2^γ to α_s significantly and that a single-parameter pointlike fit as well as a five-parameter full (pointlike and hadronic) fit to PETRA, TRISTAN, and LEP data yield results, which are not only consistent with the world average, but also have competitive experimental and theoretical errors.

We work in a fixed flavor number scheme with three active quark flavors (u, d, s). It is well known [21] that for current measurements of F_2^γ the available hadronic energy squared $W^2 = Q^2(1-x)/x$ is not much larger than the production threshold $4m_h^2$ of the heavy quarks ($h = c, b, t$), so that mass effects can not be neglected and the massive, fixed order $\mathcal{O}(\alpha)$ expression for the Bethe-Heitler process $\gamma^*(Q^2)\gamma \rightarrow h\bar{h}$ [22] should be used instead of the massless, factorized $\mathcal{O}(\alpha/\alpha_s)$ expression. For a consistent NLO ($\mathcal{O}(\alpha)$) analysis, we do not include the known [23], but numerically small, $\mathcal{O}(\alpha\alpha_s)$ corrections to the Bethe-Heitler process and omit the $\mathcal{O}(\alpha\alpha_s^2)$ contributions from the process $\gamma^*(Q^2)g \rightarrow h\bar{h}$. The heavy quark

masses are not well constrained from measurements of F_2^γ . We adopt a charm quark mass of $m_c = 1.5 \pm 0.1$ GeV in good agreement with recent precise determinations from threshold production at e^+e^- -colliders [24]. At large Q^2 , charm quarks contribute up to 40 % to F_2^γ in the whole x -range, while at small Q^2 they contribute at most 10% below $x \sim 0.2$. The contribution from bottom quarks is suppressed by a relative factor 1/16 from the different quark charges, while the threshold for top quark production lies at extremely small $x = 10^{-5} \dots 10^{-3}$, so that these contributions are both numerically negligible.

Since we wish to omit spurious higher order terms, which arise from the convolution of NLO contributions to the parton densities with the NLO Wilson coefficients and lead to instabilities at large x [21], we work in Mellin moment space, where the convolutions reduce to simple products, the evolution can be done analytically and without any approximations, and spurious higher order terms can be consistently omitted. The resulting prediction for F_2^γ is then converted back to x -space using a fast inverse Mellin transform with logarithmic mapping and fitted to experimental measurements with the multi-dimensional minimization algorithm MINUIT [25]. The quality of the fit is measured in terms of the χ^2 value per degree of freedom, χ^2/DF , for all selected data points.

We include in our analysis all published measurements of F_2^γ collected at the high-energy e^+e^- -colliders PETRA [26–28], TRISTAN [29–31], and LEP [32–38]. If more than one set of statistically overlapping data exists, the most recent publication is used. We exclude from our fit the data published by the TPC/2 γ Collaboration at PEP [39, 40], since several data points, mainly at low x , are inconsistent with measurements published by PLUTO [27], L3 [34], and OPAL [37] in the range $1.9 < Q^2 < 5.1$ GeV². Data where the charm component has been subtracted are also discarded. Statistical uncertainties and correlations between data points due to the experimental unfolding are taken into account as provided by the experiments, while systematic uncertainties are assumed to be uncorrelated. Due to this assumption the values of χ^2/DF are expected to be on average slightly less than unity. If asymmetric errors are given by the experiments, the data points are taken at the center of the full error interval. Most experiments have not corrected for the finite virtuality of the target photon P^2 . We neglect P^2 in this analysis, since usually $P^2 \ll Q^2$.

For our pointlike fit, we identify the starting scale Q_0 with the asymptotic scale parameter Λ , so that the hadronic input vanishes automatically and only a single parameter (Λ , or equivalently $\alpha_s(m_Z)$) has to be fitted. As discussed above, this is only justified at large x and Q^2 , where the residue of the pointlike singularity is expected to be small. Therefore, we perform our single-parameter pointlike fit only to a subset of data points with $x \geq 0.45$ and $Q^2 \geq 59$ GeV². Very similar results are obtained with the widely used values of $Q_0 = 0.5 \dots 0.6$

TABLE I: χ^2/DF and $\alpha_s(m_Z)$ values obtained in LO and NLO in the $\overline{\text{MS}}$ and DIS_γ factorization schemes with a single-parameter fit of the pointlike photon structure function F_2^γ . Also shown are the results obtained without LEP data and with very high Q^2 data.

Scheme	χ^2/DF	$\alpha_s(m_Z)$
LO	7.9/ 19	$0.1260 \pm 0.0055(\text{ex})^{+0.0061}_{-0.0055}(\text{th})$
$\overline{\text{MS}}$	9.1/ 19	$0.1183 \pm 0.0050(\text{ex})^{+0.0029}_{-0.0028}(\text{th})$
DIS_γ	8.1/ 19	$0.1195 \pm 0.0051(\text{ex})^{+0.0031}_{-0.0028}(\text{th})$
w/o LEP	3.2/ 7	$0.1244 \pm 0.0126(\text{ex})^{+0.0033}_{-0.0032}(\text{th})$
high Q^2	11.9/ 8	$0.1159 \pm 0.0125(\text{ex})^{+0.0018}_{-0.0018}(\text{th})$

GeV [16–18, 20], while choosing $Q_0 = 1$ GeV significantly increases the value of χ^2/DF ; two-parameter pointlike fits of α_s and Q_0 are driven to $Q_0 \simeq \Lambda$. In the first three lines of Tab. I we list the χ^2/DF and $\alpha_s(m_Z)$ values obtained in LO and NLO. The NLO fit is performed in two factorization schemes ($\overline{\text{MS}}$ and DIS_γ [21]) with different treatment of the pointlike Wilson coefficient in F_2^γ , but the numerical variation is found to be small. The total values of χ^2/DF as well as those for the individual data sets (not shown) lie around unity or below, indicating that the pointlike photon structure function and the fitted values of $\alpha_s(m_Z)$ describe the data sets well within their statistical and systematic uncertainties. The experimental errors are determined by varying $\alpha_s(m_Z)$ until the total value of χ^2 is increased by one unit. To estimate the theoretical error, we vary the charm quark mass as indicated above and follow the common convention of varying the factorization and renormalization scales by factors of two about their central value, the physical scale Q . We then add these three individual errors in quadrature. The LO value of $\alpha_s(m_Z)$ is consistent with the NLO value within the expected accuracy, $\mathcal{O}(\alpha_s^2)$, and the theoretical error is reduced from LO to NLO as expected. In the fourth line of Tab. I, we list the result of a fit without the LEP data. The experimental error is more than doubled, showing that the LEP data have considerably increased the sensitivity of F_2^γ to α_s at high x and Q^2 . When data at all values of x , but very high Q^2 ($Q^2 \geq 284$ GeV²) are fitted, the central value of $\alpha_s(m_Z)$ remains virtually unchanged (last line of Tab. I). At very high Q^2 , the theoretical error drops by a factor of two, whereas the experimental error increases. Measurements of F_2^γ at a future linear e^+e^- - or $e\gamma$ -collider like TESLA at very high values of Q^2 and with small experimental errors will therefore lead to even more precise determinations of α_s .

The goodness of our pointlike fit may also be judged from Fig. 1, where the fitted data points are shown as full circles, while those that have been omitted from the fit are shown as open circles, and where the statistical and systematic errors have been added in quadrature. The theoretical curves are perturbatively stable, *i.e.* LO and




FIG. 1: Single-parameter fits of the pointlike photon structure function, compared to PETRA [26], TRISTAN [29, 31], and LEP [32, 34–36, 38] data at large Q^2 . The data points marked by open circles have not been used in the fits. Also shown is the hadronic contribution from a five-parameter NLO fit of the full photon structure function in the DIS_γ scheme.

NLO fits differ only by small amounts. The choice of factorization scheme affects the region of very large x , but it has only a minor effect on the description of the data. Also shown in Fig. 1 is the hadronic contribution from a five-parameter NLO fit of the full photon structure function in the DIS_γ scheme. It clearly falls from small to large x and Q^2 and amounts to only a few percent in the region that has been used in the pointlike fit.

For our full (pointlike and hadronic) fit, we start from the observations that F_2^γ is dominated by the u -quark density in the photon and is only sensitive to the combined density of d - and s -quarks, whose contribution is furthermore suppressed by the smaller d - and s -quark charges. The gluon contributes to F_2^γ in LO only through a rather weak coupling to the quark singlet density in the evolution equations. A consecutive fit of the u -quark, d - and s -quark, and gluon densities shows, that only the first is well constrained by F_2^γ data and that the fit does not improve, when more degrees of freedom are added. Therefore we do not impose a hadronic boundary condition for the gluon and assume, that the hadronic fluctuations of the photon are insensitive to the quark charge, *i.e.* we identify the hadronic boundary conditions for u -quarks and d - and s -quarks at the starting scale Q_0 . To-

TABLE II: Q_0 , χ^2/DF , and $\alpha_s(m_Z)$ values obtained in LO and NLO in the $\overline{\text{MS}}$ and DIS_γ factorization schemes with a five-parameter fit of the hadronic photon structure function F_2^γ . Also shown are the results obtained without LEP data.

Scheme	Q_0/GeV	χ^2/DF	$\alpha_s(m_Z)$
LO	0.79 ± 0.18	121/129	$0.1475 \pm 0.0074(\text{ex})^{+0.0141}_{-0.0072}(\text{th})$
$\overline{\text{MS}}$	0.83 ± 0.09	118/129	$0.1198 \pm 0.0028(\text{ex})^{+0.0034}_{-0.0046}(\text{th})$
DIS_γ	0.85 ± 0.09	115/129	$0.1216 \pm 0.0028(\text{ex})^{+0.0033}_{-0.0050}(\text{th})$
w/o LEP	0.46 ± 0.10	37/ 38	$0.1147 \pm 0.0047(\text{ex})^{+0.0282}_{-0.0033}(\text{th})$

gether with $\alpha_s(m_Z)$ and Q_0 , we then fit the parameters N , α , and β of our ansatz $f_{u,d+s}^\gamma(x, Q_0^2) = Nx^\alpha(1-x)^\beta$ to the full data set described above. In the first three lines of Tab. II we list the Q_0 , χ^2/DF , and $\alpha_s(m_Z)$ values obtained with this five-parameter fit in LO and NLO. The starting scale Q_0 is perturbatively stable and is found to be close to the masses of the light vector mesons ρ , ω , and ϕ in contrast to earlier claims that the perturbative evolution of F_2^γ sets in only at rather high values of $Q_0 \sim 2 \text{ GeV}$ [41]. The individual and total values of χ^2/DF lie again around unity or below, so that the fitted full photon structure functions describe the full data set well within the experimental uncertainties. Note that the χ^2 value for the four TPC/2 γ points at $Q^2 = 2.8 \text{ GeV}^2$, which have not been used in the fits, is 18.0 and thus very large. The gluon density, generated with $f_g^\gamma(x, Q_0^2) = 0$, turns out to be in good agreement with recent H1 dijet data [42]. The experimental errors on the values of Q_0 and $\alpha_s(m_Z)$ reflect an increase in χ^2 by one unit, when all other fit parameters are kept fixed. Due to the larger number of data points in the full fit, the experimental error turns out much smaller than in the pointlike fit. When the full fit is performed without the LEP data (last line of Tab. II), the experimental error is almost doubled, *i.e.* the impact of the LEP data is again impressive. A fit to LEP data only leads to almost identical results as the full fit. The theoretical error in LO and without the LEP data gets a large asymmetric contribution from doubling the factorization scale, which is highly correlated with an increase in the fitted value of Q_0 and which is drastically reduced in the full NLO fit. Similar results as those listed in Tab. II are obtained, when only u -quarks are assigned a hadronic boundary condition.

In Fig. 2 we compare our results to the fitted F_2^γ data in the region of low x and Q^2 . This region is clearly dominated by the hadronic contribution and by the impact of the LEP data. A fit without the LEP data results in a rise of F_2^γ at low x , which is much too steep. The fits are perturbatively stable and the data are described almost equally well in the $\overline{\text{MS}}$ and DIS_γ scheme.

Since the total error on $\alpha_s(m_Z)$ is smaller in the full fit than in the pointlike fit due to the larger number of data points, we adopt as our final result




FIG. 2: Five-parameter fits of the full photon structure function, compared to data from PETRA [27], TRISTAN [30, 31], and LEP [32–35, 37] at small Q^2 . The data points marked by open circles refer to the second experiment and/or Q^2 value. Also shown are the hadronic and pointlike contributions to the NLO fit in the DIS_γ scheme.

$$\alpha_s(m_Z) = 0.1198 \pm 0.0054 \quad (1)$$

in NLO and the $\overline{\text{MS}}$ scheme, where the larger theoretical error has been added to the experimental error in quadrature. While our total error is slightly larger than those obtained in Z -boson- and τ -decays at LEP, it is comparable to the errors obtained in deep-inelastic scattering at HERA and heavy quarkonium decays. This encourages us to combine our result with the current world average of 0.1172 ± 0.0014 [1] to a new world average

$$\alpha_s(m_Z) = 0.1175 \pm 0.0014, \quad (2)$$

where the errors are assumed to be uncorrelated.

In conclusion, we have for the first time fitted the now final PETRA, TRISTAN, and LEP data on the photon structure function F_2^γ in NLO of perturbative QCD. We have extracted the value of the strong coupling constant $\alpha_s(m_Z)$ with competitive experimental and theoretical errors from a single-parameter pointlike fit to data at large x and Q^2 and from a five-parameter full (pointlike and hadronic) fit at all x and Q^2 . Our analysis proves that the available F_2^γ data contribute significantly to a precise determination of α_s and that future measurements of F_2^γ at linear colliders will have a large impact.

We thank G. Kramer for many valuable discussions and a careful reading of the manuscript. S. A. and M. K. are supported by the Deutsche Forschungsgemeinschaft through Grant No. KL 1266/1-2.

* michael.klasen@desy.de

† Heisenberg Fellow

- [1] K. Hagiwara *et al.*, Phys. Rev. D **66** (2002) 010001.
- [2] E. Witten, Nucl. Phys. B **120** (1977) 189.
- [3] W. Bardeen and A. Buras, Phys. Rev. D **20** (1979) 166; **21** (1979) 2041(E).
- [4] D. Duke and J. Owens, Phys. Rev. D **22** (1980) 2280.
- [5] G. Rossi, Phys. Lett. B **130** (1983) 105.
- [6] I. Antoniadis, G. Grunberg, Nucl. Phys. B **213** (1983) 445.
- [7] J. Field, F. Kapusta, L. Poggioli, Phys. Lett. B **181** (1986) 362.
- [8] W. Frazer, Phys. Lett. B **194** (1987) 287.
- [9] M. Glück and E. Reya, Phys. Rev. D **28** (1983) 2749; G. Rossi, *ibid.* **29** (1984) 852.
- [10] R. DeWitt, L. Jones, J. Sullivan, D. Willen and H. Wyld, Phys. Rev. D **19** (1979) 2046; **20** (1979) 1751(E).
- [11] W. Wagner, UCD-86-29, XXIII ICHEP, Berkeley (1986).
- [12] G. Yost *et al.*, Phys. Lett. B **204** (1988) 1.
- [13] J. Hernandez *et al.*, Phys. Lett. B **239** (1990) 1; **253** (1990) 524(E).
- [14] K. Hikasa *et al.*, Phys. Rev. D **45** (1992) S1; **46** (1992) 5210(E).
- [15] L. Montanet *et al.*, Phys. Rev. D **50** (1994) 1173.
- [16] M. Glück, E. Reya, A. Vogt, Phys. Rev. D **46** (1992) 1973.
- [17] P. Aurenche, M. Fontannaz and J. P. Guillet, Z. Phys. C **64** (1994) 621.
- [18] G. Schuler and T. Sjöstrand, Z. Phys. C **68** (1995) 607.
- [19] L. Gordon and J. Storrow, Nucl. Phys. B **489** (1997) 405.
- [20] M. Glück, E. Reya and I. Schienbein, Phys. Rev. D **60** (1999) 054019; **62** (1999) 019902(E).
- [21] M. Glück, E. Reya, A. Vogt, Phys. Rev. D **45** (1992) 3986.
- [22] V. Budnev, I. Ginzburg, G. Meledin and V. Serbo, Phys. Rept. **15** (1974) 181.
- [23] E. Laenen, S. Riemersma, J. Smith and W. van Neerven, Phys. Rev. D **49** (1994) 5753.
- [24] J. Kühn, M. Steinhauser, Nucl. Phys. B **619** (2001) 588.
- [25] F. James, M. Roos, Comp. Phys. Comm. **10** (1975) 343.
- [26] W. Bartel *et al.*, Z. Phys. C **24** (1984) 231.
- [27] C. Berger *et al.*, Phys. Lett. B **142** (1984) 111; Nucl. Phys. B **281** (1987) 365.
- [28] M. Althoff *et al.*, Z. Phys. C **31** (1986) 527.
- [29] S. Sahu *et al.*, Phys. Lett. B **346** (1995) 208.
- [30] T. Kojima *et al.*, Phys. Lett. B **400** (1997) 395.
- [31] K. Muramatsu *et al.*, Phys. Lett. B **332** (1994) 477.
- [32] R. Barate *et al.*, Phys. Lett. B **458** (1999) 152.
- [33] P. Abreu *et al.*, Z. Phys. C **69** (1996) 223.
- [34] M. Acciarri *et al.*, Phys. Lett. B **436** (1998) 403; *ibid.*, **447** (1999) 147; *ibid.*, **483** (2000) 373.
- [35] K. Ackerstaff *et al.*, Phys. Lett. B **411** (1997) 387.
- [36] K. Ackerstaff *et al.*, Z. Phys. C **74** (1997) 33.
- [37] G. Abbiendi *et al.*, Eur. Phys. J. C **18** (2000) 15.
- [38] G. Abbiendi *et al.*, arXiv:hep-ex/0202035.
- [39] H. Aihara *et al.*, Phys. Rev. Lett. **58** (1987) 97.
- [40] H. Aihara *et al.*, Z. Phys. C **34** (1987) 1.
- [41] L. Gordon, D. Holling, J. Storrow, J. Phys. G **20** (1994) 549.
- [42] C. Adloff *et al.*, Phys. Lett. B **483** (2000) 36.