
The Thiele Machine

No Free Insight



DEVON THIELE

DEVON THIELE

The Thiele Machine

No Free Insight

The Thiele Machine: No Free Insight

Devon Thiele

Copyright 2025–2026 Devon Thiele. Licensed under Creative Commons Attribution-ShareAlike 4.0 (CC BY-SA 4.0).

The source of this book, its proofs and its build are public: github.com/sethirus/The-Thiele-Machine. Archived at doi.org/10.5281/zenodo.17316437.

Set in Charter. Built from its source with pdfTeX.

The cover and part illustrations were generated with AI in the engraving tradition of Gustave Doré. They are new illustrations, not reproductions of his work.

For Julie, and for my dad.

Julie,

Three years ago, a few months into the rendering engine, I came out of my office and told you I thought I was onto something. You said okay. You didn't ask what it was, or what it was for, or when it would be done, and you didn't ask me for anything I could have shown you, because there was nothing to show. There was a program that had done something it shouldn't have been able to do, and a man who didn't yet know what a Turing machine was.

You gave me the three years anyway. The nights after the lot closed. The weekends. The hours I was in the house and not in the house, because my head was in a wall I couldn't get past. Every dead end, and there were more of those than there are pages here. You never once asked me to stop, and you never once said it was fine when it wasn't. You just kept giving me the next evening.

This book is what I did with them. It's the one thing I've built from nothing that I'm sure of, because I threw out every piece of it I couldn't prove, and what's left holds. I know that isn't what you gave the years for. You gave them for me. But it's what I have to show, and I wanted it to be worth showing.

Dad,

The name on the cover was yours first. I asked you before I used it and you said go ahead. You and Mom taught me that nothing is true just because somebody says it is. This book is that lesson, hundreds of pages long. I put our name on it because I wanted one thing in the world with that name on it that nobody handed me, that I couldn't have faked, and that I can stand behind in front of anyone. Here it is.

Julie carries the name now too. So the machine has it third. This is for the two of you who had it first.

Contents

What this is 9

1 Any machine that steps 12
- 1.1 Is the pencil thing a computer? 12
- 1.2 How far does that go? 13
- 1.3 Try to break it 14
- 1.4 Now ask it about keeping 15
- 1.5 The machine that stops 15
- 1.6 What Turing actually proved 16

2 Running out of room 18
- 2.1 Four people, three chairs 18
- 2.2 This is very old 19
- 2.3 The hotel that never runs out 21
- 2.4 Cut the strip 22
- 2.5 How long can a finite machine hold off? 23
- 2.6 What's being counted 24

3 Why charge for it 25
- 3.1 A rule you can refuse 25
- 3.2 The first door: endless paper 26
- 3.3 The second door: let it come back down 26
- 3.4 The third door: don't charge for it 27
- 3.5 How much 28
- 3.6 Found, assumed, chosen 29

4 The smallest one 31
- 4.1 Clay, seals, and seven witnesses 31
- 4.2 Two columns and a stamp 32
- 4.3 Closer to now 33
- 4.4 Five jobs 34
- 4.5 Two boxes and a coin 35
- 4.6 What the tally is and isn't 36
- 4.7 Any machine can wear one 37
- 4.8 Which question gets charged? 37
- 4.9 The cheapest price list 38

5 More settled 39
- 5.1 A row of cups 39

5.2 Naming what's on the table 41
5.3 The toll on the whole direction 42
5.4 One cup isn't enough 43
5.5 Somebody got here first, from the side 44
5.6 Why crossing a line has to cost 44

6 Looking isn't keeping 46
6.1 A small being at a small door 46
6.2 Bennett moves where the cost is 47
6.3 The demon on a sheet of paper 48
6.4 Two rows of dots 49
6.5 Who holds the pen 50
6.6 Learning after the first look 51
6.7 Finishing together 51
6.8 The certainty count 52
6.9 Who gets charged is partly bookkeeping 52
6.10 Somebody got there first, goddamn it 53
6.11 What the toll is for, said at last 54

What this is

I think in shapes, so I'll start with one.

Take a thing. Anything. A coin on a table, a light switch, a page in a notebook. Right now it's one way. It could be another.

Now do something to it. Flip the coin, hit the switch, write on the page. That's a move: a direction that takes the thing from how it is to how it is next.

Here's a computer you can run with a pencil. Write 3 on a scrap of paper. The move is: erase the number and write the next one up. Do it twice. 3, 4, 5. You just added 2 and 3. Nothing happened there but a thing, a move, and the next thing. Hold on to that picture; whether it covers every computer is something I'll have to earn, and I will, later, with you watching.

Here's the part I couldn't let go of. Some moves just shuffle. Flip the coin and you can flip it back, and nothing is different for good. But some moves make a difference that stays. Write a word in pen. Blow a fuse. Check an answer and mark it checked. After that move something is the case that wasn't before, and it stays the case.

I figured there had to be a cost in that step. Not in the shuffling. In the step where something goes from not settled to settled and stays there. That was a hunch, so test it on the pencil computer.

Tell it when to stop: count up, and when you reach 5, stop. Ask one question of each number: is it done? 3, no. 4, no. 5, yes. And once it says yes it keeps saying yes, because the machine has stopped.

A machine that has stopped is still somewhere on the next tick. It stays where it is: 5 goes to 5. Draw the arrows. 4 goes to 5. 5 goes to 5. Two different starting places, one landing place. Standing on 5, you can't tell whether you just got there or have been there all along.

Could you dodge that? Try letting the stopped machine rock back and forth instead of sitting still: 5-tick, 5-tock, 5-tick, both of them saying done. Draw the arrows. 4 goes to 5-tick. 5-tock goes to 5-tick. Two arrows land on 5-tick, the spot where you came in. Rocking only moved the place where they meet. The one way out is to never land anywhere twice: make every tick write something new, a tally mark, the time, anything, so the paper never repeats. That works as long as there's paper. On a finite sheet you run out, and two arrows land on one spot again.

So whenever this machine settles its answer and keeps it, something gives. Either two states become one, or it uses up fresh paper on every tick for as long as it runs.

That part isn't a hunch. You drew it. You drew it for one small counter; the book proves it for any machine that goes from one state to a next, and shows you where.

What I do with it is a choice, and I'll say so: I charge for the move where the answer goes from not done to done. Charging for it is up to me. That something has to give once it makes that move isn't. The two arrows don't always meet on that very move. Give done two spots, 5a and 5b: 4 goes to 5a, 5a goes to 5b, and 5b stays put. Now the arrows from 5a and 5b meet on 5b, a tick after the answer settled. But after the move from not done to done, there's no avoiding it: two arrows meet somewhere, or the paper never repeats.

Two more things have to be pinned down before the rule means anything, and you can see both with what's already on the table.

First, take the coin again. Flip it, flip it back, forever, free. Ask whether it's settled. It never is. A rule that says "a move that settles something has to cost" never asks the coin for anything, because the coin never settles anything. So the rule alone would let a coin that does nothing count. The machine has to actually settle something, at least once.

Second, suppose a later move could pay you back. The move that settles charges you one, the next move hands you one back, and the whole run comes out free. So prices can be nothing or something, never a refund.

One last thing before the whole definition. Settled doesn't have to be just yes or no. Take a checklist with three boxes. None ticked, one, two, all three: each tick settles it a little more, and ticks don't come off. "Done or not" is the smallest case, a checklist with one box. So when I say "more settled" from here on, picture the checklist.

So here's the Thiele Machine, all of it. A thing. Moves that take it to the next thing. A way to tell how settled the thing is, where settled never goes back to unsettled. Some run, from a start you pick, that actually settles it more than it was. A price on every move, nothing or more, never a refund. And one rule: a move that settles something more than it was costs something.

That's the whole definition. The rest of the book works out what follows from it, one step at a time, and tries to break it.

Now look back at the pencil computer with a stop in it. That's Turing's machine, in miniature. Turing's machine, from his 1936 paper, is a tape and a table of rules: from where it is, the table says where it goes next. A thing and moves. It already has something that settles and stays settled: whether it has stopped, the same question you asked of 3, 4 and 5. What Turing never put in is a price on the move that stops it. Put one in, and a Turing machine that ever stops fits every line above. So this isn't a rival to Turing's machine, and it computes nothing his can't. His machine is one kind of Thiele machine, with the price left out.

Now try this. Give two people the same messy list of numbers. The first sorts it, then goes back down it checking that every number is no bigger than the next. The second copies a sorted list off somebody else's sheet. Now look at the two sheets. Same numbers, same order. Look as hard as you like: nothing on either sheet tells you which one was checked. The difference was real, and neither sheet kept it. It's the same thing you saw standing on 5: you can see where you are, and not how you got there.

So put a box on each sheet that says checked. A plain Thiele machine charges for ticking the box, but nothing in the rules above stops the copier from paying and ticking it anyway. Paying for the tick doesn't make it true. So add one more condition: the box can only be ticked right after a check of that very list passes. Now the copier's tick is refused, and a ticked box tells you a check of that list passed. I call a box like that established. It's a condition you put on the machine. The bare definition doesn't hand it to you.

It's called the Thiele Machine. The afterword says why.

CHAPTER 1

Any machine that steps

When I started all this I didn't know what a Turing machine was. I knew it had something to do with computers, and that's it. I had a program that had done something I didn't expect, and a hunch, and no words for either. Most of the words in this book I found the same way: I'd hit a wall, go looking for whoever had been there before me, and not move on until I could watch the idea actually work. This chapter is the first of those trips. I'll take you on it the way I took it.

The question I had was the one I left you with on page one. I'd been calling that pencil-and-scrap-of-paper thing a computer. Is it? Or did I just make it small enough to look like one?

1.1 Is the pencil thing a computer?

Start with what the word used to mean. Before about 1940 a computer wasn't a machine. It was a job. In the late 1800s the Harvard Observatory hired rooms full of women to do the arithmetic of the sky, cataloguing stars by hand for a fraction of what a man was paid. Some of them found things nobody else had; Henrietta Leavitt worked out a way to tell how far away a star is. (I got that from Dava Sobel's history of those women and from the historians at the American Institute of Physics.) Rooms like that worked out artillery tables in both world wars; that one's from David Alan Grier's history of human computers.

So the pencil thing wasn't a toy version of a computer. For a long time it was the only kind there was. Good start, but that's history, not proof. A person with a pencil can do a lot of things that aren't following rules. I wanted to know what was left if you took the person away and kept only the rules.

That's exactly the question Alan Turing asked, in a paper in 1936. Going in, all I knew was that it had something to do with computers. What's in there is a clerk. He wrote down, as plainly as he could, everything a person sitting at a desk with a pencil does when they follow a procedure, and threw out everything else. What was left had four parts, and you can build it on the table in front of you right now.

Draw a strip of three squares. That's the tape. Write 1 in the first square, 1 in the second, leave the third blank.

Put a coin on the first square. That's where the clerk is looking. Turing called it the head.

On a scrap of paper write the letter A. That's which step of the procedure the clerk is on. Turing called it the state.

Now the rules. Write these two on a card:

> If you're on A and looking at a 1: leave it, move the coin one square right, stay on A.
>
> If you're on A and looking at a blank: write a 1 there, don't move, and change the letter to STOP.

That's the whole machine: a tape, a head, a state, and a card of rules. Now run it. Each tick, read the card and do what it says. After each tick, write down everything about the table: what's on each square, where the coin is, and the letter. I'll put brackets around the square the coin is on.

[1]	1	␣	A
1	**[1]**	␣	A
1	1	**[␣]**	A
1	1	**[1]**	STOP

It started with two 1s and ended with three. It added one. That's the same thing the pencil computer did when it went from 4 to 5.

Now look at what you wrote down on each line. The whole tape, where the coin is, and the letter. Is there anything about this machine at that moment that isn't on the line? Try to think of something. The rules don't change. The tape, the coin and the letter are all there is. So one line is the whole thing, and the card takes one line to the next line. Thing, move, next thing.

So the pencil computer wasn't a cheat. Turing's machine is the pencil computer with a bigger piece of paper.

1.2 How far does that go?

It would be nice if the answer were "every computer, done." It isn't quite. Push on it and it came apart into three different claims that people tend to run together. Here they are, and what each one stands on.

The first one is the one we just did. Every machine you can write down as a whole state and a rule for the next state has the shape from page one. You checked that yourself for Turing's machine: one line, one card, next line.

The second is that the other ways people have written computation down, with registers instead of a tape, or with functions handed to functions (that's Alonzo Church's lambda calculus, from the same year as Turing's paper), compute exactly the same things Turing's machine does. I didn't prove that, and I won't pretend I did. Other people proved it, one pairing at a time; Turing himself sketched the

match with Church's version in an appendix to his 1936 paper. What you can check yourself is that each of those is also a whole state and a rule for the next one, the same way you just did with the coin.

The third is the big one, and here's where I had to be careful. Does this cover everything a person could ever do by following rules? Does it cover every machine anybody could ever build out of real stuff? Those turned out to be claims nobody can prove, and for a good reason. "Everything a person could do by following rules" isn't a precise enough thing to prove anything about. That claim has a name, the Church-Turing thesis. "Every machine you could build" is a claim about physics, and the only way to knock it down would be to build one that beats Turing's, which nobody has done. Neither is a theorem, and in this book neither will get treated like one.

So the debt from page one comes out like this. For every machine you can write down as a whole state and a rule for the next state, paid: you just checked it. For every machine anybody could ever build, that's a claim about the world, it's labeled as a claim, and I'm not going to lean on it.

1.3 Try to break it

Now the question that actually mattered to me. Page one showed that when the counter settled on 5, two arrows had to meet, unless it used fresh paper forever. That was one little counter. Is it true of every machine with this shape?

Don't take my word for it. Get a sheet of paper and do this.

Draw some dots. Ten, twenty, as many as you like. Those are states. From every dot, draw exactly one arrow to some dot. It can point anywhere, even back at the dot it starts from. That's the one rule: every state has exactly one next state.

Now pick a dot to start on and follow the arrows. Keep a list of every dot you've been on.

Do it a few times, with different drawings and different starts. Watch what happens at each step. You'll find that each step does one of three things.

It lands on a dot that isn't on your list yet. A new one.

It lands on the dot you started from. Now you're going around in a circle, and since every dot has only one arrow out, you'll go around that same circle forever.

Or it lands on some other dot that's already on your list. Look at the first time that happens. The first time you got to that dot, you came in on an arrow from the dot right before it on your list. Now you've come in again, on an arrow from the dot you were just on. Those have to be two different dots, because until this step you hadn't been on any dot twice. Two arrows, from two different dots, landing on one. That's a merge, the same thing as 4 and 5 both going to 5. And from there you're on a path you've walked before, so you walk it again, forever.

Now try to make it do a fourth thing. Really try. Draw whatever you want.

Here's why you can't. Wherever the next arrow points, that dot is either on your list or it isn't. If it isn't, it's new. If it is, it's either the start or it isn't the start. There's no other place for the arrow to point. Three cases, and they cover everything.

And notice what you never had to know: how many dots there were. Ten dots or a billion or an endless sheet, it's the same three cases. What you did need was that every dot has exactly one arrow out. Hold on to that, because it's about to matter.

1.4 Now ask it about keeping

Go back to your dots. Color the dot you start on white. Pick some other dots and color them black, and make one rule about the coloring: no arrow is allowed to go from a black dot to a white one. Once you're on black, you stay on black. That's "settled stays settled." White is not done, black is done.

Follow a run from your white start until it lands on black. Now which of the three things can happen after that?

It can't circle back to the start. The start is white, and you can never get back to white once you're on black.

So it's one of the other two. Either it keeps landing on new dots forever, or two arrows meet somewhere.

Try to draw a counterexample: a run that goes from white to black, stays black, never lands on a new dot forever, and never has two arrows meet. You'll be drawing for a while. There isn't one, and the three cases above are the reason.

That's what page one promised: "the book proves it for any machine that goes from one state to a next." You drew it for one counter there. Now you've drawn it for any machine at all, any number of dots.

What it doesn't tell you is anything about heat. Whether two arrows meeting gives off heat in a real device is a question about the device. People argue about it, and they get their own chapter. This is dots and arrows and nothing else.

1.5 The machine that stops

Here's where the "exactly one arrow out" part bites.

Go back to the tape and the coin. The last line was 1 1 [1], STOP. What does the card say to do next? Nothing. There's no rule for STOP. The machine just stops. In dots and arrows: the STOP dot has no arrow out.

That breaks the one thing the three cases needed. So try the keeping question on a machine that stops. Draw two dots. "Working" is white. "Stopped" is black. One arrow, from working to stopped, and no arrow out of stopped.

Follow it. White, then black, then nowhere. Did two arrows meet? No, there's only one arrow. Did it use up new dots forever? No, it stopped. The answer settled and nothing gave.

Now ask it the obvious question: what does a stopped machine do on the next tick? Not "nothing happens," because something is still there. The usual answer is that it stays where it is. So give stopped an arrow back to itself. Now look again. Working points to stopped. Stopped points to stopped. Two arrows, one dot. The merge is back.

What if you don't want it to stay put? Point stopped at a new dot, "parked," and point parked at itself. Now stopped and parked both point at parked. Two arrows, one dot again. The merge moved one dot over. It didn't go away. That's the rocking machine from page one.

So a machine that stops only dodges this by not answering what happens next. Answer it any way you like, and you're back in the three cases: either two arrows meet, or you draw new dots forever.

1.6 What Turing actually proved

Almost every account you'll find says Turing proved the famous halting problem: that no machine can look at every other machine and say whether it stops. Since the stopping machine is right in the middle of this chapter, I went and looked at what he actually wrote, with an AI walking me through the paper and the links it turned up, and it isn't quite what happened.

Turing's own machines in 1936 weren't built to stop at all. The ones he cared about printed out the digits of a number, forever. What he proved no machine could always decide was a question about those: whether a machine keeps printing digits, and whether it ever prints a given symbol. The second one is a few lines away from the halting problem, so in a real sense he had it. But he never wrote it down that way. Stephen Kleene stated it in 1952. Martin Davis seems to be the first to have called it "the halting problem," in 1958. I found that in a paper where Joel David Hamkins and Theodor Nenu went back to Turing's 1936 paper and checked, line by line.

So the machine that stops, the one sitting right in the hole, is Turing's machine the way people later learned to read it. It's the one page one meant when it said Turing's machine already has something that settles and stays settled: whether it has stopped.

Put it together. Read a Turing machine through "has it stopped?" Give the stopped line a next line by letting it stay put. Then the step that stops it is a step where two arrows meet, which you just drew. That's the step I charge for. Charging for it is still my choice. That something gives there isn't.

The dots in this chapter could go on forever. The next question is what happens when they can't, when the paper runs out.

CHAPTER 2

Running out of room

The last chapter left one door open. A run that settles its answer and keeps it either has two arrows meet somewhere, or it lands on a new dot every step, forever. Endless fresh paper. On a sheet that never ends, that's a real way out.

But nobody has a sheet that never ends. So what happens when the paper is finite?

2.1 Four people, three chairs

Do this one with real chairs, or with coins on a table. It takes about a minute.

Draw a box on the floor and call it the done room. Put three chairs inside it and one chair outside, by the door. Sit four people down, one per chair. The people are the states the machine could be in. The three inside are already done. The one by the door isn't.

One move of the machine moves everybody at once: each person gets up and goes to whatever chair the rule sends them to. There are two rules about where they can go.

Nobody inside may end up outside. Done stays done.

The person by the door has to end up inside. This is the move that settles their answer.

Now try to finish the move with one person in each chair.

You can't. The three inside have to stay inside, so they fill the three inside chairs between them. The person from the door has to come in too, and there's no fourth chair. Somebody has to share. Two people, one chair: two arrows, one landing place.

Try it with more chairs. Ten inside, one by the door. Eleven people have to fit into ten chairs. A thousand inside, one by the door: a thousand and one people, a thousand chairs. It doesn't matter how big the room is, as long as it has a size. The insiders take every inside chair, and the newcomer has nowhere to go but somebody's lap.

Notice who ends up sharing. It doesn't have to be the newcomer. The rule could send the newcomer to an empty-looking chair and send two insiders to the same

chair somewhere else. Page one already showed you that: 4 went to 5a, and the arrows met a tick later on 5b. What's forced is that two arrows meet somewhere in that move. Where they meet depends on the rule.

Two words I'll need from here on. A move is one-to-one when two different starting places always end up in two different places: nobody shares a chair. A move that isn't one-to-one is a merge: two arrows, one landing place.

Now say what the chairs showed in those words. Take any move that keeps done states done and takes one state from not done to done, on a machine with K done states. If the move were one-to-one, the K people already inside would fill all K inside chairs with nobody sharing, and the newcomer would have nowhere to land. So it can't be one-to-one. It merges.

That's the whole argument, and it's the chairs. The only thing finiteness did was make "it fills all K done states" true: K people into K chairs with no sharing leaves no empty chair. That's the one place the size of the room mattered.

2.2 This is very old

The piece of mathematics doing the work there is old, and it isn't mine. Put more things into boxes than there are boxes, and some box gets two. People call it the pigeonhole principle. The earliest use of it I could find is Jean Leurechon in 1622, arguing that somewhere two people must have exactly the same number of hairs on their heads. The first name for it I could find came from Dirichlet, who used it in 1842 and later called it the drawer principle. It's the kind of fact a child accepts and a mathematician spends a whole career finding new uses for.

And the use I just made of it turned out to be old too. I found that out twice, and the second time stung more.

The first time was physics. In 1876 Josef Loschmidt pointed out, against Ludwig Boltzmann's account of how a gas settles down, that the laws of motion run just as well backwards: reverse every molecule and the gas retraces its steps. Twenty years later, in 1896, a young assistant of Max Planck's in Berlin pushed harder. He used Henri Poincaré's recurrence theorem from 1890, which says a system like that comes back as close as you like to where it started, again and again, to argue that nothing in it can climb forever. Give it time and it comes back down. The assistant was Ernst Zermelo, the same Zermelo who would put set theory on firm axioms twelve years later. Boltzmann answered the same year, not very politely, that you'd have to wait so long for a return that nobody would ever see one. (Jos Uffink, who wrote the Stanford Encyclopedia's entry on Boltzmann's physics, calls that reply drenched in sarcasm and bitterness, and calls Zermelo's paper a careful logical argument.)

You can see Zermelo's point on the dots from the last chapter. Draw a finite number of dots where no two arrows ever land on the same dot. Every dot has one

arrow in and one arrow out. Follow any run. Can it ever reach a dot it's been on before, other than the start? No: that would take two arrows into one dot. So since the dots are finite and it can't keep finding new ones, it has to come back to its start. Every run goes around in a loop. Now color the start white and some dots black, done stays done. Can a run go from white to black? If it did, it would have to come back around to its white start, and black can't go back to white. So it can't. A machine where no two arrows ever meet, on a finite sheet, can never settle anything new for good.

That's Zermelo's objection, with the infinities taken out. On a finite set you don't need Poincaré's theorem. And the wait isn't enormous; it's at most as many steps as there are dots. Zermelo was aiming it at the second law of thermodynamics, and settling an answer never came into it. A quantum version came in 1957, when P. Bocchieri and A. Loinger proved that a closed quantum system comes back close to where it started. That's a neighbour, not the same proof: even a small quantum system has endlessly many states, and you'd have to say how a record is read off it.

The second time was computer science, and that's the one that stung. Picture a small machine that reads letters, one move per letter. If every letter just reshuffles its states, no two arrows meeting, people call it a permutation automaton, and the sets of inputs such machines can pick out are called group languages. Jean-Éric Pin's 1992 paper on reversible automata sorts out exactly what those machines can recognise, and in that world the chairs argument is folklore: a machine like that can't recognise "has seen a b at some point," because recognising it would mean a done that stays done. Going quantum doesn't get around it. Alex Brodsky and Nicholas Pippenger showed in 2002 that a quantum version that only measures once, at the end, recognises exactly the same group languages. One that's allowed to measure after every letter, and stop on what it sees, does better, and "has seen a b" is one of the things it picks up. A measurement that can stop the machine is a record that stays. That's the extra thing that lets it in.

Then the closest one, and it's recent. In 2023 Thomas Ouldridge and David Wolpert worked out the physics of a small machine built the way real computers are: the same physical step run over and over. They sorted what such machines recognise into two kinds. If the smallest machine for a job has a letter where two arrows meet, every machine for that job does, and on their derivation running it gives off entropy that no cleverer design can get back. If the smallest machine only reshuffles, they find it can run with no unavoidable entropy at all. One of their worked examples is exactly "has not seen a b," which is "has seen a b" turned inside out. So the counting, and a physical price on it, were in their paper three years before this book. Their price comes out of the physics of the device. Mine, in the next chapter, is a choice I make and say so.

There's a cousin that speaks a different language. Constructor theory is David Deutsch and Chiara Marletto's way of stating physics as which transformations a device that runs in a cycle can and can't bring about, and in it a task is irreversible when it's possible and its reverse isn't. Marletto, Vlatko Vedral and their coauthors in 2022 built an irreversible task out of a machine of quantum bits that swap a little with each other, and ran it with photons. Maria Violaris and Marletto argued the same year that this machine, run in a cycle, can turn a pure quantum bit into a mixed one and never back, even though every piece of it can be reversed, and that this suggests erasing may cost more than the usual bound. The chairs are a small, exact, classical relative of that: a set of states the step can't leave, a task that's possible, and no way back. Theirs is quantum and approximate and aimed at physics. Mine is counted. Deutsch and Marletto's paper on information, from 2015, has another neighbour in its section 8.6: measuring one quantity disturbs another for good, and nothing can undo the disturbance while the result is still being kept. That's a second way of saying that a record that stays makes the step impossible to run backwards.

And Morgan Rogers, in his papers of 2019 and 2021 on what mathematicians call toposes of monoid actions, gives the general home for one more way to see it: if every move can be undone by some other sequence of moves, you can never enter a room you're not allowed to leave, because the undoing sequence would have to take you back out. On a finite set, a move where no two arrows meet can always be undone that way: repeat it enough times and everybody comes home.

So the counting isn't mine. It had homes in physics and in automata long before me, and a physical price on it a few years before me. What I do with it starts in the next chapter.

2.3 The hotel that never runs out

Now the way out. If the chairs run out, give the room more chairs. Endlessly many.

Take a strip of paper that never ends. Each move writes a 1 in the next empty box. Done means "there's at least one 1 on the strip." Check it against everything so far. Done stays done: once there's a 1, there's always a 1. The first move takes the empty strip to done. And no two arrows ever meet: two different strips always become two different strips, because you can always erase the last 1 and get back where you were. So here's a machine that settles something, keeps it, and never merges anything.

It got out of the chairs. It didn't get out of paying. Look at what every move does: it writes in a box nobody has used before. The first move does it, and so does every move after that, forever. That's the other column from the last chapter: fresh paper on every tick.

If the endless strip feels like a cheat, it's a famous one. David Hilbert told a story in a lecture course in the winter of 1924 to 1925 about a hotel with endlessly many rooms, every one of them taken, where a new guest still gets a room. Everybody moves from room n to room n + 1, and room 1 is left empty. Nobody shares. The one-to-one move leaves a gap. That's the strip.

The part worth checking is that Hilbert never published it. The hotel got famous because George Gamow put it in a popular book in 1947, and for a long time it wasn't clear Hilbert had told it at all. The historian of science Helge Kragh first guessed that Gamow made it up. Then Tilman Sauer, who had helped edit a volume of Hilbert's lectures, pointed him to Hilbert's lecture notes, which had just been printed in 2013, and in 2014 Kragh took his guess back in a paper called "The True (?) Story of Hilbert's Infinite Hotel." Keep the question mark. Even Kragh's paper puts the lecture in January 1924 in one place and January 1925 in another. Still, this time a story everybody repeated had a real lecture behind it. I've learned not to count on that.

Bring it back to the chairs. In the closed room the person from the door had nowhere to sit because the chairs ran out. In Hilbert's hotel they never do. So the whole argument was about running out of room, all along.

You can run the chairs backwards to see exactly how much room. Take a move that keeps done states done, settles one new state, and never lands two arrows on one place. Everybody already in the done room moves to a different done chair. The newcomer gets a done chair too, and nobody already inside is moving into it, because that would be two arrows meeting. So everybody already inside got seated in the done room with at least one chair to spare. With n chairs, that's n people into n - 1 chairs, which can't be done. So a move like that needs a done room with no end to it. Give the room any fixed size, and the move merges.

That's Hilbert's front desk run backwards. It's also exactly how Richard Dedekind defined "infinite" in 1888: a collection is infinite when you can match it one-to-one with a part of itself that leaves something out. So taking fresh paper isn't a way out of paying. It's the other way to pay, and only a machine with endlessly many states can keep paying it.

2.4 Cut the strip

Now cut the strip to eight boxes. Same rule: fill the next empty box from the left, never touch the earlier marks. There are nine possible strips, from empty to full. Write them down:

0 marks	`00000000`	next: `10000000`
1 mark	`10000000`	next: `11000000`
⋮	⋮	⋮
7 marks	`11111110`	next: `11111111`
8 marks	`11111111`	next: ?

After eight moves the strip is full. What should the ninth move do? Try each answer.

It stays full: then the strip with seven marks and the full strip both go to the full strip. Two arrows, one landing. The chairs.

It stops: then it has no next state, and that's the stopping hole from the last chapter.

It clears the strip: then done went back to not done.

Every answer gives up one of the things the chairs needed. None of them gets past the chairs.

Keeping the whole history like this is the trick Charles Bennett used in 1973, when he showed that any computation can be done without two arrows ever meeting, if you let it keep a copy of everything it did. Rolf Landauer, back in 1961, had already written that for a machine that runs a program that doesn't stop, the space to hold every step "cannot be there." Keeping the history is a real way to pay. It costs a memory that never fills, and every machine you can buy has a memory that fills.

Bennett wasn't the first there, or the last. Yves Lecerf showed in a short note to the French Academy of Sciences in 1963 that machines with no merging can imitate any Turing machine, and hardly anybody noticed. Tommaso Toffoli in 1980, and Edward Fredkin and Toffoli together in 1982, both then at MIT, built logic gates and circuits that compute anything and lose nothing, by carrying extra inputs in and extra outputs out. Their basic gate wasn't new even then: Landauer's own 1961 paper writes one down, calls it reversible and able to do an AND, and doubts anything like it could be useful. What Toffoli added was the proof that it computes everything. And Kenichi Morita proved in 1996 that even a tiny machine with two counters can compute everything with no two arrows ever meeting. Every one of them keeps what it doesn't forget somewhere: a history, extra outputs, a counter that codes the past. They all pay in fresh paper.

2.5 How long can a finite machine hold off?

One more question, because the 5a and 5b example on page one hinted at it. If the machine has a few spare done states, can it put off the merge for a while? Yes, and you can count exactly how long.

At three o'clock a screen says 42. The program is finished. Leave the camera running. At 3:01 it says 42. At 3:02 it says 42. The screen hasn't changed, but

the box behind it still has a next state every tick. It can stay exactly where it is, or something inside it can move while the screen stays put: picture a wheel turning from “first holding position” to “second holding position.”

Say the box has K different ways of holding the answer 42. Start one tick before it finishes. Suppose the run goes through its holding ticks without two arrows ever meeting along the way. Could two of those holding ticks be the exact same state? If they were, follow the arrows backwards from both, one tick at a time. Since no two arrows meet along the way, being on the same state now means they were on the same state a tick ago. Keep walking both back. Eventually the earlier one reaches the tick before it finished, the state that wasn’t done yet, while the later one is still on a done state. A state that isn’t done would have to be the same state as one that is: impossible. So every holding tick needs its own state, and with K of them, it can last at most K ticks. Within K + 1 ticks of finishing, two arrows meet.

You can build one that lasts exactly that long: the unfinished state, then K holding states in a row, the last one stepping to itself. The answer holds for K ticks with nothing merging, and on the next tick the last two arrows meet. A bigger memory puts off the merge. It doesn’t make it go away.

2.6 What’s being counted

One thing to keep straight before going on. Whether a move merges depends on what you count as the state. The endless strip keeps its whole history in the state, so nothing ever merges; cut it to eight boxes and the same rule merges. Keep a list of every earlier state alongside the current one, and any machine’s moves become one-to-one, because two different histories never look the same; a later chapter proves that. Keep only the state one step back, and it isn’t enough: two states that differ only in what came before them still land together.

So “this move merges” is always a fact about a move on a stated set of states. Every merge in this book is counted on the whole state the machine says it has, and nothing outside that state is counted.

What I’ve got now is counting and nothing else: on a finite machine, settling something for good means two arrows meet. The next chapter is about what I do with that, and which parts of it are my choice.

CHAPTER 3

Why charge for it

Everything so far has been counting, and the counting isn't mine. On a finite machine, settling something for good means two arrows meet somewhere. On an endless one, it can mean fresh paper forever instead. Nobody gets to choose that. You drew it.

Page one also said something I do get to choose: I charge for the move where the answer goes from not done to done. This chapter is about how much of a choice that is. If it's nothing but my preference, then the whole machine is a long way of telling you my favourite colour, and you should treat it that way. So I want to find out what it rests on.

3.1 A rule you can refuse

I'm not going to ask you to believe a price. I'm going to write one down as a rule, in the smallest form I can, and you can accept it or refuse it.

Any move where two arrows meet costs at least one.

That's all it says. It doesn't mention done or not done. It doesn't mention checking or settling. It puts a price on one thing: a move that sends two different states to the same place.

It isn't my rule. Charles Bennett, who spent decades working out what computing has to cost, put it this way in 2003: what has to cost is "the erasure of a bit or the merging of two computation paths." Two arrows meeting is two paths merging. I'm taking his rule as it stands and writing it on the move.

Now put it next to the chairs. The chairs said: on a finite machine, a move that settles something for good is a move where two arrows meet. The rule says: a move where two arrows meet costs at least one. Put those two sentences side by side and read them together. A move that settles something for good costs at least one.

That's the rule from page one, the one I said was my choice. I call it the toll. That was my word for it before I could prove any of this: there had to be a toll, a cost in the transition, in the step itself. And look at where it came from just now. Nobody handed it to the machine. It fell out of the chairs and Bennett's rule together.

So it isn't only my preference. But it isn't free of choices either. Count what went into it. There are three, and you can refuse each one.

The done states have to be finitely many, so the chairs run out.

Done has to stay done.

And two arrows meeting has to cost something.

Accept all three and you don't get to refuse the toll. It comes with them. Refuse any one, and the toll goes back to being my choice. So the question "did I have to pick it?" has turned into three smaller questions, and those are better things to argue about. For each one there's a machine that walks out through that door, and you should see each one, because a rule you can't picture breaking is a rule you're taking on faith.

3.2 The first door: endless paper

You've already seen this machine. It's the strip from the last chapter that never runs out. Each move writes a 1 in the next empty box. Done stays done, the first move settles it, and no two arrows ever meet.

So Bennett's rule never charges it anything. Two arrows never meet, so there's nothing to charge. The strip settles its answer at no cost on the price list.

But it didn't get out of paying. It pays in fresh paper, on every move, forever. A price list that only charges for arrows meeting just isn't looking at that column. And on any real machine, the paper runs out, and the chairs come back.

3.3 The second door: let it come back down

Take one coin. The only move is: turn it over. Heads is done, tails is not done.

Draw the arrows. Tails goes to heads. Heads goes to tails. Two dots, two arrows, and they never meet. So Bennett's rule never charges it.

And it does settle something: tails goes to heads, not done to done. At no cost.

What it broke is the second condition. The same move that took tails to heads takes heads back to tails. Done doesn't stay done. So it's not keeping anything. It's flipping a switch back and forth, and the page-one definition already throws it out: a Thiele machine's done never goes back.

Here's the part of this I found sharpest. Forget about done staying done entirely, and keep only the finite chairs. Take any move that takes some state from not done to done. Now go through the done states, one at a time, and ask: does this move send any of them back to not done?

If it does, you've found the coin: the move takes something back somewhere.

If it doesn't, then this move keeps every done state done. That's all the chairs needed, for this one move. So two arrows meet.

There's no third answer. On a finite machine, a move that settles something either has two arrows meet, or it takes something back somewhere else. A move that settles for free and never takes anything back doesn't exist.

That's narrower than "every record costs." It's still true, and I'll take it.

One more picture, so you don't think it's tidier than it is. Three cards, T0, T1 and T2, with a star on T1. The move sends T0 to T1, T1 to T2, and T2 to T1. Look at what it does. T0 gains the star: something settled. T1 loses the star: something taken back. And T0 and T2 both land on T1: two arrows meet. So Bennett's rule charges this move, and yet it doesn't keep anything for good. Charged moves and permanent moves aren't the same list. What's true is one direction: on a finite machine, every move that settles something for good is a move Bennett's rule charges.

And there's an exact way to say which moves Bennett's rule charges no matter what the rest of the price list looks like. A move where two arrows meet gets charged by every price list that follows the rule; that's what the rule says. A move where they never meet doesn't have to be: price that one move at zero and every other move at one, and the rule is still followed. So the moves the rule forces you to charge are exactly the moves where two arrows meet. That one holds on any machine, finite or not.

3.4 The third door: don't charge for it

Take a sheet of paper. It's either blank or stamped. One move: stamp it. Blank goes to stamped, and stamped stays stamped.

Check the conditions. Two states, so the chairs run out. Stamped stays stamped. Blank goes to stamped, so something settles. And draw the arrows: blank goes to stamped, stamped goes to stamped. Two arrows meet. Everything the chairs needed is here.

Now set the price of the stamp to zero.

Nothing in counting stops you. The stamp merges two states, and you just decided merges are free. The answer settled, and the bill says zero.

Price the stamp at one instead, and every condition holds at once, so the three can live together. That stamp at one is the smallest Thiele machine there is, and it gets its own chapter. But the third door asks whether they have to live together, and counting can't say so. The only thing that could close this door is a claim about the world: that two arrows meeting really does cost something, out there, in a real device.

The nearest such claim is Rolf Landauer's, from 1961: that forgetting gives off heat. It's measured for a few devices and argued over for everything else, and it gets its own chapter, with the people who doubt it named. I'll say one thing about it now, because it matters here. Bennett's rule as I wrote it charges every move where two

arrows meet, whether or not you already knew which state you were in. Landauer's heat, as that chapter will show, is only owed for what you didn't know. The rule I'm using is the worst case on purpose: a move that has to work whatever it's handed pays for the worst it could do.

So that's where the toll stands. Accept finite chairs, done stays done, and a price on two arrows meeting, and the toll isn't a choice; it's what the three say together. Each of the three is a choice, and I've shown you a machine walking out through each door. The physics that would close the third door is a premise, and I'll always call it one.

3.5 How much

The toll says at least one. Counting can say more, if you let it.

Take four cards and write A, B, C and DONE on them. Pick one without showing me. Now cover its face with a label that says DONE. Whatever you picked, the face now says DONE. Before the label, how many yes-or-no questions would it take me to find out which card you had? Two: "Is it A or B?" and then one more. After the label, I don't need to ask anything, because there's only one thing the face can say. Two questions' worth of difference got squeezed out.

That's four states landing on one. Each question you no longer need halves the number of places you could have been: four to two, two to one. Two halvings.

If you're willing to go one step further than Bennett's rule, and say each unit of cost pays for at most one halving, then that move costs at least two. And two is enough. In general: if m states were already done and the move settles k more, all m + k of them land on the m done ones, and the move has to pay at least as many halvings as it takes to get from m + k down to m. One newcomer next to one done state is one halving: the toll again.

(The label can be peeled off, so the cards only show the squeeze on what the face says. Whether anything physical got erased is a different question, for the heat chapter.)

There's a plainer way to say what the halving rule asks for. Look at one move and find the place where the most arrows land. Call that its worst pile-up. A cost of one allows piles of two. A cost of two allows piles of four. A pile of three already needs two. That's why the m + k states in the done room had to pay at least that much: spread m + k arrows across m chairs as evenly as you like, and the fullest chair still gets at least (m + k) / m of them. So the rule asks every move to pay for the worst it could ever do, whichever state it's actually handed.

That worst case has a home in physics. Philippe Faist, Frédéric Dupuis, Jonathan Oppenheim and Renato Renner worked out in 2015 the least work any logical step needs when it has to succeed every time, on every input, with no averaging. It

comes to exactly that logarithm of the worst pile-up. An AND gate sends three of its four inputs to 0, so it needs about 1.6 halvings' worth, two in whole units. My rule counts the same thing in units of cost. Joules don't come into it.

You can also count the whole run this way. Write N different starting states on cards. Run each card through the rule once, and stack the cards that land on the same state into one pile. Do it again with the piles. Each tick, count how many different states you're down to. The drops add up exactly: what you started with equals everything lost along the way plus what's left at the end. If you end up with fewer than you started with, some tick had two arrows meet. Turn the cards over first, though. Two runs can show the same answer on the front and still be different underneath, and those are different states.

A word about the k in all this. Physicists write kT for heat at a temperature, and when the counting meets the physics in the heat chapter you'll see k with a little B hung under it, for Boltzmann. That letter has a story I didn't expect. Boltzmann's grave in Vienna has $S = k \log W$ carved on it: entropy is a constant times the logarithm of the number of ways the parts could be arranged. The idea is his. But the constant, and the equation written that way, came from Max Planck, around 1900. In his Nobel lecture in 1920 Planck said the constant "is frequently termed Boltzmann's constant, although to the best of my knowledge Boltzmann himself never introduced it." So the most famous constant in this corner of physics wears the name of a man who never wrote it down. Take the constant away and what's left is the logarithm of a count, which is a number of halvings, which is what the cards just gave you. The constant is the exchange rate between halvings and heat. Whether the world honours that rate is the premise.

3.6 Found, assumed, chosen

That's the place to sort everything so far into three piles, because the rest of the book depends on where each thing falls.

I see it like this: I didn't invent a machine. I found one. That walks straight into one of the oldest fights in mathematics, so here's a short trip into it. G. H. Hardy, the Cambridge number theorist who wrote A Mathematician's Apology in 1940, came down on the found side about as hard as anyone ever has: he wrote that mathematical reality lies outside the people who study it, and the theorems they like to call their creations are really notes on what they saw. The German mathematician Leopold Kronecker supposedly split the difference: God made the whole numbers, and everything else is the work of people. Supposedly, because push on that line and it gets thin fast. It comes down through Heinrich Weber, in a memorial piece written after Kronecker died, pointing back to a lecture from 1886, and that's the only source I could find anybody citing. It's a great line with one witness.

None of that settles whether I found the Thiele Machine or made it up, and a proof wouldn't settle it either. So it's a conviction, and I'll call it one. What it changes is how I hold the thing: like something that was already there, which means I can be wrong about its shape and the shape won't care. And "found" only covers what counting forced. Here are the three piles.

Found: what counting forces on any machine.

Every step of a machine with one next state for each state lands somewhere new, circles back to its start, or has two arrows meet, and after it comes back it repeats forever. You drew that in chapter one.

A run that settles something and keeps it can't circle back to its start, so it has two arrows meet or takes fresh states forever. When only finitely many states are done, the move that settles it has two arrows meet. You sat people in chairs for that.

A machine that stops, with no next state, slips past all of it, and gets caught the moment you say what it does on the next tick.

Assumed: what the found pile is taken to stand on in the world.

That two arrows meeting in a real device, when the device really could have been in either state, gives off heat. That's Landauer's principle, and it's an assumption here, never proved, with its critics named in its chapter.

That a real device can only tell apart finitely many states in any bounded space, so its fresh paper runs out. Nobody has measured where that ceiling is.

That the machine starts clean: if done means "this happened," that's only because it started at not done. That start is put in by hand.

Chosen: what I add, each of which could have gone another way.

To put the price inside the step, as part of what a step is.

That done stays done. That's in the definition because I put it there.

The toll: at least one on the move that settles something. With finite chairs and done staying done, it follows from a price on two arrows meeting, and that price is a choice too.

Which thing gets to count as settling, and what the price list charges. Nothing I've shown you forces either one.

Everything after this chapter is built on those three piles, and I'll keep saying which pile each piece comes from.

CHAPTER 4

The smallest one

The last chapter put a price on the move that settles something. Before I build the smallest machine that carries that price, there's a question I had to ask first. People have been making marks that other people are supposed to rely on for a very long time. Did any of them put a price on the moment the mark goes on and stays? I wanted to know what they thought they were doing, and what it cost them. So here's what I could reach, and where the search ran out.

4.1 Clay, seals, and seven witnesses

Start a little over five thousand years ago, in the Near East, with people who counted sheep with little clay tokens. To send a count across a distance without the courier changing it, they put the tokens inside a hollow clay ball, pressed the shapes of the tokens into the outside while the clay was soft, and sealed it. The archaeologist Denise Schmandt-Besserat has spent a career on these, and she puts the practice at around 3300 BC; the ball is called a bulla.

You can do a version of it with a cookie tin. Glue a paper seal across the lid and the rim, let it dry, and then try to open the lid without tearing the seal. The bare lid would come straight off. The seal is what turns opening it into something you can see afterwards. A substitute seal could fool you, so the tin shows that somebody disturbed it, not that nobody could ever fake it. The clay ball does the same: the outside shows what's inside, and opening it breaks it. Keep that idea, because it comes back: a record is worth something only if it can't be quietly undone.

Now law, which is where people get serious about who's allowed to say what happened. The book of Deuteronomy, chapter nineteen: nobody is convicted on the word of one witness; it takes "two or three." The Roman jurist Gaius, writing in the second century, says a will the court will honour is one sealed by seven witnesses. Why seven? Gaius doesn't say. The legal historian Henry Maine, in 1861, had an answer: the older Roman will was a mock sale, done in front of five witnesses, a man holding the scales, and the buyer of the estate, seven people in all, and when the ceremony fell away the seven seals kept the count. Maine's is the only account of that I found, so treat it as his.

Try it. Write “paid five dollars, signed A” on a napkin and ask a friend to forge it. It takes them ten seconds. Now have two other people initial it at the moment of payment and each keep a copy. If whoever reads it later compares those copies and trusts the people holding them, changing your napkin alone gets caught. Stolen copies, forged initials or a reader who never compares them can still beat it. The claim didn’t change. What stands behind it did.

Which brings me to seals, and the medieval historian Michael Clanchy, whose book is called From Memory to Written Record. England between the Norman Conquest and the early 1300s was learning to trust writing. Until then it had trusted people who remembered. The lesson was painful: seals were forged, charters were forged, and historians today have a whole craft of telling real charters from fakes because there are so many fakes. A seal is only as good as whoever holds the stamp. In 2011 a company called DigiNotar, which issued the certificates web browsers trust, found out the same thing at the cost of its existence: a mathematically perfect signature is worth exactly what the custody of the key is worth, and intruders had taken over the machines that signed. It’s the same failure nine hundred years apart, and I find that oddly comforting.

And then the obvious next question: if seals could be forged, what caught the forgeries? It wasn’t the seals. The most famous forged record in European history is the Donation of Constantine, a document in which the emperor Constantine supposedly handed rule over Rome and the western empire to the Pope. It was most likely written in the eighth century, some four hundred years after Constantine died, and it was leaned on for centuries. Around 1440 a scholar named Lorenzo Valla took it apart by its Latin: the language belonged to a much later age, and it used words, “satraps” among them, that no Roman of Constantine’s day would have used for his officials. (Valla wasn’t even first. Nicholas of Cusa, a German churchman who later became a cardinal, had already called it a forgery in 1433. I know that from the reference works; I haven’t read his book. Valla’s is the version people remember, which is a pattern that keeps turning up.)

Look at what kind of check that was. The question wasn’t “is the stamp real.” It was “could these words have been written when they say they were.” A record can pass the first and fail the second. That was my first hint that what I kept calling one thing, a record, was doing more than one job.

4.2 Two columns and a stamp

The one I’d never have guessed mattered is bookkeeping. Around 1300, the ledger of a Florentine company called Farolfi, kept at its branch in Provence, shows the earliest complete example anyone has found of a trick merchants were working out: write every transaction twice, once where the money came from and once where

it went, and the two columns have to balance. The treasurers of the city of Genoa kept their ledger for 1340 that way. And in 1494 a friar named Luca Pacioli printed the method in a mathematics book, and it spread.

Do it for one day. Keep two columns of every dollar that moves, and total them at night. Have a friend secretly change one entry in one column. The totals disagree; you catch it. Now have them change the matching entry in both columns. The totals agree, and you've caught nothing. So that's what double entry actually is: a check for slips, which a careful liar walks straight through.

Did Pacioli see that? I can only get at it secondhand. A scholar named Forrester, in a paper on the University of Strathclyde's accounting-history pages, reads him as reporting that in many places the books were taken to the local consul to be stamped before a single entry went in. The stamp is a witness again, from outside. The two columns are a method. The stamp is somebody else agreeing to stand behind the fact that these are the books. Keep those two apart, because one is a habit and the other is a commitment.

Then I found where this got turned into a motto. The Royal Society, founded in London in 1660 and still running, picked as its motto Nullius in verba: take nobody's word for it. Where did they get it? From somebody's words, naturally. It's half a line of the Roman poet Horace, from a letter to his patron, saying he isn't bound to swear by the words of any master. So the society that takes nobody's word for it took its motto from a poet and cut the line to fit. That's not a contradiction. It's a nice picture of the method: borrow what's useful, say whose it was, and don't swear by it.

The historian Steven Shapin wrote about what that motto meant in practice, and the answer is more interesting than the slogan. Robert Boyle, pumping the air out of a glass globe to see what happens to a candle or a bird, had three problems: the machine, the report, and the witnesses. He solved the third by having gentlemen in the room who'd sign their names, and the second by writing the report so richly that a reader who wasn't there could picture the scene and count as a witness anyway. Shapin calls that virtual witnessing, and he notes, dryly, that the air pump was very expensive, the big science of the 1600s. So there was a cost. It just wasn't a cost of declaring the fact. It was the cost of the instrument and of getting enough people into the room. Hold on to that difference, because the rest of the book hangs on it.

4.3 Closer to now

The rest of the lineage points the same way, so I'll go through it quickly.

Regulators who made stockbrokers keep their records on media that can't be rewritten (the SEC required exactly that, write once and read many, from 1997 until

2022, when it allowed an audit trail instead) were paying for the hardware. The rule put no charge on the act of writing.

Stuart Haber and Scott Stornetta, in 1991, chained time-stamped records together so that changing an old entry breaks every entry after it. Bitcoin's footnotes lead back to them. Try the idea with a running total. Take five entries, 1, 2, 3, 4, 5, and add each to the total so far: 1, 3, 6, 10, 15. Now change the second entry to 3. The totals become 1, 4, 7, 11, 16. The change travels all the way down. Now also lower the third entry to 2, and the last total is 15 again. Simple addition can be fooled by changes that cancel out. A real chain uses a much harder function than addition, and somebody has to keep a trusted copy of the last value to compare against. (Ralph Merkle's hash tree, from his 1979 Stanford thesis, lets one short value stand for a great many records the same way.) What the running total shows you is how each entry depends on the ones before it. It doesn't put a price on writing one.

Proof of work, which Cynthia Dwork and Moni Naor proposed in 1992 to make junk mail expensive, and which Bitcoin later chained, is the one place where a record really does cost something. But what it charges for is a search: how many tries it took to find a lucky number. An empty block and a full one face the same target. The work pays for the search. Nothing extra gets charged for the moment anything becomes settled.

Robert Floyd and Tony Hoare, asking in 1967 and 1969 how you'd prove a program does what it's supposed to, attached claims to points in a program: "whenever you get here, this is true." The claim is backed by a proof, and the proof is paid for somewhere else, off the machine.

The certificates of complexity theory, going back to Stephen Cook in 1971, are answers designed to be quick to check: hand me the certificate and I can confirm the claim fast. Whether they're also hard to find is the famous open question of the field, P versus NP, and nobody knows. Either way, what's bounded is the checking. Nothing is charged for raising a flag.

And from statistics, a ledger that runs the other way. Dean Foster and Robert Stine's alpha-investing, from 2008, runs a long string of tests against a budget of allowed false alarms: every test spends some of the budget, and every test that finds something earns some back. That's a budget for how often you're allowed to be wrong. A reward there doesn't mean the discovery was true.

4.4 Five jobs

Lay them all down together, and what shows through isn't one thing. It's five different jobs the word "record" has been doing, and I'd been letting one word carry all five. Most of them come with a corpse.

Durability: the record can't be quietly undone. The clay ball, the write-once disk and the hash chain do this one. The corpse is the editable log, and the company that kept its debts off the books: Enron hid debt in side companies set up to keep it off the balance sheet, which is how the Justice Department described it in 2002, and its auditor, Arthur Andersen, shredded paper. Andersen was convicted that year, and the Supreme Court threw the conviction out in 2005 over how the jury had been instructed. Notice that this job doesn't need a witness at all. A copy will do, as long as somebody else holds it.

Authenticity: the record came from who it says. Seals, signatures and the consul's stamp do this one. The corpse is the forged charter, and DigiNotar. A perfectly authentic entry can still be erased, and a perfectly durable one can still be a forgery. They're different jobs.

Evidence: enough independent people stand behind the claim that lying gets expensive. Two witnesses, seven seals and Boyle's gentlemen. This one is about how many people put their own risk next to the claim. Paper has nothing to do with it.

Accounting: the record checks itself for slips. The two columns. And the second column isn't an outside witness. It's you, checking you. It catches a dropped digit, and a careful liar gets past it.

Quick checking: a certificate, signature or fingerprint that makes one particular check fast. Checking doesn't have to be cheaper than making, and I once thought it had to. Plenty of records are checked by doing the whole thing over, and that's fine when the thing is small.

Now the thing none of them says. Not one of those jobs puts a price on the moment a record goes on and stays. Proof of work prices a search. Bookkeeping prices a habit. Witnesses and seals price showing up and guarding the stamp. I looked, with everything I had, and that's where it ran out. If somebody priced that moment itself, they're on a shelf I never found.

So I had to try it myself, on the smallest thing I could build.

4.5 Two boxes and a coin

Take a sheet of paper. Draw two boxes and label them NO and YES. NO is not done, YES is done. Put a coin in NO. Draw a line underneath for tally marks; that's where you keep score.

You get one move. Call it the stamp. From NO, the coin slides to YES. From YES, it stays in YES. Every time you make the move, whichever box you're in, you put down one tally mark.

Play. Start in NO, make the move as many times as you like, and look at the tally. Now try to get to YES with no marks at all. You can't, and you can see why: the only way from NO to YES is the stamp, and the stamp makes a mark.

Now cross out the mark in the stamp's rule and play again. You get to YES with no marks, every time. That's how little the mark is. The board doesn't force it. Chapter three is the reason it's there: finite chairs, done stays done, and Bennett's rule. Here it's written on a card.

Draw the arrows for the stamp. NO goes to YES. YES goes to YES. Two arrows, one landing place. It's the same picture as 4 and 5 both going to 5 on page one.

That's a Thiele machine. A thing (where the coin is). A move (the stamp). A way to tell how settled it is (NO or YES), where YES never goes back to NO. A run that actually settles it (start in NO, stamp once). A price on the move (one mark, never a refund). And the toll holds: the move that takes NO to YES costs one.

Is there a smaller one? Try.

One box? Then there's nothing to settle. The coin is always in the same place, so no run ever takes it from not done to done.

Two boxes, then. A move either sends the two boxes to two different boxes, or it sends both to the same one. If it keeps them apart, there are only two ways to do that: leave each where it is, or swap them. Leaving them where they are never settles anything. Swapping them is the coin from the second door: it takes NO to YES, but it also takes YES back to NO, and done has to stay done. So a two-box machine that settles anything has to have a move that sends both boxes to the same box, and since YES has to stay YES, that box is YES. That's the stamp.

So nothing smaller works, and look at what that means. Chapters one and two found that keeping something takes two arrows meeting or endless fresh paper. Here there are only two boxes, so there's no fresh paper to be had, and the arrows meet. The smallest machine there is already has the merge inside it, in plain view. Nobody added it. A fuse that blows, a bead that drops into a dip and stays, a stamp pressed onto a box: each one is two states and one move that sends both to the done side, and each one, with that move priced at one, is this machine.

I'll call it the latch, because that's what it is: something that catches and holds.

4.6 What the tally is and isn't

Why tally marks? Because they're about the oldest receipt there is. The real ones, notched sticks split down the middle, get their own chapter later, fire and all. The tally on your paper is nothing like those. Nobody holds the other half. You keep it, beside the board, and you could rub a mark out and nobody would ever know.

Hold on to that. Of the five jobs, the latch does a small piece of the first one: none of its moves takes YES back to NO. It has no witnesses, no signature, no second column, and nothing in it checks anything. YES means the coin is in YES. That's all it means. Whether a YES can ever be made to mean "this was checked and can be relied on" is a real question, and it's the one the second half of the book is about.

For now the tally is a scorekeeper's receipt: you know what you paid, and nobody else has any reason to believe you.

4.7 Any machine can wear one

Here's something the latch lets you do to any machine at all.

Take any machine, any states, any moves. Pick one state, call it x, that some move lands on. Now build a new machine right beside it: the same states and the same moves, with one extra coin riding along in its own two boxes, NO and YES. The coin starts in NO. The first time a move lands on x, the coin goes to YES, and it never comes back. Price every move at one.

Check it against the definition. The coin never goes back to NO. Start anywhere the move that lands on x can be made, with the coin in NO, and make it: the coin goes to YES. Every move costs one, so the toll holds. It's a Thiele machine.

Now cover the extra coin with your hand and watch the rest. What you see is the machine you started with, move for move. Nothing about it changed. The coin rode along and the old machine never knew.

And with your hand over the coin, you can't tell which box it's in. Let any spot with the coin in NO count as a place to start. Now find a moment where the old machine is sitting on x. Did it just arrive there, which put the coin up? Or did it start there, with the coin still down? Both can happen, and the part you can see looks exactly the same. The coin is the only place the difference lives.

So every machine that has a move at all is what's left of a Thiele machine once you cover the coin. Even a machine with one state and one move that goes nowhere: the move lands on that one state, so the coin goes up, and what you've built is the latch. Turing's machine read through "has it stopped?" doesn't even need the extra coin: chapter one showed it already has a done that stays done, and stopping is already a step where two arrows meet. Both ways, it points the same direction page one did. The machine with the price in it is the general thing. The computer is that machine with the price left out.

4.8 Which question gets charged?

One more puzzle, because it shows why the definition picks one question and charges only that one.

Put a tile on a table and slide it one place to the right. Let that move be free. Now ask a new question: is the tile in the place it just reached? Before the slide, no. After it, yes. If every question had to follow the toll, that slide would have to cost at least one.

Try it with any free move that changes anything. Ask “is the machine where this move just put it?” The answer goes from no to yes. So if the toll applied to every question you could ask, every move that changes anything would cost something, and nothing could ever be free. That’s why the toll belongs to one question, the one the machine says is “how settled,” and not to every question you could think of.

And for that one question, a free move can’t touch it. It can’t take NO to YES, because the toll charges that. It can’t take YES to NO, because done stays done. So a free move leaves the settled question exactly where it found it. That’s the whole reason the free moves and the settling moves come apart so cleanly.

4.9 The cheapest price list

Last thing about the latch. Put three price cards next to a button: zero, one and two. Somewhere in the machine, pressing that button takes NO to YES. Cross out zero; the toll won’t allow it. One works. Two works too. The toll picked the smallest card it could, without forbidding the bigger ones.

Now take a button that never takes NO to YES, from anywhere. Zero works. So go through the buttons: one on each button that can settle something somewhere, zero on each button that never does. That’s the cheapest price list that follows the toll, and there’s only one cheapest. Notice the price stays on the button: the stamp costs one even when you press it on YES and nothing changes. The toll is a floor, not an exact bill.

And it counts. Every time a run goes from NO to YES, that step costs at least one, so a run costs at least as much as the number of times it settled something. On the latch that’s at most once, because YES stays YES. On a switch that can be turned off for free, you could settle it and unsettle it a hundred times, and you’d pay at least a hundred. A latch pays once, and keeps it.

CHAPTER 5


More settled

Up to now everything has had two places: not done and done, NO and YES, the coin in one box or the other. Page one promised more than that. It said settled doesn't have to be yes or no, and to picture a checklist.

That promise matters to me more than it might look. The hunch this whole thing started from was that there's another direction, one ordinary computing doesn't look along. A direction isn't a switch. A switch is just the first line drawn across it. So this chapter takes the switch and finds out what happens when there are many lines.

5.1 A row of cups

Get four cups and a strip of paper. Write five stages of an experiment along the strip, left to right: proposed, set up, run, read, recorded. Agree to do them in that order. Put an upside-down cup between each pair of neighbours. The first cup means "the setup happened," the second "the run happened," the third "the result was read," the fourth "the result was recorded." When you finish a stage, turn its cup right side up. Cups only ever get turned up. None of them ever gets turned back down.



Read down the drawing. Zero means cup down, one means cup up. Four cups give five stages along this run because each new stage turns up one unused cup.


Two things to notice with the cups in front of you.

Each cup on its own is the latch from the last chapter: down or up, and once up, it stays up.

And four cups were enough for five stages. Try it with three. Turn them up one at a time, never back down, and count how many different arrangements you pass through. None up, one up, two up, three up. Four arrangements, and you needed five. You can't do it. Every time the arrangement changes, at least one more cup goes up, and with three cups that can only happen three times. So three cups give you at most four stages, and in general a row of cups that only go up can tell apart at most one more stage than it has cups. A straight line of stages needs one cup fewer than it has stages, and no fewer.

An ordinary binary counter holds a thousand stages in ten bits. That doesn't break anything here: a counter's bits go down as well as up. These cups don't.

Now a harder strip. Some checks don't line up. One person repeats the experiment. Somebody else, separately, works through the calculation. Neither one includes the other. Draw a diamond: "neither" at the bottom, "experiment repeated" on the left, "calculation checked" on the right, "both" at the top. There are two ways up, and the two middle points don't come before or after each other.



Read the cups as repeat, calculation, both. Neither middle point includes the other. The third cup repeats information already supplied by the first two; it is useful for showing thresholds, not needed for storage.

Put a cup on each point except the bottom: one for the repeat, one for the calculation, one for both. Each cup is still a latch. The cups tell you which things have happened. They don't tell you whether the experiment was set up right or the calculation started from the right place. That's a different question, and it comes later.

5.2 Naming what's on the table

Now that you've seen the pieces, here are the names I'll use for them.

The set of places the record can be, whether it's a straight strip or a diamond or something with a hundred points, I'll call the direction.

Each question of the form "has the record reached at least this place?" I'll call a line across it. One line is one cup.

To say "at least this place," you need to be able to compare two places: is this one at or below that one? The strip and the diamond let you see that by looking. Said exactly, it's a test you can put to any two places, and it has to follow two rules. Every place is at or below itself. And if one place is at or below a second, and the second is at or below a third, then the first is at or below the third.

There's a third rule you might expect: if two places are each at or below the other, they're the same place. Every strip and diamond on the table follows it. To see what a rule like that is doing, take it away.

So take two places, left and right, and say each one is at or below the other, without them being the same place. Now there are two cups: "at least left?" and "at least right?" Put the record on left. Both cups are up, because left is at or below

itself and at or below right. Put it on right. Both cups are up again. Look at the cups and you can't tell where the record is.

Now let the record swap between left and right on every step, for free. Is that allowed? It never goes below where it was, since each is at or below the other. No cup ever moves, so nothing ever settles, so the toll never asks for anything. The record moved on every single step, and nothing on the table noticed. It never settles anything either, so it isn't a Thiele machine.

Put the third rule back, and the cups tell you exactly where the record is. Suppose two places leave exactly the same cups up. The cup for the first place is up when the record sits on the first place, so it's up on the second place too, which says the first is at or below the second. The same goes the other way round. Each at or below the other: by the third rule, they're the same place. Without the third rule, the cups can miss moves. Neither is wrong. Dropping the rule just tells the cups not to care about moves between places on the same rung, and they don't.

5.3 The toll on the whole direction

One cup had one rule: the step that turns it up pays. Now there's a row of cups, or a diamond, and one step can turn up several at once. Here's the rule I'd guess first, and it turns out to be the right one: a step that turns up any cup pays at least one. Three cups at once still only has to pay one. The rule asks whether a step turned something up, not how many.

Does that still give you what one cup gave you? Take any run where some cup is up at the end that was down at the start. Somewhere along the run, one step turned that cup up, and that step paid. So the run paid at least one. And every step that turned anything up paid, so the run paid at least once for each of them.

Look at what that argument never used. It didn't care whether the cups sat in a row or a diamond. It didn't care how many cups there were. It didn't even use the fact that cups never come back down: if something could knock a cup back over, the argument still works, because all it needed was the one step where that cup went up.

There's a simple way to tell, from places alone, when a step turns up a cup. Say a step takes the record from place x to place y. The cup at y is up afterwards, because y is at or below itself. Was it down before? It was down exactly when y isn't at or below x. And if y is at or below x, every cup that's up at y was already up at x, so nothing turned up. So a step turns up some cup exactly when it lands somewhere that isn't at or below where it started. That's the toll from page one, written on any direction.

And checking it one cup at a time is the same as checking it on the whole thing. If every cup on its own follows the two-box toll, the whole direction follows the toll, and the other way round.

5.4 One cup isn't enough

Could one cleverer cup do the job of all four? Say you keep one latch and try to read the record's place off the latch together with whatever else the machine shows.

Take the smallest case: a machine with nothing else to show, and a record that climbs through three places, bottom, middle, top. One latch gives you two answers, down or up. Three places, two answers. Some two places get read the same. One cup can't do it.

Try a second guess, closer to how a single cup works. A cup's rule is: the new value is the old value, or the event that turns it up. For a whole row, the obvious copy is: the new place is the lowest place that's at or above both the old place and the event's place. Test it. Take the strip bottom, middle, top, with a cup at middle and a cup at top, and a record that jumps from bottom straight to top in one step, and stays put at middle. With nothing else for the event to look at, the event has to be one fixed place. From bottom, it has to take the record to top, so the event is top. But then from middle, the same rule takes the record to top too, and the record was supposed to stay at middle. The rule breaks.

What works is giving every cup its own event, and letting each cup's event look at where the record already is. A later chapter builds that out over any machine.

How many cups does a direction need? That depends on what you ask the cups to do, so start with the smallest job: tell where the record is, using cups that only ever go up as the record climbs. For that job a cup doesn't have to be "at least this one place." It can go up at several places, as long as climbing never knocks it back down.

A straight strip needs one fewer cup than it has stages; you counted that above. A direction that branches needs at least as many cups as the steps in its longest climb, and at least enough to tell all its places apart.

Put three places above a bottom, none of them above another. The longest climb is one step, but one cup can't tell four places apart, so it takes at least two. Two do it, if you let a cup go up at more than one place: one cup that goes up at the first and third places, one that goes up at the second and third. Every place gets its own pattern. But look at those two cups alone and the third place seems to sit above the first, because every cup that's up at the first is up at the third too. If you also want the cups to show which places sit above which, it takes three: one "at least this place" cup for each of the three.

And sometimes just telling the places apart needs more than both of those. Take two places, each below two others. One step of climb, four places, and two cups still won't do it, because both upper places would have to keep up every cup either lower place has up, and two cups don't leave room for that. It takes three, and showing which sits above which takes four.

5.5 Somebody got here first, from the side

People got here before me, from a direction I didn't expect: programming for many processors at once. Lindsey Kuper and Ryan Newton's LVars, from 2013, are memory cells whose value only climbs, where a program can't ask for the value itself, only whether it has climbed past some point. That's a cup. Two writers running side by side can push the value up in different orders, and a point once passed stays passed. Their programs come out the same every time partly because of which questions the language lets you ask: a program allowed to ask whether the cup is up yet could still see the other writer's timing.

Amin Timany and Lars Birkedal, coming at it from proofs about programs, show in their 2021 paper, as I read it, that any order at all can be carried this way in a logic for proving programs correct, as a value that only climbs and that the running program never reads. That's the closest I've found to somebody else cutting a direction into lines.

The two of them rest on the same shape, a value that only climbs and a question that only asks whether it's past a point, and they use it for opposite jobs. In an LVar the question steers the program. In Timany and Birkedal's logic, the value lives in the proof and the program never looks at it. The shape is theirs first. What I can't find in either is a price on crossing a line.

5.6 Why crossing a line has to cost

Chapters two and three got the toll out of the chairs: on a finite machine, a move that settles something for good lands two states on one. Does that survive a whole row of cups?

The chairs need one yes-or-no question to count with, and now there are many cups. So which cup? The first one I'd reach for is the cup at the place where the record starts. That goes nowhere: the starting state already has that cup up, so there's no newcomer coming in from outside. The cup that works is the one the step turns up, the cup at the place where the record lands. It's up afterwards and it was down before. So put everybody who has that cup up in the done room, and the state the step starts from is the person by the door.

Now it's the chairs again, once for that cup. If the move keeps everybody in the room inside and never lands two of them on one chair, they fill every chair, and there's none left for the newcomer. So one of two things happened: two arrows met, or somebody inside got moved out, which means that cup went back down. If cups never come back down, it has to be the first one. Two arrows meet.

And then Bennett's rule from chapter three does the rest. A move where two arrows meet costs at least one, so a move that turns up a cup costs at least one. The toll on the whole direction rests on the same three things as the toll on one cup: finitely many states with each cup up, cups that never come back down, and a price on two arrows meeting.

Each of those can be knocked out, and each one goes the same way it did for one cup. Let the states be all the whole numbers and let the record count up by one on every step, free: nothing ever merges, the price list has nothing to charge, and every step turns up a new cup. That's the endless strip, paying in fresh paper. Let a two-state record swap back and forth: nothing merges, and the cup it turns up gets knocked back down. That's the coin. Let a two-state record stamp every state up and charge nothing: the arrows meet and nobody asked them to pay. That's the stamp at zero.

So nothing new got smuggled in when one cup became a row, and nothing the one cup needed got left behind. You might expect the diamond to need some extra rule for the two middle cups that don't come before or after each other. It doesn't. Every cup gets its own room of chairs, and the argument in each room is the old one.

CHAPTER 6

Looking isn't keeping

The toll charges the move that settles something for good. But there's another thing a machine does that feels like it ought to cost at least as much, maybe more, and that's finding out. Every time a machine rules out a possibility, it knows more than it did. If two arrows meeting costs something, does learning cost something too?

Most people's gut says yes, and finding out is the expensive part. Most physicists felt that way too, for a long time. It's why one little imaginary creature kept them busy for over a century. This chapter is about where that feeling goes wrong, and where it's right. The short version: looking doesn't have to make two arrows meet. It uses up a blank page instead, and you can hand the page back. Wiping the page to use it again is where two arrows meet. And making a record that stays, one nobody can hand back, is paying for good.

6.1 A small being at a small door

In December 1867 James Clerk Maxwell, who had published his equations for electricity and magnetism two years before, wrote a letter to his friend Peter Tait about a problem he couldn't get rid of. Take a box of gas at one temperature, with a wall down the middle and a tiny door in the wall. Now imagine, he said, a "very observant and neat-fingered being" sitting at the door, who can see the molecules coming. When a fast one comes from the left, it opens the door and lets it through. When a slow one comes from the right, same. Otherwise it keeps the door shut. It never pushes anything. It only opens and closes a frictionless door at the right moments.

After a while the right side is hot and the left side is cold, and you've got a difference you could run an engine off, made out of nothing but looking. That's supposed to be impossible. It's exactly what the second law of thermodynamics says can't happen, and the second law is the one law physicists are least willing to give up. Arthur Eddington wrote in a 1928 book of popular lectures that a theory found to contradict it has nothing for it but to collapse in deepest humiliation.

Maxwell put the being in his textbook in 1871, and it got named Maxwell's demon, though not by Maxwell. William Thomson, later Lord Kelvin, called it a

demon in the journal Nature in 1874. Maxwell, in an undated note to Tait headed "Concerning Demons," tried to take the name back: "Call him no more a demon but a valve." Nobody listened, which is probably for the best, because "Maxwell's valve" wouldn't have lasted a century. But his correction is exactly right, and it's the whole problem: the being does no work. It opens and shuts a door.

So physicists spent the next century trying to find where the being was cheating. The obvious answer pulls hard: the being has to see the molecules, and seeing costs. In 1929 Leo Szilard cut the box down to a single molecule and did the arithmetic. The second law survives, he said, if finding out which side the molecule is on costs at least as much as the being gains. In 1951 the French physicist Léon Brillouin made it concrete: to see a molecule you have to bounce light off it, and, the way the philosopher of physics Jeffrey Bub sums it up, finding out that way costs at least one halving's worth of entropy for every bit. For thirty years that was the answer. Information costs because getting it costs.

But that's a price on one way of looking. It doesn't show that every way of looking has to pay it.

Szilard is worth one more minute, for a reason that has nothing to do with the demon. Four years after the one-molecule engine, in September 1933, by his own later telling, he was waiting at a traffic light on Southampton Row in London. He had just read a newspaper report in which Ernest Rutherford, the physicist who announced the atomic nucleus in 1911, called the hope of getting power out of atoms moonshine. As the light changed, it occurred to him that an element which gave off two neutrons when it swallowed one could keep a reaction going by itself. That's the nuclear chain reaction, thought up crossing a street. It's his memory of himself, and Richard Rhodes's history of the bomb opens with it, in a chapter called "Moonshine," so hold it as the man's own story, not a record made at the time. It changes nothing about the demon. What it adds is a picture of the man: the same mind looked at one molecule in 1929 and one neutron in 1933, and both times the question was what a single small step forces downstream.

6.2 Bennett moves where the cost is

In 1982 Charles Bennett, who had already shown in 1973 that computing can be done with no two arrows meeting if you keep the whole history, took the demon apart one step at a time.

The being looks at a molecule and writes down which side it's on. Draw it. Before the look, there are two ways the world could be: molecule on the left with a blank notepad, or molecule on the right with a blank notepad. After the look, there are two ways: molecule left with the pad saying "left," or molecule right with the pad

saying "right." Two went in, two came out, and you can read the before off the after. No two arrows met.

But something got used up: the blank page. Before the look, the pad was blank and ready. After it, the pad is full. That's the fresh paper from chapter one. The look didn't merge anything. It borrowed a page.

So where's the cost? The being has one notepad and a lot of molecules. After the first molecule the page is full, and to do the next one it has to wipe it. Draw the wipe. Pad says "left" goes to blank. Pad says "right" goes to blank. Two arrows, one landing place. There it is. Looking feels like the active part, the thing the being does, and wiping feels like housekeeping. The housekeeping is where two arrows meet.

That's Bennett's, and it's the same rule chapter three wrote on the move: what has to cost is "the erasure of a bit or the merging of two computation paths." In the same 2003 paper he lists copying onto a blank place among the operations that can, in principle, be done without any such cost.

6.3 The demon on a sheet of paper

Build it, because the whole thing is four dots, and you should see the arrows meet or not meet for yourself.

The machine has a hidden bit, which is the molecule's side, and a display, which is the notepad. Write its four states as pairs, hidden bit first and display second: 00, 01, 10, 11. It has two moves.

MEASURE copies the hidden bit onto the display the way a light switch adds: if the hidden bit is 1, flip the display; if it's 0, leave it alone. Draw the arrows. 00 goes to 00. 01 goes to 01. 10 goes to 11. 11 goes to 10. Every state has exactly one arrow landing on it. Nothing met.

Look at the two states where the display starts blank, 00 and 10. They go to 00 and 11: the hidden bit got copied onto the blank display. That's the look.

WIPE sets the display to 0 no matter what it held. 00 goes to 00, and 01 goes to 00 as well. 10 goes to 10, and 11 goes to 10 as well. Two arrows land on 00 and two land on 10. That's the wipe, and arrows meet twice.

Hidden bit, display before	After MEASURE	After WIPE
00	00	00
01	01	00
10	11	10
11	10	10

So Bennett's rule charges WIPE at least one, and it's allowed to charge MEASURE nothing. Zero on the price list doesn't mean zero spent in a real device; a real MEASURE takes time and energy, and it uses up the blank display. What it doesn't do is make two arrows meet.

One detail worth seeing. Follow only the two cases you actually set up, the ones with a blank display: after MEASURE they're 00 and 11. WIPE sends those to 00 and 10. Still two different states, because the hidden bit still tells them apart. So on those two cases, WIPE didn't lose anything. The rule charges it anyway, because the rule has to answer for every state it might be handed, and on the full table the arrows meet. The price is the worst case on purpose.

Now be the one watching. You can see the display and not the hidden bit. The display starts blank, so you've got two candidates for what's going on: 00 or 10. One bit you don't know. The machine runs MEASURE, and you look. If the display shows 1, only 10 could have done that. If it shows 0, only 00. Your two candidates went down to one. You learned the hidden bit.

The price was zero. And the machine's own list of where it could be went from 00 and 10 to 00 and 11. Still two. Nothing it could have been got merged with anything else.

You narrowed. It didn't.

6.4 Two rows of dots

That last line is the thing I had to get straight before any of this made sense, so here it is as a picture.

Draw the machine's possible states as a row of dots. Now draw a second row above it: the dots that you, watching from outside, still think it might be on. The first row is a fact about the machine. The second is a fact about you, about what you've seen and what you've crossed off. Both rows can shrink, and they shrink for different reasons.

The machine's row shrinks only when two of its arrows meet. That's what the price charges. Run the halving rule from chapter three along a whole run and the counts just multiply: if the first move costs one, the row can at most halve; if the next costs two, it can shrink to a quarter of that; and so on. So a run can't shrink the machine's own row by more halvings than it paid for, all told. (Count with whole questions instead and you lose a little: four places and three places both take two yes-or-no questions to pin down, so the rounded count can't see that squeeze. The plain count can.)

Your row shrinks when you see something some of your candidates couldn't have shown you, and you cross them off. Nothing happened to the machine. It didn't merge anything. It doesn't know you're watching. It would have done the same thing with nobody there.

The physicist Edwin Jaynes had a name for mixing those two rows up, and it's one of the most useful names I know: the mind projection fallacy, treating what you know about a thing as if it were a property of the thing. Your list of candidates is in

your head. The machine's row is in the machine. The price is about the machine's row, and crossing a candidate off yours doesn't move it.

I have to be fair to Jaynes, though, because he'd argue with where I just took his phrase. In a 1965 paper he wrote that entropy is "an anthropomorphic concept," a phrase he credited to the physicist Eugene Wigner, meaning it belongs to the particular experiments you or I choose to do on a system, not to the system alone. For Jaynes, the second law is about what's known too, and the line I'd like to draw, that physics cares about the machine's row and not yours, is exactly the line he'd refuse. That fight, Jaynes against Boltzmann, is one of the oldest in the subject, and I can't settle it and don't have to. What I need is only this: the two rows are different rows, and only one of them belongs to the machine.

6.5 Who holds the pen

Here's the most beautiful thing in this chapter, and you find it by playing with the four dots, not by thinking about them.

Run MEASURE twice. The first time copies the hidden bit onto the display. The second time adds the hidden bit in again, and adding it twice the light-switch way gets you back where you started. Check the arrows: 10 goes to 11, and 11 goes back to 10. The display went full and then blank again, no two arrows met, and the price was zero both times.

So a display that was blank before the first MEASURE can be put back to blank with no arrows meeting, by a second MEASURE. Not by WIPE. This is the borrowed page being handed back: while the original is still there to check against, the copy can be un-copied.

What's the difference? MEASURE looks at the hidden bit while it clears the display. It has the thing the copy came from, so it can undo the copy. WIPE doesn't look. It sets the display to 0 regardless, and "regardless" is exactly where the arrows meet. Bennett has this in his 2003 paper too: he lists erasing "one of two copies of data known to be identical" as a step that loses nothing, because you're only removing a duplicate. And Lídia del Rio and her colleagues argued in 2011 that the cost of erasing a system depends on how much the eraser already knows about it, dropping to zero, and in the quantum case below zero. The cost of clearing a record falls on whoever clears it, and it's lower the more they know.

Try it with a card under a cup, red or blue, and a note that's blank or marked. Start with the note blank. Lift the cup: for red, flip the note, blank to marked or marked to blank; for blue, leave it. Now the note tells you the colour. Do the same thing again while the card is still there, and the note goes back to blank. Check all four card-and-note combinations and you'll see this rule undoes itself, so no two combinations land together. Now erase the note regardless of the card: for either

card, blank and marked both land on blank. Arrows meet. The cards don't measure anything about heat. What they show is which information each rule gets to use.

6.6 Learning after the first look

A fair objection: the demon's watcher learned something by looking, but any watcher learns something on the very first look, before the machine does anything. So the better question is, after that first look, can a step teach you more, at no cost?

Yes. The smallest machine that does it has three states in a ring: U goes to V, V goes to W, W goes back to U. The window shows "true" only on W. Start with two candidates, U or V. Both show "false," so the first look tells you nothing. Take one step. From U you'd now see "false"; from V you'd see "true." One step told them apart, and a ring never has two arrows meeting, so the price can be zero.

Two states can't do it, ever, with any arrows. If you can't tell two states apart at the first look, the window shows the same thing on both, and those two are all the states there are. So the window shows the same thing everywhere, forever. Try to draw a counterexample with two dots. You'll draw the same two dots a few times and give up, and giving up is the proof. Three is the smallest number of states where two that look alike at first can come apart later.

6.7 Finishing together

One more count before the big picture, because it's about how much room learning needs.

Run a machine from several different starts, and photograph its whole state at every tick. Lay the photographs out in rows, one row per start, one column per tick. Say every run finishes for the first time at the same tick. And say that in every column up to the one just before the finish, no two states step to the same state. That includes the last step, the one onto the finish: if two runs sat on two different states just before the finish, they can't both step onto one finished state.

Could two photographs on that sheet show the same state? Two in the same column: walk both back a tick at a time, and since no arrows met, they were equal the tick before, all the way back to the starts. But the starts were different. Two in different columns: give them both the same remaining steps. The one that's ahead would finish first, and it was supposed to finish at the same tick as everybody else. So every photograph shows a different state. With N starts and T ticks, the machine needs room for N times (T + 1) states. Running in step and finishing together has a size.

6.8 The certainty count

Now I can put the whole path from not knowing to knowing on one page.

You don't know which state the machine started in. There are a few it could have been, each with its chance. Run the machine. Take any two of those possible starts and watch where they go. Only two things can happen. They land on the same state, and the difference between them is gone from the machine: two arrows met, and under the halving rule from chapter three the price covered it, at most one halving per unit. Or they land on two different states, and the difference is still sitting in the machine, kept as two states held apart. There's no third thing a pair of starts can do.

So everything you didn't know about the start splits cleanly in two: the part the run destroyed, which was paid for, and the part the end state still holds.

Now put a watcher at the window. Whatever they see depends only on where the machine started, so whatever they learn is a fact about the start. And the same split holds for it: what they learned is either still held in the machine's end state, or it's gone from the machine and two arrows meeting paid for losing it. A run that costs nothing leaves everything the watcher learned still held in the machine.

That's the exact sense of "looking borrows." The watcher's copy is a copy, and the machine still holds the original. When the machine lets go of the original, two arrows meet, and the bill comes due.

What that doesn't say: how much heat. That's for the heat chapter. It doesn't say the watcher's own notes are free; they need a blank page too. And if the start is known for certain, there's nothing to learn and nothing destroyed, so there's no floor from any of this at all.

6.9 Who gets charged is partly bookkeeping

The physicists had this freedom on the books before me. Paul Fahn in 1996, doing the accounting of Szilard's one-molecule engine step by step, argued that whether you charge the looking step or the wiping step is partly a choice of how you keep the books, and that the total is what's fixed. Takahiro Sagawa and Masahito Ueda in 2009 worked out a bound on that total, with the split depending on the memory. So "a price list could charge MEASURE instead" is a known freedom, not a loophole I found.

The other side is alive too. Ruth Kastner argued in 2025 that Brillouin was essentially right after all: put the real physics of measuring back in, and it's the measuring that pays and stops the demon, and Bennett's free copy fails as physics. And John Norton argued in 2013 that at the scale of molecules, jiggling from heat keeps any process that's supposed to be perfectly reversible from ever finishing, and

finishing it costs more than Landauer's amount. If either of them is right, a real demon's look has a cost after all.

What's on the page here survives that, because it's about price lists: a price list that follows Bennett's rule isn't forced to charge the copy, whatever a real device spends. What wouldn't survive is the physics I've been hanging on it, that a real demon can look for free. If Norton is right, the line between looking and keeping keeps its counting and loses its physics. I can't referee that one.

And in 2010 a group in Tokyo, Shoichi Toyabe, Takahiro Sagawa, Masahito Ueda, Eiro Muneyuki and Masaki Sano, ran a real demon: a tiny particle walked up a staircase by feedback on what a camera saw. The energy it gained was paid for by the information the camera took, and nothing was violated. The demon is real, it works, and it pays. Where it pays is the part that took a century.

6.10 Somebody got there first, goddamn it

I'd been sitting on this idea, that making a result stay so it can't be handed back is what pays, and finding out only borrows, and I thought: this one's mine. Then I found Carlo Rovelli's 2024 paper, "The Thermodynamic Cost of Choosing." A real physicist, with a real paper, saying that making a choice that sticks gives off at least the heat Landauer charges for erasing a bit. He even quotes a German saying, Wahl ist Leiden, to choose is to suffer, and at that point what are you going to do but laugh.

On its face he contradicts Bennett. Bennett says measuring doesn't have to cost; Rovelli says any measurement costs. So look at what Rovelli calls a choice. A coin in the air hasn't chosen anything; it has to hit the ground and stop. And then the sentence I'd put on my wall: "stabilization must be irreversible, otherwise it is not stabilization." Bennett's free look is one that can still be undone, which is why it's free. Rovelli's choice is one that's been made to stay, which is why it costs. The way I read them, they're standing on opposite sides of the same line: looking, which borrows, on Bennett's side, and keeping, which pays, on Rovelli's.

That reading is mine, and I owe you exactly how much of it. Push on it with his paper open and it gives in four places. He saves the word "choice" for an outcome that's stable and not predictable, and rules out a thermostat because it runs on a predictable mechanism; the settling move in this book is fixed in advance by a rule, so by his own definition his result doesn't cover it. His choice is one possibility branching into two, which he calls in a sense the opposite of erasing; the toll comes from two arrows meeting, the opposite shape. His stability lasts a while, not forever; mine is forever, by definition. And he doesn't stand beside Bennett at all: he holds that any measurement costs, credits that to the philosopher Hans Reichenbach, and doesn't cite Bennett. So what Rovelli and I share is a sentence: a choice that sticks

costs something. Push on it and his coin hitting the ground, a fact nobody could predict, sits closer to looking than to keeping. I still think it's the same instinct. You've got everything you need to disagree.

And on copying, the nearest physics isn't Rovelli's. It's in biology. Thomas Ouldridge and Pieter Rein ten Wolde worked out in 2017 the basic costs of making a copy of a molecule that stays matched to its template after the two separate, and of destroying one. A companion paper the same year, with Christopher Govern, does the same kind of accounting for how a cell reads out its own sensors, which is a measurement in all but name. The way I read them, the copy that has to stay is the one that pays. Then Jenny Poulton, ten Wolde and Ouldridge in 2019 found the sharp form: in their simple models, when a copy grows with the weakest possible push, its advantage over random copying, and the efficiency of storing that information, both vanish. Pushing harder, with chemical work, makes copies that stay matched after the two let go. That's finding out against keeping, in chemistry, with the cost attached, and biology had the distinction before I did. Theirs is a correlation between molecules, with heat in it. Mine is a reading in a model of computation, with no temperature anywhere.

Two more neighbours have "learning" and "commit" in their titles, so the line here shouldn't be read against them by accident. Sebastian Goldt and Udo Seifert in 2017 did the physics of a neural network learning a rule, and argued that learning has a cost: the entropy the network produces limits how much it can learn. That doesn't contradict this chapter. Their learner isn't copying onto a blank page; it's a driven, noisy physical system, and that's what they price. This chapter only says one way of finding out, a copy onto a blank, is allowed to be free. Nothing here says every way of learning is cheap, and theirs isn't. And from statistics, Aurélien Garivier, Tor Lattimore and Emilie Kaufmann in 2016 compared strategies that explore for a while and then commit to one choice against strategies that keep adjusting as they go. In their setting, over a long run, the best strategies that commit lose about twice as much reward as the best ones that keep adjusting. That's the closest published sentence I've found to "committing costs something" with a number on it. It's lost reward, though, not a count on a ledger, so it's a cousin, not a twin.

6.11 What the toll is for, said at last

Now I can say exactly what the toll charges. Not looking. Not copying. Not finding out. A look can be done with no arrows meeting, on a price list that charges nothing for it; the demon is a machine where that happens. Looking uses up a blank page, and while the original is still there, the page can be handed back. What gets charged is the one thing chapter two showed can't be done on a finite machine without two arrows meeting: settling something so it stays, a record nothing hands back.

That's what "No Free Insight," the name I gave the toll before I understood it, really covers, and I don't want the name doing work it hasn't earned. Insight in the sense of finding out can come free; it borrows a page. The insight the toll prices is the kind made to stay. Whether that narrower thing deserves the name is fair to argue about. I kept the name because it's the question I started with, and because narrowing it down is the finding.

One more thing this tells you, for later. A later chapter counts what it takes to narrow a list of candidates down to a shorter one, and puts a cost on it. Narrowing alone doesn't force anybody to pay anything; the demon just narrowed two candidates to one for free. So wherever this book puts a price on narrowing, the payment goes in as an assumption, out in the open, and I'll keep calling it one.