

## Advanced Statistical Inference

### Gaussian Processes

Maurizio Filippone  
Maurizio.Filippone@eurecom.fr

Department of Data Science  
EURECOM

## Gaussian Processes

- ▶ Linear models requires specifying a set of basis functions
  - ▶ Polynomials, Trigonometric, ...??
- ▶ Can we use Bayesian inference to let data tell us?
- ▶ Gaussian Processes work implicitly with an infinite set of basis functions and learn a probabilistic combination of these

## Suggested readings

Gaussian Processes for Machine Learning

Carl E. Rasmussen and Christopher K. I. Williams

Pattern Recognition and Machine Learning

C. Bishop

## Gaussian Processes

Gaussian Processes can be explained in two ways

- ▶ Weight Space View
  - ▶ Bayesian linear regression with infinite basis functions
- ▶ Function Space View
  - ▶ Defined as priors over functions

## Gaussian Processes

Gaussian Processes can be explained in two ways

- ▶ **Weight Space View**
  - ▶ **Bayesian linear regression with infinite basis functions**
- ▶ Function Space View
  - ▶ Defined as priors over functions

## Bayesian Linear Regression - recap

- ▶ Posterior **must be** Gaussian

$$p(\mathbf{w}|\mathbf{X}, \mathbf{y}, \sigma^2) = \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

- ▶ Covariance:

$$\boldsymbol{\Sigma} = \left( \frac{1}{\sigma^2} \mathbf{X}^\top \mathbf{X} + \mathbf{S}^{-1} \right)^{-1}$$

- ▶ Mean:

$$\boldsymbol{\mu} = \frac{1}{\sigma^2} \boldsymbol{\Sigma} \mathbf{X}^\top \mathbf{y}$$

- ▶ Predictions

$$p(y_*|\mathbf{X}, \mathbf{y}, \mathbf{x}_*, \sigma^2) = \mathcal{N}(\mathbf{x}_*^\top \boldsymbol{\mu}, \sigma^2 + \mathbf{x}_*^\top \boldsymbol{\Sigma} \mathbf{x}_*)$$

## Bayesian Linear Regression - recap

- ▶ Modeling observations as noisy realizations of a linear combination of the features:

$$p(\mathbf{y}|\mathbf{w}, \mathbf{X}, \sigma^2) = \mathcal{N}(\mathbf{X}\mathbf{w}, \sigma^2 \mathbf{I})$$

- ▶ Gaussian prior over model parameters:

$$p(\mathbf{w}) = \mathcal{N}(\mathbf{0}, \mathbf{S})$$

## Introducing basis functions

- ▶ Imagine transforming the inputs using a set of  $D$  functions

$$\mathbf{x} \rightarrow \phi(\mathbf{x}) = (\phi_1(\mathbf{x}), \dots, \phi_D(\mathbf{x}))^\top$$

- ▶ The functions  $\phi_1(\mathbf{x})$  are also known as basis functions
- ▶ Define:

$$\boldsymbol{\Phi} = \begin{bmatrix} \phi_1(\mathbf{x}_1) & \dots & \phi_D(\mathbf{x}_1) \\ \vdots & \ddots & \vdots \\ \phi_1(\mathbf{x}_N) & \dots & \phi_D(\mathbf{x}_N) \end{bmatrix}$$

## Introducing basis functions

- ▶ Applying Bayesian Linear Regression on the transformed features gives

$$p(\mathbf{w}|\mathbf{X}, \mathbf{y}, \sigma^2) = \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

- ▶ Covariance:

$$\boldsymbol{\Sigma} = \left( \frac{1}{\sigma^2} \boldsymbol{\Phi}^\top \boldsymbol{\Phi} + \mathbf{S}^{-1} \right)^{-1}$$

- ▶ Mean:

$$\boldsymbol{\mu} = \frac{1}{\sigma^2} \boldsymbol{\Sigma} \boldsymbol{\Phi}^\top \mathbf{y}$$

- ▶ Predictions:

$$p(y_*|\mathbf{X}, \mathbf{y}, \mathbf{x}_*, \sigma^2) = \mathcal{N}(\boldsymbol{\phi}_*^\top \boldsymbol{\mu}, \sigma^2 + \boldsymbol{\phi}_*^\top \boldsymbol{\Sigma} \boldsymbol{\phi}_*)$$

## Bayesian Linear Regression as a Kernel Machine

- ▶ Working with  $\psi(\cdot)$  costs  $O(D^2)$  storage,  $O(D^3)$  time
- ▶ Working with  $k(\cdot, \cdot)$  costs  $O(N^2)$  storage,  $O(N^3)$  time
- ▶ Pick the one that makes computations faster ... or
- ▶ What if we could pick  $k(\cdot, \cdot)$  so that  $\psi(\cdot)$  is infinite dimensional?

## Bayesian Linear Regression as a Kernel Machine

- ▶ We are going to show that predictions can be expressed exclusively in terms of scalar products as follows

$$k(\mathbf{x}, \mathbf{x}') = \boldsymbol{\psi}(\mathbf{x})^\top \boldsymbol{\psi}(\mathbf{x}')$$

- ▶ This allows us to work with either  $k(\cdot, \cdot)$  or  $\boldsymbol{\psi}(\cdot)$
- ▶ Why is this useful??

## Kernels

- ▶ It is possible to show that for

$$k(\mathbf{x}, \mathbf{x}') = \exp\left(-\frac{\|\mathbf{x} - \mathbf{x}'\|^2}{2}\right)$$

there exists a corresponding  $\boldsymbol{\psi}(\cdot)$  that is infinite dimensional!!!

- ▶ There are other kernels satisfying this property

## Kernels

Proof that the Gaussian kernel induces an infinite dimensional  $\psi(\cdot)$

- ▶ For simplicity consider one dimensional inputs  $x, z$
- ▶ Expand the Gaussian kernel  $k(x, z)$  as

$$\exp\left(-\frac{(x-z)^2}{2}\right) = \exp\left(-\frac{x^2}{2}\right) \exp\left(-\frac{z^2}{2}\right) \exp(xz)$$

- ▶ Focusing on the last term and applying the Taylor expansion of the  $\exp(\cdot)$  function

$$\exp(xz) = 1 + (xz) + \frac{(xz)^2}{2!} + \frac{(xz)^3}{3!} + \frac{(xz)^4}{4!} + \dots$$

## Bayesian Linear Regression as a Kernel Machine

Proof

- ▶ To show that Bayesian Linear Regression can be formulated through scalar products only, we need Woodbury identity:

$$(A + UCV)^{-1} = A^{-1} - A^{-1}U(C^{-1} + VA^{-1}U)^{-1}VA^{-1}$$

- ▶ Do not memorize this!

## Kernels

Proof that the Gaussian kernel induces an infinite dimensional  $\psi(\cdot)$

- ▶ Define the infinite dimensional mapping

$$\psi(x) = \exp\left(-\frac{x^2}{2}\right) \left(1, x, \frac{x^2}{\sqrt{2!}}, \frac{x^3}{\sqrt{3!}}, \frac{x^4}{\sqrt{4!}}, \dots\right)^\top$$

- ▶ It is easy to verify that

$$k(x, z) = \exp\left(-\frac{(x-z)^2}{2}\right) = \psi(x)^\top \psi(z)$$

## Bayesian Linear Regression as a Kernel Machine

Proof

- ▶ Woodbury identity:

$$(A + UCV)^{-1} = A^{-1} - A^{-1}U(C^{-1} + VA^{-1}U)^{-1}VA^{-1}$$

- ▶ We can rewrite:

$$\begin{aligned} \Sigma &= \left(\frac{1}{\sigma^2} \Phi^\top \Phi + \mathbf{S}^{-1}\right)^{-1} \\ &= \mathbf{S} - \mathbf{S} \Phi^\top \left(\sigma^2 \mathbf{I} + \Phi \mathbf{S} \Phi^\top\right)^{-1} \Phi \mathbf{S} \end{aligned}$$

- ▶ We set  $A = \mathbf{S}$ ,  $U = V^\top = \Phi^\top$ , and  $C = \frac{1}{\sigma^2} \mathbf{I}$

## Bayesian Linear Regression as a Kernel Machine

Proof

- Mean and variance of the predictions:

$$p(y_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_*, \sigma^2) = \mathcal{N}(\phi_*^\top \boldsymbol{\mu}, \sigma^2 + \phi_*^\top \boldsymbol{\Sigma} \phi_*)$$

- Rewrite the variance:

$$\sigma^2 + \phi_*^\top \boldsymbol{\Sigma} \phi_* =$$

$$\sigma^2 + \phi_*^\top \mathbf{S} \phi_* - \phi_*^\top \mathbf{S} \boldsymbol{\Phi}^\top (\sigma^2 \mathbf{I} + \boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top)^{-1} \boldsymbol{\Phi} \mathbf{S} \phi_*$$

... continued

## Bayesian Linear Regression as a Kernel Machine

Proof

- Mean and variance of the predictions:

$$p(y_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_*, \sigma^2) = \mathcal{N}(\phi_*^\top \boldsymbol{\mu}, \sigma^2 + \phi_*^\top \boldsymbol{\Sigma} \phi_*)$$

- Rewrite the mean:

$$\begin{aligned} \phi_*^\top \boldsymbol{\mu} &= \frac{1}{\sigma^2} \phi_*^\top \boldsymbol{\Sigma} \boldsymbol{\Phi}^\top \mathbf{y} \\ &= \frac{1}{\sigma^2} \phi_*^\top \left( \mathbf{S} - \mathbf{S} \boldsymbol{\Phi}^\top (\sigma^2 \mathbf{I} + \boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top)^{-1} \boldsymbol{\Phi} \mathbf{S} \right) \boldsymbol{\Phi}^\top \mathbf{y} \\ &= \frac{1}{\sigma^2} \phi_*^\top \mathbf{S} \boldsymbol{\Phi}^\top \left( \mathbf{I} - (\sigma^2 \mathbf{I} + \boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top)^{-1} \boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top \right) \mathbf{y} \\ &= \frac{1}{\sigma^2} \phi_*^\top \mathbf{S} \boldsymbol{\Phi}^\top \left( \mathbf{I} - \left( \mathbf{I} + \frac{\boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top}{\sigma^2} \right)^{-1} \frac{\boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top}{\sigma^2} \right) \mathbf{y} \end{aligned}$$

... continued

## Bayesian Linear Regression as a Kernel Machine

Proof

- Mean and variance of the predictions:

$$p(y_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_*, \sigma^2) = \mathcal{N}(\phi_*^\top \boldsymbol{\mu}, \sigma^2 + \phi_*^\top \boldsymbol{\Sigma} \phi_*)$$

- Rewrite the variance:

$$\sigma^2 + \phi_*^\top \mathbf{S} \phi_* - \phi_*^\top \mathbf{S} \boldsymbol{\Phi}^\top (\sigma^2 \mathbf{I} + \boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top)^{-1} \boldsymbol{\Phi} \mathbf{S} \phi_* =$$

$$\sigma^2 + k_{**} - \mathbf{k}_*^\top (\sigma^2 \mathbf{I} + \mathbf{K})^{-1} \mathbf{k}_*$$

- Where the mapping defining the kernel is

$$\boldsymbol{\psi}(\mathbf{x}) = \mathbf{S}^{1/2} \boldsymbol{\phi}(\mathbf{x}) \quad \text{and}$$

$$k_{**} = k(\mathbf{x}_*, \mathbf{x}_*) = \boldsymbol{\psi}(\mathbf{x}_*)^\top \boldsymbol{\psi}(\mathbf{x}_*)$$

$$(\mathbf{k}_*)_i = k(\mathbf{x}_*, \mathbf{x}_i) = \boldsymbol{\psi}(\mathbf{x}_*)^\top \boldsymbol{\psi}(\mathbf{x}_i)$$

$$(\mathbf{K})_{ij} = k(\mathbf{x}_i, \mathbf{x}_j) = \boldsymbol{\psi}(\mathbf{x}_i)^\top \boldsymbol{\psi}(\mathbf{x}_j)$$

## Bayesian Linear Regression as a Kernel Machine

Proof

- Define  $\mathbf{H} = \frac{\boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top}{\sigma^2}$

- The term in the parenthesis

$$\left( \mathbf{I} - \left( \mathbf{I} + \frac{\boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top}{\sigma^2} \right)^{-1} \frac{\boldsymbol{\Phi} \mathbf{S} \boldsymbol{\Phi}^\top}{\sigma^2} \right)$$

becomes

$$(\mathbf{I} - (\mathbf{I} + \mathbf{H})^{-1} \mathbf{H}) = \mathbf{I} - (\mathbf{H}^{-1} + \mathbf{I})^{-1}$$

- Using Woodbury ( $A, U, V = \mathbf{I}$  and  $C = \mathbf{H}^{-1}$ )

$$\mathbf{I} - (\mathbf{H}^{-1} + \mathbf{I})^{-1} = (\mathbf{I} + \mathbf{H})^{-1}$$

## Bayesian Linear Regression as a Kernel Machine

Proof



- ▶ Substituting into the expression of the predictive mean

$$\begin{aligned}\phi_*^\top \mu &= \frac{1}{\sigma^2} \phi_*^\top \mathbf{S} \Phi^\top \left( \mathbf{I} - \left( \mathbf{I} + \frac{\Phi \mathbf{S} \Phi^\top}{\sigma^2} \right)^{-1} \frac{\Phi \mathbf{S} \Phi^\top}{\sigma^2} \right) \mathbf{y} \\ &= \frac{1}{\sigma^2} \phi_*^\top \mathbf{S} \Phi^\top \left( \mathbf{I} + \frac{\Phi \mathbf{S} \Phi^\top}{\sigma^2} \right)^{-1} \mathbf{y} \\ &= \phi_*^\top \mathbf{S} \Phi^\top \left( \sigma^2 \mathbf{I} + \Phi \mathbf{S} \Phi^\top \right)^{-1} \mathbf{y} \\ &= \mathbf{k}_*^\top \left( \sigma^2 \mathbf{I} + \mathbf{K} \right)^{-1} \mathbf{y}\end{aligned}$$

- ▶ All definitions as in the case of the variance

## Gaussian Processes - Prior over Functions

- ▶ Consider an infinite number of Gaussian random variables
- ▶ Think of them as indexed by the real line and as independent
- ▶ Denote them as  $f(\mathbf{x})$



## Gaussian Processes

Gaussian Processes can be explained in two ways

- ▶ Weight Space View
  - ▶ Bayesian linear regression with infinite basis functions
- ▶ **Function Space View**
  - ▶ Defined as priors over functions

## Kernel



- ▶ Consider the Gaussian kernel again

$$k(\mathbf{x}, \mathbf{x}') = \alpha \exp(-\beta \|\mathbf{x} - \mathbf{x}'\|^2)$$

- ▶ We introduced some parameters for added flexibility



## Gaussian Processes - Prior over Functions

- Impose covariance using the kernel function





## Gaussian Processes - Prior over Functions

- Draw the infinite random variables again fixing one of them (the one at  $x = 0$ )





## Gaussian Processes - Prior over Functions

- Draw the infinite random variables again allowing the one at  $x = 0$  to be random too




## Gaussian Processes - Prior over Functions

- This can be used as a prior over functions!




## Gaussian Processes - Priors over Functions

- Infinite Gaussian random variables with parameterized and input-dependent covariance



## Gaussian Processes - Prior over Functions


- The distribution of  $N$  random variables  $f(\mathbf{x}_1), \dots, f(\mathbf{x}_N)$  depends exclusively on the corresponding rows and columns of the infinite by infinite kernel matrix  $K$



$$K = \begin{matrix} & \underbrace{\hspace{1cm}}_n \\ \begin{matrix} \uparrow \\ n \end{matrix} & \begin{matrix} \text{[Grid of } n \times n \text{ blocks]} \end{matrix} \end{matrix}$$

## Gaussian Processes - Prior over Functions

- The distribution of  $N$  random variables  $f(\mathbf{x}_1), \dots, f(\mathbf{x}_N)$  depends exclusively on the corresponding rows and columns of the infinite by infinite kernel matrix  $K$



$$K = \begin{matrix} \begin{matrix} \uparrow \\ n \end{matrix} & \begin{matrix} \text{[Grid of } n \times n \text{ blocks]} \end{matrix} \end{matrix}$$

## Gaussian Processes - Prior over Functions

- The marginal distribution of  $\mathbf{f} = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_N))^T$  is

$$p(\mathbf{f}|\mathbf{X}) = \mathcal{N}(\mathbf{0}, \mathbf{K})$$

- The conditional distribution of  $f_*$  given  $\mathbf{f}$

$$p(f_*|\mathbf{f}, \mathbf{x}_*, \mathbf{X}) = \mathcal{N}(\bar{m}, \bar{s}^2)$$

with

$$\bar{m} = \mathbf{k}_*^T \mathbf{K}^{-1} \mathbf{f}$$

$$\bar{s}^2 = k_{**} - \mathbf{k}_*^T \mathbf{K}^{-1} \mathbf{k}_*$$



## Gaussian Processes - Prior over Functions

- ▶ Remember that when we modeled labels  $\mathbf{y}$  in the linear model we assumed noise with variance  $\sigma$  around  $\mathbf{w}^\top \mathbf{x}$
- ▶ We can do the same in Gaussian processes

$$p(\mathbf{y}|\mathbf{f}) = \prod_{i=1}^N p(y_i|f_i)$$

with

$$p(y_i|f_i) = \mathcal{N}(y_i|f_i, \sigma^2)$$

- ▶ Likelihood and prior are both Gaussian - conjugate!
- ▶ We can integrate out the Gaussian process prior over  $\mathbf{f}$

$$p(\mathbf{y}|\mathbf{X}) = \int p(\mathbf{y}|\mathbf{f})p(\mathbf{f}|\mathbf{X})d\mathbf{f}$$

- ▶ This gives

$$p(\mathbf{y}|\mathbf{X}) = \mathcal{N}(\mathbf{0}, \mathbf{K} + \sigma^2\mathbf{I})$$

## Gaussian Processes - Prior over Functions

- ▶ We can also make predictions as follows:

$$\begin{aligned} p(y_*|\mathbf{y}, \mathbf{x}_*\mathbf{X}) &= \int p(y_*|f_*)p(f_*|\mathbf{f}, \mathbf{x}_*, \mathbf{X})p(\mathbf{f}|\mathbf{y}, \mathbf{X})d\mathbf{f}df_* \\ &= \mathcal{N}(m_y, s_y^2) \end{aligned}$$

with

$$m_y = \mathbf{k}_*^\top (\mathbf{K} + \sigma^2\mathbf{I})^{-1} \mathbf{y}$$

$$s_y^2 = \sigma^2 + k_{**} - \mathbf{k}_*^\top (\mathbf{K} + \sigma^2\mathbf{I})^{-1} \mathbf{k}_*$$

- ▶ Same expression as in the “Weight-Space View” section

## Gaussian Processes - Prior over Functions

- ▶ We can derive the predictive distribution as follows:

$$p(f_*|\mathbf{y}, \mathbf{x}_*\mathbf{X}) = \int p(f_*|\mathbf{f}, \mathbf{x}_*, \mathbf{X})p(\mathbf{f}|\mathbf{y}, \mathbf{X})d\mathbf{f}df_* = \mathcal{N}(m, s^2)$$

with


$$m = \mathbf{k}_*^\top (\mathbf{K} + \sigma^2\mathbf{I})^{-1} \mathbf{y}$$

$$s^2 = k_{**} - \mathbf{k}_*^\top (\mathbf{K} + \sigma^2\mathbf{I})^{-1} \mathbf{k}_*$$

- ▶ Same expression as in the “Weight-Space View” section


## Gaussian Processes - Regression example

- ▶ Some data generated as a noisy version of some function



## Gaussian Processes - Regression example

- Draws from the posterior distribution over  $f_*$  on the real line



## Optimization of Gaussian Process parameters

- Define  $\mathbf{C} = \mathbf{K} + \sigma^2 \mathbf{I}$
- Maximize the logarithm of the likelihood

$$p(\mathbf{y}|\mathbf{X}, \theta) = \mathcal{N}(\mathbf{0}, \mathbf{C})$$

that is

$$-\frac{1}{2} \log |\mathbf{C}| - \frac{1}{2} \mathbf{y}^\top \mathbf{C}^{-1} \mathbf{y} + \text{const.}$$

- Derivatives can be useful for gradient-based optimization

$$\frac{\partial \log[p(\mathbf{y}|\mathbf{X}, \theta)]}{\partial \theta_i}$$

## Optimization of Gaussian Process parameters

- The kernel has parameters that have to be tuned

$$k(\mathbf{x}, \mathbf{x}') = \alpha \exp(-\beta \|\mathbf{x} - \mathbf{x}'\|^2)$$

... and there is also the noise parameter  $\sigma^2$ .

- Define  $\theta = (\alpha, \beta, \sigma^2)$
- How should we tune them?

## Optimization of Gaussian Process parameters

- Log-likelihood

$$-\frac{1}{2} \log |\mathbf{C}| - \frac{1}{2} \mathbf{y}^\top \mathbf{C}^{-1} \mathbf{y} + \text{const.}$$

- Derivatives can be useful for gradient-based optimization:

$$\frac{\partial \log[p(\mathbf{y}|\mathbf{X}, \theta)]}{\partial \theta_i} = -\frac{1}{2} \text{Tr} \left( \mathbf{C}^{-1} \frac{\partial \mathbf{C}}{\partial \theta_i} \right) + \frac{1}{2} \mathbf{y}^\top \mathbf{C}^{-1} \frac{\partial \mathbf{C}}{\partial \theta_i} \mathbf{C}^{-1} \mathbf{y}$$

## Summary

- ▶ Introduced Gaussian Processes
  - ▶ Weight space view
  - ▶ Function space view
- ▶ Gaussian processes for regression
- ▶ Optimization of kernel parameters
- ▶ To think about:
  - ▶ Gaussian processes for classification?
  - ▶ Scalability?