

Advanced Statistical Inference

Refresher on linear algebra

Maurizio Filippone
`Maurizio.Filippone@eurecom.fr`

Department of Data Science
EURECOM

Overview

Basics

Spectral decomposition

Positive definite matrices

Square root matrices

Basic definitions and properties

- Vectors $\mathbf{v} = [v_i]$

$$\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$$

Basic definitions and properties

- Vectors $\mathbf{v} = [v_i]$

$$\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$$

- Matrices $\mathbf{A} = [a_{ij}]$

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

Basic definitions and properties

- ▶ Matrix addition $\mathbf{A} + \mathbf{B} = [a_{ij} + b_{ij}]$

Basic definitions and properties

- ▶ Matrix addition $\mathbf{A} + \mathbf{B} = [a_{ij} + b_{ij}]$
- ▶ Scalar multiplication $\gamma \mathbf{B} = [\gamma b_{ij}]$

Basic definitions and properties

- ▶ Matrix addition $\mathbf{A} + \mathbf{B} = [a_{ij} + b_{ij}]$
- ▶ Scalar multiplication $\gamma \mathbf{B} = [\gamma b_{ij}]$
- ▶ Matrix-vector multiplication $\mathbf{A}\mathbf{v} = \left[\sum_k a_{ik} v_k \right]$

Basic definitions and properties

- ▶ Matrix addition $\mathbf{A} + \mathbf{B} = [a_{ij} + b_{ij}]$
- ▶ Scalar multiplication $\gamma \mathbf{B} = [\gamma b_{ij}]$
- ▶ Matrix-vector multiplication $\mathbf{A}\mathbf{v} = \left[\sum_k a_{ik} v_k \right]$
- ▶ Matrix-matrix multiplication $\mathbf{A}\mathbf{B} = \left[\sum_k a_{ik} b_{kj} \right]$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{AA}^{-1} = \mathbf{I}$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^{\top} = [a_{ji}]$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^{\top} = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{A}\mathbf{B})^{\top} = \mathbf{B}^{\top}\mathbf{A}^{\top}$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^{\top} = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{A}\mathbf{B})^{\top} = \mathbf{B}^{\top}\mathbf{A}^{\top}$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^{\top}$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^\top = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{A}\mathbf{B})^\top = \mathbf{B}^\top \mathbf{A}^\top$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^\top$
- ▶ Determinant $|\mathbf{A}|$ (on the board)

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^{\top} = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{AB})^{\top} = \mathbf{B}^{\top}\mathbf{A}^{\top}$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^{\top}$
- ▶ Determinant $|\mathbf{A}|$ (on the board)
- ▶ Determinant $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^\top = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{A}\mathbf{B})^\top = \mathbf{B}^\top \mathbf{A}^\top$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^\top$
- ▶ Determinant $|\mathbf{A}|$ (on the board)
- ▶ Determinant $|\mathbf{A}\mathbf{B}| = |\mathbf{A}||\mathbf{B}|$
- ▶ Determinant $|\mathbf{A}^\top| = |\mathbf{A}|$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^\top = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{A}\mathbf{B})^\top = \mathbf{B}^\top \mathbf{A}^\top$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^\top$
- ▶ Determinant $|\mathbf{A}|$ (on the board)
- ▶ Determinant $|\mathbf{A}\mathbf{B}| = |\mathbf{A}||\mathbf{B}|$
- ▶ Determinant $|\mathbf{A}^\top| = |\mathbf{A}|$
- ▶ Trace $\text{Tr}(\mathbf{A}) = \sum_k a_{kk}$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^\top = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^\top$
- ▶ Determinant $|\mathbf{A}|$ (on the board)
- ▶ Determinant $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$
- ▶ Determinant $|\mathbf{A}^\top| = |\mathbf{A}|$
- ▶ Trace $\text{Tr}(\mathbf{A}) = \sum_k a_{kk}$
- ▶ Trace is permutation invariant
 $\text{Tr}(\mathbf{ABC}) = \text{Tr}(\mathbf{CAB}) = \text{Tr}(\mathbf{BCA})$

Eigenvalues and eigenvectors

- ▶ Let $\mathbf{A} = [a_{i,j}]$ be an $n \times n$ matrix
- ▶ \mathbf{A} is said to have an eigenvalue λ and (non-zero) eigenvector \mathbf{x} corresponding to λ if

$$\mathbf{Ax} = \lambda\mathbf{x}.$$

- ▶ Eigenvalues are the λ values that solve the determinantal equation $|\mathbf{A} - \lambda\mathbf{I}| = 0$.

Spectral decomposition of symmetric matrices

- ▶ The Spectral decomposition theorem says that every square and symmetric matrix $\mathbf{A} = [a_{ij}]$ may be written

$$\mathbf{A} = \mathbf{C}\mathbf{D}\mathbf{C}^\top,$$

- ▶ The columns of \mathbf{C} are the eigenvectors of \mathbf{A}
- ▶ The diagonal matrix \mathbf{D} contains the corresponding eigenvalues

$$\mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}$$

- ▶ The eigenvectors may be chosen to be orthonormal, so that $\mathbf{C}\mathbf{C}^\top = \mathbf{C}^\top\mathbf{C} = \mathbf{I}$.
- ▶ Note the useful property: $|\mathbf{A}| = \prod_{i=1}^n \lambda_i$

Positive definite matrices

The $n \times n$ matrix \mathbf{A} is said to be positive definite if

$$\mathbf{y}^\top \mathbf{A} \mathbf{y} > 0$$

for all $n \times 1$ vectors $\mathbf{y} \neq \mathbf{0}$.

Positive semidefinite matrices

The $n \times n$ matrix \mathbf{A} is said to be positive semidefinite if

$$\mathbf{y}^\top \mathbf{A} \mathbf{y} \geq 0$$

for all $n \times 1$ vectors $\mathbf{y} \neq \mathbf{0}$.

Example: Show $\mathbf{X}^\top \mathbf{X}$ is positive semidefinite

Let \mathbf{X} be an $n \times p$ matrix of real constants and \mathbf{y} be $p \times 1$. Then $\mathbf{Z} = \mathbf{X}\mathbf{y}$ is $n \times 1$, and

$$\begin{aligned} & \mathbf{y}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{y} \\ = & (\mathbf{X}\mathbf{y})^\top (\mathbf{X}\mathbf{y}) \\ = & \mathbf{Z}^\top \mathbf{Z} \\ = & \sum_{i=1}^n z_i^2 \geq 0 \end{aligned}$$

Some properties of symmetric positive definite matrices

For a symmetric matrix,

Positive definite



All eigenvalues positive

Showing Positive definite \Rightarrow Eigenvalues positive

Let \mathbf{A} be symmetric and positive definite.

- ▶ Spectral decomposition says $\mathbf{A} = \mathbf{C}\mathbf{D}\mathbf{C}^\top$.
- ▶ Using $\mathbf{y}^\top \mathbf{A} \mathbf{y} > 0$, let \mathbf{y} be an eigenvector, say the third one.
- ▶ Because eigenvectors are orthonormal,

$$\begin{aligned}\mathbf{y}^\top \mathbf{A} \mathbf{y} &= \mathbf{y}^\top \mathbf{C} \mathbf{D} \mathbf{C}^\top \mathbf{y} \\ &= \begin{pmatrix} 0 & 0 & 1 & \cdots & 0 \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} \\ &= \lambda_3 \\ &> 0\end{aligned}$$

Square root matrices

For symmetric, non-negative definite matrices

Define

$$\mathbf{D}^{1/2} = \begin{pmatrix} \sqrt{\lambda_1} & 0 & \cdots & 0 \\ 0 & \sqrt{\lambda_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sqrt{\lambda_n} \end{pmatrix}$$

So that

$$\begin{aligned} \mathbf{D}^{1/2} \mathbf{D}^{1/2} &= \begin{pmatrix} \sqrt{\lambda_1} & 0 & \cdots & 0 \\ 0 & \sqrt{\lambda_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sqrt{\lambda_n} \end{pmatrix} \begin{pmatrix} \sqrt{\lambda_1} & 0 & \cdots & 0 \\ 0 & \sqrt{\lambda_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sqrt{\lambda_n} \end{pmatrix} \\ &= \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = \mathbf{D} \end{aligned}$$

For a non-negative definite, symmetric matrix **A**

Define

$$\mathbf{A}^{1/2} = \mathbf{C}\mathbf{D}^{1/2}\mathbf{C}^\top$$

So that

$$\begin{aligned}\mathbf{A}^{1/2}\mathbf{A}^{1/2} &= \mathbf{C}\mathbf{D}^{1/2}\mathbf{C}^\top\mathbf{C}\mathbf{D}^{1/2}\mathbf{C}^\top \\ &= \mathbf{C}\mathbf{D}^{1/2}\mathbf{I}\mathbf{D}^{1/2}\mathbf{C}^\top \\ &= \mathbf{C}\mathbf{D}^{1/2}\mathbf{D}^{1/2}\mathbf{C}^\top \\ &= \mathbf{C}\mathbf{D}\mathbf{C}^\top \\ &= \mathbf{A}\end{aligned}$$

Cholesky decomposition

- ▶ Define lower triangular matrix \mathbf{L}

$$\mathbf{L} = \begin{bmatrix} L_{11} & 0 & \cdots & 0 \\ L_{21} & L_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{bmatrix}$$

so that $\mathbf{A} = \mathbf{L}\mathbf{L}^\top$

- ▶ Cholesky algorithm computes \mathbf{L} from \mathbf{A}

- ▶ $|\mathbf{A}| = |\mathbf{L}\mathbf{L}^\top| = |\mathbf{L}|^2 = \left(\prod_{i=1}^n L_{ii} \right)^2$