

Advanced Statistical Inference

Refresher on linear algebra

Motonobu Kanagawa
`motonobu.kanagawa@eurecom.fr`

Data Science Department
EURECOM

March 6, 2024

Overview

Basics

Spectral decomposition

Positive definite matrices

Square root matrices

Outline

Basics

Spectral decomposition

Positive definite matrices

Square root matrices

Basic definitions and properties

- ▶ Vectors $\mathbf{v} = [v_i]$

$$\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$$

Basic definitions and properties

- Vectors $\mathbf{v} = [v_i]$

$$\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$$

- Matrices $\mathbf{A} = [a_{ij}]$

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

Basic definitions and properties

- ▶ Matrix addition $\mathbf{A} + \mathbf{B} = [a_{ij} + b_{ij}]$

Basic definitions and properties

- ▶ Matrix addition $\mathbf{A} + \mathbf{B} = [a_{ij} + b_{ij}]$
- ▶ Scalar multiplication $\gamma \mathbf{B} = [\gamma b_{ij}]$

Basic definitions and properties

- ▶ Matrix addition $\mathbf{A} + \mathbf{B} = [a_{ij} + b_{ij}]$
- ▶ Scalar multiplication $\gamma \mathbf{B} = [\gamma b_{ij}]$
- ▶ Matrix-vector multiplication $\mathbf{A}\mathbf{v} = \left[\sum_k a_{ik} v_k \right]$

Basic definitions and properties

- ▶ Matrix addition $\mathbf{A} + \mathbf{B} = [a_{ij} + b_{ij}]$
- ▶ Scalar multiplication $\gamma \mathbf{B} = [\gamma b_{ij}]$
- ▶ Matrix-vector multiplication $\mathbf{A}\mathbf{v} = \begin{bmatrix} \sum_k a_{ik} v_k \end{bmatrix}$
- ▶ Matrix-matrix multiplication $\mathbf{AB} = \begin{bmatrix} \sum_k a_{ik} b_{kj} \end{bmatrix}$

Basic definitions and properties

► Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^{\top} = [a_{ji}]$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^{\top} = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{A}\mathbf{B})^{\top} = \mathbf{B}^{\top}\mathbf{A}^{\top}$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^\top = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{A}\mathbf{B})^\top = \mathbf{B}^\top \mathbf{A}^\top$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^\top$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^{\top} = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{A}\mathbf{B})^{\top} = \mathbf{B}^{\top}\mathbf{A}^{\top}$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^{\top}$
- ▶ Determinant $|\mathbf{A}|$ (defined and explained later)

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^\top = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^\top$
- ▶ Determinant $|\mathbf{A}|$ (defined and explained later)
- ▶ Determinant $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^\top = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^\top$
- ▶ Determinant $|\mathbf{A}|$ (defined and explained later)
- ▶ Determinant $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$
- ▶ Determinant $|\mathbf{A}^\top| = |\mathbf{A}|$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^{\top} = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{AB})^{\top} = \mathbf{B}^{\top}\mathbf{A}^{\top}$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^{\top}$
- ▶ Determinant $|\mathbf{A}|$ (defined and explained later)
- ▶ Determinant $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$
- ▶ Determinant $|\mathbf{A}^{\top}| = |\mathbf{A}|$
- ▶ Trace $\text{Tr}(\mathbf{A}) = \sum_k a_{kk}$

Basic definitions and properties

- ▶ Inverse $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$
- ▶ Transpose $\mathbf{A}^\top = [a_{ji}]$
- ▶ Transpose of products $(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$
- ▶ Symmetric matrices $\mathbf{A} = \mathbf{A}^\top$
- ▶ Determinant $|\mathbf{A}|$ (defined and explained later)
- ▶ Determinant $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$
- ▶ Determinant $|\mathbf{A}^\top| = |\mathbf{A}|$
- ▶ Trace $\text{Tr}(\mathbf{A}) = \sum_k a_{kk}$
- ▶ $\text{Tr}(\mathbf{AB}) = \text{Tr}(\mathbf{BA})$

Matrix Determinant

For a 2x2 matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, the determinant is defined as:

$$|A| = ad - bc$$

For a 3x3 matrix $A = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}$, the determinant can be calculated as:

$$|A| = a(ei - fh) - b(di - fg) + c(dh - eg)$$

Laplace Expansion

The Laplace expansion is a method to compute the determinant of a matrix by expanding it along a row or column. For any $n \times n$ matrix A , the determinant can be expressed as:

$$|A| = \sum_{j=1}^n (-1)^{i+j} a_{ij} |M_{ij}|$$

where:

- ▶ a_{ij} is an element of A
- ▶ M_{ij} is the $(n-1) \times (n-1)$ submatrix obtained by removing the i -th row and j -th column from A

This recursive process continues until reaching 2×2 matrices, where the determinant is directly calculable.

Outline

Basics

Spectral decomposition

Positive definite matrices

Square root matrices

Eigenvalues and eigenvectors

- ▶ Let $\mathbf{A} = [a_{i,j}]$ be an $n \times n$ matrix
- ▶ \mathbf{A} is said to have an eigenvalue λ and (non-zero) eigenvector \mathbf{x} corresponding to λ if

$$\mathbf{Ax} = \lambda\mathbf{x}.$$

- ▶ Eigenvalues are the λ values that solve the determinantal equation $|\mathbf{A} - \lambda\mathbf{I}| = 0$.

Spectral decomposition of symmetric matrices

- ▶ The Spectral decomposition theorem says that every square and symmetric matrix $\mathbf{A} = [a_{ij}]$ may be written

$$\mathbf{A} = \mathbf{C}\mathbf{D}\mathbf{C}^\top,$$

- ▶ The columns of \mathbf{C} are the eigenvectors of \mathbf{A}
- ▶ The diagonal matrix \mathbf{D} contains the corresponding eigenvalues

$$\mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}$$

- ▶ The eigenvectors may be chosen to be orthonormal, so that $\mathbf{C}\mathbf{C}^\top = \mathbf{C}^\top\mathbf{C} = \mathbf{I}$.
- ▶ Note the useful property: $|\mathbf{A}| = \prod_{i=1}^n \lambda_i$

Outline

Basics

Spectral decomposition

Positive definite matrices

Square root matrices

Positive definite matrices

The $n \times n$ matrix \mathbf{A} is said to be positive definite if

$$\mathbf{y}^\top \mathbf{A} \mathbf{y} > 0$$

for all $n \times 1$ vectors $\mathbf{y} \neq \mathbf{0}$.

Positive semidefinite matrices

The $n \times n$ matrix \mathbf{A} is said to be positive semidefinite if

$$\mathbf{y}^T \mathbf{A} \mathbf{y} \geq 0$$

for all $n \times 1$ vectors $\mathbf{y} \neq \mathbf{0}$.

Example: Show $\mathbf{X}^\top \mathbf{X}$ is positive semidefinite

Let \mathbf{X} be an $n \times p$ matrix of real constants and \mathbf{y} be $p \times 1$. Then $\mathbf{Z} = \mathbf{X}\mathbf{y}$ is $n \times 1$, and

$$\begin{aligned} & \mathbf{y}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{y} \\ = & (\mathbf{X}\mathbf{y})^\top (\mathbf{X}\mathbf{y}) \\ = & \mathbf{Z}^\top \mathbf{Z} \\ = & \sum_{i=1}^n z_i^2 \geq 0 \end{aligned}$$

Some properties of symmetric positive definite matrices

For a symmetric matrix,

Positive definite



All eigenvalues positive

Showing Positive definite \Rightarrow Eigenvalues positive

Let \mathbf{A} be symmetric and positive definite.

- ▶ Spectral decomposition says $\mathbf{A} = \mathbf{C}\mathbf{D}\mathbf{C}^\top$.
- ▶ Using $\mathbf{y}^\top \mathbf{A} \mathbf{y} > 0$, let \mathbf{y} be an eigenvector, say the third one.
- ▶ Because eigenvectors are orthonormal,

$$\begin{aligned}\mathbf{y}^\top \mathbf{A} \mathbf{y} &= \mathbf{y}^\top \mathbf{C} \mathbf{D} \mathbf{C}^\top \mathbf{y} \\ &= \begin{pmatrix} 0 & 0 & 1 & \cdots & 0 \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} \\ &= \lambda_3 \\ &> 0\end{aligned}$$

Outline

Basics

Spectral decomposition

Positive definite matrices

Square root matrices

Square root matrices

For symmetric, non-negative definite matrices

Define

$$\mathbf{D}^{1/2} = \begin{pmatrix} \sqrt{\lambda_1} & 0 & \cdots & 0 \\ 0 & \sqrt{\lambda_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sqrt{\lambda_n} \end{pmatrix}$$

So that

$$\begin{aligned} \mathbf{D}^{1/2} \mathbf{D}^{1/2} &= \begin{pmatrix} \sqrt{\lambda_1} & 0 & \cdots & 0 \\ 0 & \sqrt{\lambda_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sqrt{\lambda_n} \end{pmatrix} \begin{pmatrix} \sqrt{\lambda_1} & 0 & \cdots & 0 \\ 0 & \sqrt{\lambda_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sqrt{\lambda_n} \end{pmatrix} \\ &= \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = \mathbf{D} \end{aligned}$$

For a non-negative definite, symmetric matrix \mathbf{A}

Define

$$\mathbf{A}^{1/2} = \mathbf{C}\mathbf{D}^{1/2}\mathbf{C}^\top$$

So that

$$\begin{aligned}\mathbf{A}^{1/2}\mathbf{A}^{1/2} &= \mathbf{C}\mathbf{D}^{1/2}\mathbf{C}^\top\mathbf{C}\mathbf{D}^{1/2}\mathbf{C}^\top \\ &= \mathbf{C}\mathbf{D}^{1/2}\mathbf{I}\mathbf{D}^{1/2}\mathbf{C}^\top \\ &= \mathbf{C}\mathbf{D}^{1/2}\mathbf{D}^{1/2}\mathbf{C}^\top \\ &= \mathbf{C}\mathbf{D}\mathbf{C}^\top \\ &= \mathbf{A}\end{aligned}$$

Cholesky decomposition

- Define lower triangular matrix \mathbf{L}

$$\mathbf{L} = \begin{bmatrix} L_{11} & 0 & \cdots & 0 \\ L_{21} & L_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{bmatrix}$$

so that $\mathbf{A} = \mathbf{L}\mathbf{L}^\top$ (here, \mathbf{A} is non-negative definite).

- Cholesky algorithm computes \mathbf{L} from \mathbf{A}

- $|\mathbf{A}| = |\mathbf{L}\mathbf{L}^\top| = |\mathbf{L}|^2 = \left(\prod_{i=1}^n L_{ii} \right)^2$