

Parametric Models and Maximum Likelihood Estimation

Motonobu Kanagawa

Introduction to Statistics, EURECOM

March 30, 2023

Outline

- 1 Estimation in Parametric Models
- 2 Maximum Likelihood Estimation
- 3 MLE as Kullback-Leibler Divergence Minimization
- 4 Consistency of MLE
- 5 Conclusions and Further Readings

Density Estimation Problem

- Let P be an **unknown** probability distribution on a measurable set $\mathcal{X} \subset \mathbb{R}^d$.
- Assume that P has a **probability density function** $p : \mathcal{X} \rightarrow \mathbb{R}$.
- Given i.i.d. data X_1, \dots, X_n from the unknown P , we are interested in **estimating the density function** p .
- This is the task of **density estimation**.

Notation

We may write $X_1, \dots, X_n \sim p$ (i.i.d.) with the density function p .

Density Estimation Problem

- There are mainly two approaches to this problem: **parametric** and **nonparametric**.

Parametric approach

- Define a model of a **finite degree of freedom** for the unknown density p .
- This is called a **parametric model**, and indexed by a finite number of **parameters**.
- Assumptions of the model are often made on the **shape** of the unknown density p .
- Density estimation is done by **estimating the parameters** from the data $X_1, \dots, X_n \sim p$.

Density Estimation Problem

Nonparametric approach

- Define a model with **infinite degree of freedom**.
- **Increase the complexity** of the model **as more data become available**.
- Assumptions of the model are often made on the **smoothness** of the unknown density p .
- e.g., kernel density estimation [Silverman, 1986].

- In this course we'll only focus on the **parametric approach** (while the nonparametric approach is also important).

Parameter Approach to Density Estimation

- In the parametric approach, we define a **parametric model** for the unknown density function p generating data X_1, \dots, X_n .

Parametric Model

- Let Θ be a set of **parameter vectors** (e.g., $\Theta \subset \mathbb{R}^q$).
- For each $\theta \in \Theta$, define a **probability density function** $p_\theta : \mathcal{X} \rightarrow [0, \infty)$.
- A **parametric model** is defined as the **set** of such density functions:

$$\mathcal{P}_\Theta := \{p_\theta \mid \theta \in \Theta\}.$$

Parametric Approach to Density Estimation

Remarks on the Term “Parametric Models”

- The parametric model can be seen as a function $f : \mathcal{X} \times \Theta \rightarrow [0, \infty)$ such that

$$f(x, \theta) := p_{\theta}(x), \quad x \in \mathcal{X}, \theta \in \Theta.$$

We may say that f is a parametric model.

- Alternatively, regarding $\theta \in \Theta$ as a variable, we also say p_{θ} is a parametric model for simplicity.

Parametric Approach to Density Estimation

- The parametric model \mathcal{P}_Θ should be designed so that the unknown density p belongs to \mathcal{P}_Θ , i.e., $p \in \mathcal{P}_\Theta$;

- $p \in \mathcal{P}_\Theta = \{p_\theta \mid \theta \in \Theta\}$ is equivalent to the existence of some $\theta^* \in \Theta$ such that

$$p = p_{\theta^*} \in \mathcal{P}_\Theta.$$

- We may call such θ^* the true parameter (vector).

- Therefore the model \mathcal{P}_Θ should reflect our knowledge/belief about the unknown p .

Parametric Approach to Density Estimation

- If $p \in \mathcal{P}_\Theta = \{p_\theta \mid \theta \in \Theta\}$, we say that the model \mathcal{P}_Θ is **correctly specified**.
 - In this case, estimation of the unknown density $p = p_{\theta^*}$ can be done by **estimating the true parameter θ^*** from the data X_1, \dots, X_n .
- If $p \notin \mathcal{P}_\Theta = \{p_\theta \mid \theta \in \Theta\}$, we say that the model \mathcal{P}_Θ is **misspecified**.

Example: Gaussian Models

- Recall that the density function of a **Gaussian distribution** on $\mathcal{X} = \mathbb{R}$ is given by

$$p_{\text{gauss}}(x; \mu, \sigma^2) := \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right), \quad x \in \mathbb{R}$$

where

- $\mu \in \mathbb{R}$ is the **mean** of p_{gauss}
- $\sigma^2 > 0$ is the **variance** of p_{gauss} .

Example: Gaussian Density Models




Figure 1: Gaussian density functions; From Wikipedia “Normal distribution.”

Example: Gaussian Models

There are several ways to define a probabilistic model.

1. Parametrizing the mean

- Assume that we know/believe that the **variance** of the unknown density p is σ^2 .
- Then we can define a parametric model p_θ by **treating the mean μ as a parameter θ** :

$$p_\theta(x) := p_{\text{gauss}}(x; \theta, \sigma^2).$$

- In this case, the parameter set may be defined as $\Theta := [-a, a] \subset \mathbb{R}$ for some $a > 0$.

Remarks

- Note that the **definition** of the parameter set Θ is a **part of the model**.
- e.g., the choice of the interval $[-a, a]$ implicitly represents our belief that the mean μ of p should satisfy $\mu \in [-a, a]$.

Example: Gaussian Models

2. Parametrizing both the mean and variance

- We can treat **both mean μ and variance σ^2** as parameters.
- In this case, we can define a parametric model p_θ as

$$P_\theta(x) := p_{\text{gauss}}(x; \theta_1, \theta_2).$$

where

$$\theta := (\theta_1, \theta_2) \in \Theta \subset \mathbb{R} \times (0, \infty).$$

- The parameter set may be defined as

$$\Theta := [-a, a] \times [b, c] \subset \mathbb{R} \times (0, \infty)$$

for some $a, b, c > 0$.

Example: Gaussian Models

- By using the Gaussian model p_θ , we implicitly makes several assumptions about the unknown p :

Assumptions about the true p made in the Gaussian model

- 1 There is only **one mode** (or the “bump”) in the density p .
- 2 $X \sim p$ may take an **arbitrarily large value**, but with an **exponentially small probability**.
- 3 $X \sim p$ takes both **positive** and **negative** values.
- 4 All the **moments** of p exist: $-\infty < \mathbb{E}_{X \sim p}[X^k] < \infty$ for all $k \in \mathbb{N}$.

- Gaussian models have been widely used in practice.

— This is because there are several **mathematically and computationally convenient** properties (we'll see this soon).

Example: Gaussian Mixture Models

- Assume instead that we know/believe that there **two bumps** in the true density p .

- Then the use of the above Gaussian model might be inappropriate.
- We can instead consider a **two-component Gaussian mixture model**:

$$p_{\theta}(x) := \frac{1}{2}p_{\text{gauss}}(x; \theta_1, \theta_2) + \frac{1}{2}p_{\text{gauss}}(x; \theta_3, \theta_4), \quad x \in \mathbb{R}$$

where

$$\theta := (\theta_1, \theta_2, \theta_3, \theta_4) \in \Theta \subset \mathbb{R} \times (0, \infty) \times \mathbb{R} \times (0, \infty).$$

Outline

- 1 Estimation in Parametric Models
- 2 Maximum Likelihood Estimation**
- 3 MLE as Kullback-Leibler Divergence Minimization
- 4 Consistency of MLE
- 5 Conclusions and Further Readings

Maximum Likelihood Estimation

- **Maximum likelihood estimation (MLE)** is a classic but still widely used approach to **estimating the parameter of a parametric model**, advocated by [Fisher, 1922].
- The approach defines an estimator of the true parameter θ^* (in the correctly specified case) as a maximizer of the **likelihood function**.

Notation

In this lecture, we will use the notation

$$\arg \max_{\theta \in \Theta} A(\theta) = \left\{ \theta^* \in \Theta \mid A(\theta^*) = \max_{\theta \in \Theta} A(\theta) \right\}$$

as a **set** of elements in Θ that maximize the objective function $A(\theta)$.

- Thus, if there are **multiple** maximizers of $A(\theta)$, then $\arg \max_{\theta \in \Theta} A(\theta)$ consists of **multiple** elements.

($\arg \min_{\theta \in \Theta} A(\theta)$ is defined in a similar way.)

Likelihood Function

- Let $X_1, \dots, X_n \in \mathcal{X} \subset \mathbb{R}^d$ be i.i.d. data.

Likelihood Function

For a parametric model $\mathcal{P}_\Theta := \{p_\theta(x) \mid \theta \in \Theta\}$, the **likelihood function** $\ell_n : \Theta \rightarrow [0, \infty)$ for the data X_1, \dots, X_n is defined by:

$$\ell_n(\theta) := \prod_{i=1}^n p_\theta(X_i), \quad \theta \in \Theta.$$

Remarks

- $\ell_n(\theta)$ is a function of the parameter vector $\theta \in \Theta$ (with X_1, \dots, X_n being fixed).
- $\ell_n(\theta)$ is **not** a **probability density function** of $\theta \in \Theta$.

In fact, its integral may not be 1:

$$\int \ell_n(\theta) d\theta = \int \left(\prod_{i=1}^n p_\theta(X_i) \right) d\theta \neq 1.$$

Maximum Likelihood Estimation (MLE)

- Let $X_1, \dots, X_n \sim p$ be i.i.d. data from the unknown density function p .
- Let $\ell_n(\theta) := \prod_{i=1}^n p_\theta(X_i)$ be the likelihood function.

Maximum Likelihood Estimation (MLE)

- Assume that there exists a **true parameter** $\theta^* \in \Theta$ such that $p = p_{\theta^*}$ (i.e., the correctly specified case $p \in \mathcal{P}_\Theta = \{p_\theta \mid \theta \in \Theta\}$).
- MLE defines an **estimate** $\hat{\theta}_n$ of the **true parameter** $\theta^* \in \Theta$ as a solution to the following **optimization problem**:

$$\hat{\theta}_n \in \arg \max_{\theta \in \Theta} \ell_n(\theta) := \left\{ \theta' \in \Theta \mid \ell_n(\theta') = \max_{\theta \in \Theta} \ell_n(\theta) \right\}$$

- i.e., the estimate $\hat{\theta}_n$ is a **maximizer of the likelihood function**:

$$\ell_n(\hat{\theta}_n) = \max_{\theta \in \Theta} \ell_n(\theta).$$

Maximum Likelihood Estimation: Intuition

- We may interpret a parametric model p_θ as a **conditional probability density function** on \mathcal{X} given $\theta \in \Theta$:

$$p(x | \theta) := p_\theta(x), \quad x \in \mathcal{X}, \theta \in \Theta.$$

- Thus, the likelihood function may be interpreted as the **conditional joint probability density** of i.i.d. observations X_1, \dots, X_n :

$$\ell_n(\theta) = \prod_{i=1}^n p_\theta(X_i) = \prod_{i=1}^n p(X_i | \theta).$$

- Note that the **product form** is due to the **independence** assumption of X_1, \dots, X_n .

- Thus the MLE may be interpreted as searching for the parameter vector θ^* that **maximizes the conditional probability (density) of the data** X_1, \dots, X_n .
- This interpretation of the likelihood function becomes important in Bayesian inference (we'll see this in a coming lecture)

MLE as Maximizing the Log Likelihood Function

- MLE can be equivalently defined as a maximizer of the **log likelihood**:

$$\hat{\theta} \in \arg \max_{\theta \in \Theta} \ell_n(\theta) = \arg \max_{\theta \in \Theta} \log \ell_n(\theta)$$

- This is because the logarithm is a **monotonically increasing function**:

$$\log(t) > \log(s) \iff t > s > 0.$$

- The log likelihood function is often easier to work with in practice, because the **product** becomes the **sum**.

$$\log \ell_n(\theta) = \log \prod_{i=1}^n p_{\theta}(X_i) = \sum_{i=1}^n \log p_{\theta}(X_i).$$

- We'll also see the use of log likelihood leads to a **deeper understanding of MLE** [Akaike, 1998].

Example: MLE with a Gaussian Density Model

- Consider a Gaussian density model on $\mathcal{X} = \mathbb{R}$, with a parametrized mean $\mu = \theta$ and a fixed variance $\sigma^2 > 0$:

$$p_{\theta}(x) := p_{\text{gauss}}(x; \theta, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \theta)^2}{2\sigma^2}\right)$$

- Assume that i.i.d. data X_1, \dots, X_n are given.

Example: MLE with a Gaussian Density Model

- Then the log likelihood function is given as

$$\begin{aligned}\log \ell_n(\theta) &:= \sum_{i=1}^n \log p_{\text{gauss}}(X_i; \theta, \sigma^2) \\&= \sum_{i=1}^n \log \left(\frac{1}{\sqrt{2\pi\sigma^2}} \exp \left(-\frac{(X_i - \theta)^2}{2\sigma^2} \right) \right) \\&= \sum_{i=1}^n \left(\log \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right) - \frac{(X_i - \theta)^2}{2\sigma^2} \right) \\&= n \log \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right) - \sum_{i=1}^n \frac{(X_i - \theta)^2}{2\sigma^2}.\end{aligned}$$

Example: MLE with a Gaussian Density Model

- To obtain the maximizer, compute the derivative w.r.t. θ and equate it to 0:

$$\frac{d \log \ell_n(\theta)}{d\theta} = \sum_{i=1}^n \frac{(X_i - \theta)}{\sigma^2} = 0.$$

- Solving this leads to the maximum likelihood estimator for the mean:

$$\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i.$$

- This is the **empirical average** of X_1, \dots, X_n !

Exercise

- Think about **why** the empirical average is obtained as MLE for the Gaussian model.
(Hint: recall that the empirical average can be given as a solution to the **least-squares problem**).

Example: MLE with a Gaussian Density Model

Exercise

- Consider the Gaussian model with both mean $\mu = \theta_1$ and variance $\sigma^2 = \theta_2$ parametrized:

$$p_{\theta}(x) := p_{\text{gauss}}(x; \theta_1, \theta_2) = \frac{1}{\sqrt{2\pi\theta_2}} \exp\left(-\frac{(x - \theta_1)^2}{2\theta_2}\right)$$

- Show that the MLE for $\theta = (\theta_1, \theta_2)$ with i.i.d. observations X_1, \dots, X_n is given by

$$\hat{\theta} = (\hat{\theta}_1, \hat{\theta}_2) = \left(\frac{1}{n} \sum_{i=1}^n X_i, \frac{1}{n} \sum_{i=1}^n (X_i - \hat{\theta}_1)^2 \right).$$

Illustration




Figure 2: [Efron and Hastie, 2016, Fig 4.1]

Maximum Likelihood Estimation (MLE)

- In general, this optimization problem of MLE has no analytical solution (e.g., consider Gaussian mixture models)
- In that case, one needs to use **numerical optimization**. e.g.,
 - Gradient descent (see the Optim course for details.)
 - Expectation-Maximization (EM) algorithm.
- In this lecture, we'll study **statistical properties** of MLE, **assuming that we can obtain the maximizer $\hat{\theta}_n$** .

Outline

- 1 Estimation in Parametric Models
- 2 Maximum Likelihood Estimation
- 3 MLE as Kullback-Leibler Divergence Minimization**
- 4 Consistency of MLE
- 5 Conclusions and Further Readings

MLE as Kullback-Leibler (KL) Divergence Minimization

- Here we'll see an interpretation of MLE as searching for the parameter $\theta \in \Theta$ that minimizes the KL divergence between the true density p and the model density p_θ [Akaike, 1998].
- This interpretation is very important, because it provides an understanding of the MLE in the misspecified case $p \notin \mathcal{P}_\Theta$:
- To describe this, we'll look at the definition and properties of the KL divergence.

Kullback-Leibler (KL) Divergence

- KL divergence quantifies the **discrepancy** between two probability density functions.

Kullback-Leibler (KL) Divergence

- Let p and q be probability density functions on $\mathcal{X} \subset \mathbb{R}^d$ such that $p(x)/q(x) < \infty$ for all $x \in \mathcal{X}$.
- Then the KL divergence between p and q is defined as

$$\begin{aligned} KL(p||q) &:= \int p(x) \log \frac{p(x)}{q(x)} dx \\ &= \int p(x) \log p(x) dx - \int p(x) \log q(x) dx. \end{aligned}$$

Intuition: KL Divergence as a Discrepancy Measure

- If $KL(p||q)$ is **large**, then p and q are **very different**;
- If $KL(p||q)$ is **small**, then p and q are **similar**.

Properties of the KL Divergence

Nonnegativity

- The KL divergence only take **non-negative** values: for any density functions p and q ,

$$KL(p||q) \geq 0.$$

- This can be seen as follows:

$$\begin{aligned} KL(p||q) &= \int p(x) \log \left(\frac{p(x)}{q(x)} \right) dx = - \int p(x) \log \left(\frac{q(x)}{p(x)} \right) dx \\ &\geq - \log \left(\int p(x) \frac{q(x)}{p(x)} dx \right) \\ &= - \log \left(\int q(x) dx \right) = - \log(1) = 0 \end{aligned}$$

where the inequality follows from **Jensen's inequality** and $\log(t)$ being a convex function of $t > 0$ (see e.g., [Berger, 1985, Sec 1.8]).

Properties of the KL Divergence

KL Divergence as a Discrepancy

- $KL(p||q) = 0$ if and only if $p = q$ (almost everywhere).

-“if” part can be shown easily: If $p = q$, we have

$$KL(p||q) = \int p(x) \log \frac{p(x)}{q(x)} dx = \int p(x) \log(1) dx = 0.$$

Asymmetry of the KL Divergence

- That the KL divergence is not symmetric: in general,

$$KL(p||q) \neq KL(q||p)$$

- Therefore, KL divergence is not a distance (metric) between probability density functions. (A distance measure needs to be symmetric).

Properties of the KL Divergence

- KL divergence has its origin in [Information Theory](#).
- Indeed, the KL divergence can be written as

$$\begin{aligned}
 KL(p||q) &:= \int p(x) \log \frac{p(x)}{q(x)} dx \\
 &= - \underbrace{\int p(x) \log \frac{1}{p(x)} dx}_{\text{Entropy of } p} + \underbrace{\int p(x) \log \frac{1}{q(x)} dx}_{\text{Cross Entropy of } p \text{ and } q}
 \end{aligned}$$

- For details see e.g. [Gray, 2011] and the InfoTheo course.

Example: KL Divergence between Gaussians

- Consider the KL divergence between two **Gaussian densities** p and q on $\mathcal{X} := \mathbb{R}$.

KL Divergence between Univariate Gaussians

- Let $p(x) := p_{\text{gauss}}(x; \mu_p, \sigma_p^2)$ with mean $\mu_p \in \mathbb{R}$ and variance $\sigma_p^2 > 0$;
- Let $q(x) := p_{\text{gauss}}(x; \mu_q, \sigma_q^2)$ with mean $\mu_q \in \mathbb{R}$ and variance $\sigma_q^2 > 0$.
- Then the KL divergence between p and q is given by

$$KL(p\|q) = \frac{1}{2} \left(\frac{(\mu_p - \mu_q)^2}{\sigma_q^2} + \log \left(\frac{\sigma_q^2}{\sigma_p^2} \right) + \frac{\sigma_p^2}{\sigma_q^2} - 1 \right)$$

Exercise. Prove this.

Example: KL Divergence between Gaussians

- For instance, consider the equal variance case $\sigma_p^2 = \sigma_q^2 =: \sigma^2$.
- Then, the KL divergence simplifies to

$$\begin{aligned} KL(p\|q) &= \frac{1}{2} \left(\frac{(\mu_p - \mu_q)^2}{\sigma^2} + \log \left(\frac{\sigma^2}{\sigma^2} \right) + \frac{\sigma^2}{\sigma^2} - 1 \right) \\ &= \frac{(\mu_p - \mu_q)^2}{2\sigma^2}. \end{aligned}$$

- We can make the following observations:
 - As difference between the means μ_p and μ_q approaches 0, the KL divergence converges to 0.

$$KL(p\|q) \rightarrow 0 \quad \text{as} \quad (\mu_p - \mu_q)^2 \rightarrow 0.$$

- As the variance σ^2 increases, the KL divergence converges to 0

$$KL(p\|q) \rightarrow 0 \quad \text{as} \quad \sigma^2 \rightarrow \infty$$

MLE as KL Divergence Minimization

- We now look at a connection between MLE and KL divergence.
- The estimate $\hat{\theta}$ of MLE can be obtained as

$$\begin{aligned}\hat{\theta} &\in \arg \max_{\theta \in \Theta} \log \ell_n(\theta) = \arg \max_{\theta \in \Theta} \log \prod_{i=1}^n p_{\theta}(X_i) \\ &= \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log p_{\theta}(X_i) = \arg \max_{\theta \in \Theta} \frac{1}{n} \sum_{i=1}^n \log p_{\theta}(X_i).\end{aligned}$$

- The objective function in the last expression is the **empirical average** of the log density $\log p_{\theta}(x)$ with the **i.i.d. data** $X_1, \dots, X_n \sim p$:

$$\frac{1}{n} \sum_{i=1}^n \log p_{\theta}(X_i).$$

MLE as KL Divergence Minimization

- Thus, we can interpret the objective function of MLE as an **empirical approximation** to the **expected log density**:

$$\frac{1}{n} \sum_{i=1}^n \log p_{\theta}(X_i) \approx \mathbb{E}_{X \sim p}[\log p_{\theta}(X)] = \int (\log p_{\theta}(x)) p(x) dx.$$

where the expectation is with respect to the **true unknown density**, $X \sim p$.

- Thus, under an appropriate **identifiability condition** (introduced later), we may expect that

$$\hat{\theta}_n \approx \theta^* \in \arg \max_{\theta \in \Theta} \int p(x) \log p_{\theta}(x) dx.$$

where $\theta^* \in \Theta$ is a maximizer of the expected log density.

- (We use the notation θ^* as for the “true parameter” intentionally, for a reason that will be clear later).

MLE as KL Divergence Minimization

- We show that this maximizer θ^* is the **minimizer of the KL divergence** between the **true density p** and the **model density p_θ** :

$$\begin{aligned}
 \theta^* &\in \arg \max_{\theta \in \Theta} \int p(x) \log p_\theta(x) dx \\
 &= \arg \min_{\theta \in \Theta} - \int p(x) \log p_\theta(x) dx \\
 &= \arg \min_{\theta \in \Theta} - \int p(x) \log p_\theta(x) dx + \int p(x) \log p(x) dx \\
 &= \arg \min_{\theta \in \Theta} \int p(x) (-\log p_\theta(x) + \log p(x)) dx \\
 &= \arg \min_{\theta \in \Theta} \int p(x) \log \frac{p(x)}{p_\theta(x)} dx \\
 &= \arg \min_{\theta \in \Theta} KL(p \| p_\theta).
 \end{aligned}$$

MLE as KL Divergence Minimization

- Thus, we have:

$$\theta^* \in \arg \max_{\theta \in \Theta} \int p(x) \log p_{\theta}(x) dx = \arg \min_{\theta \in \Theta} KL(p \| p_{\theta}).$$

- Therefore, the estimate $\hat{\theta}_n$ of MLE

$$\hat{\theta}_n \in \arg \max_{\theta \in \Theta} \frac{1}{n} \sum_{i=1}^n \log p_{\theta}(X_i)$$

can be seen as an approximation to the **minimizer of the KL divergence**:

$$\theta^* \in \arg \min_{\theta \in \Theta} KL(p \| p_{\theta})$$

- We will look at closely the **conditions** required for this interpretation to be valid.
- These are conditions required for MLE to “succeed”, thus providing a guideline for the use of MLE in practice.

Outline

- 1 Estimation in Parametric Models
- 2 Maximum Likelihood Estimation
- 3 MLE as Kullback-Leibler Divergence Minimization
- 4 Consistency of MLE**
- 5 Conclusions and Further Readings

Consistency of MLE

- We saw that MLE may be interpreted as an **estimator** of the **optimal parameter** θ^* given by

$$\theta^* \in \arg \max_{\theta \in \Theta} \int p(x) \log p_{\theta}(x) dx = \arg \min_{\theta \in \Theta} KL(p \| p_{\theta}).$$

- We'll investigate the consistency of the estimate $\hat{\theta}_n$ in estimating such θ^* in a large sample limit $n \rightarrow \infty$.
 - This is based on [White 82]; see this paper for details.
- The purpose is to clarify **conditions** under which MLE “works well.”
- To this end, we'll introduce several **assumptions** (= conditions).

Assumptions on the Data Distribution

Assumption 1 (Data and the True Density)

The data $X_1, \dots, X_n \in \mathcal{X} \subset \mathbb{R}^d$ are i.i.d. with a distribution P with a density function p .

Assumptions on the Parametric Model

Assumption 2 (Model)

- The parameter set $\Theta \subset \mathbb{R}^q$ is compact.
— i.e., Θ is a **bounded** and **closed** subset.
- For every $x \in \mathcal{X}$, the mapping

$$\theta \rightarrow p_{\theta}(x)$$

is a **continuous function** of $\theta \in \Theta$.

Consequence of the Continuity Assumption

- The **likelihood function** $\ell_n(\theta) = \prod_{i=1}^n p_{\theta}(X_i)$ is a **continuous function** of $\theta \in \Theta$, because the mapping

$$\theta \rightarrow p_{\theta}(X_i)$$

is **continuous** for all $i = 1, \dots, n$.

Assumptions on the Parametric Model

- Assumption 2 guarantees that the **maximum** of the likelihood function is **bounded**: i.e.,

$$\max_{\theta \in \Theta} \ell_n(\theta) = \max_{\theta \in \Theta} \prod_{i=1}^n p_{\theta}(X_i) < \infty.$$

This follows from

- 1 The likelihood function $\ell_n(\theta)$ is a continuous function of $\theta \in \Theta$;
- 2 Θ is compact;
- 3 **Extreme value theorem** (a general fact): a **continuous function** on a **compact domain** is **bounded**.

Assumptions on the Parametric Model




Figure 3: From Wikipedia “Extreme value theorem”

Assumptions on the Parametric Model

- Thus, Assumption 2 guarantees that MLE

$$\hat{\theta}_n \in \arg \max_{\theta \in \Theta} \ell_n(\theta)$$

is well-defined.

- If Assumption 2 is not satisfied, then we may have

$$\max_{\theta \in \Theta} \ell_n(x) = \infty$$

- In this case, MLE $\hat{\theta}_n \in \arg \max_{\theta \in \Theta} \ell_n(x)$ is not well-defined.

Example where MLE is not Well-Defined

- Consider a 2-component Gaussian mixture model;

$$p_{\theta}(x) = \frac{1}{2}p_{\text{gauss}}(x; \theta_1, \theta_2) + \frac{1}{2}p_{\text{gauss}}(x; \theta_3, \theta_4),$$

with

$$\theta := (\theta_1, \theta_2, \theta_3, \theta_4) \in \Theta \subset \mathbb{R} \times (0, \infty) \times \mathbb{R} \times (0, \infty).$$

- Define the parameter set Θ as

$$\Theta := [-a, a] \times (0, c] \times [-a, a] \times (0, c]$$

for constants $a, c > 0$.

- In this case, Θ is **not closed** and thus **not compact**.
- Therefore Assumption 2 is **not satisfied**.

Example where MLE is not Well-Defined

- We'll show that in this case the **maximum** of the likelihood function is **unbounded**:

$$\max_{\theta \in \Theta} \ell_n(\theta) = \infty,$$

and thus **MLE is not well-defined**.

Example where MLE is not Well-Defined

- Define $\theta_1 := X_k$ for with $k \in \{1, \dots, n\}$ arbitrary, and fix θ_3 and θ_4 .

$$\begin{aligned} p_{\theta}(X_k) &= \frac{1}{2} p_{\text{gauss}}(X_k; \theta_1, \theta_2) + \frac{1}{2} p_{\text{gauss}}(X_k; \theta_3, \theta_4) \\ &= \frac{1}{2} \frac{1}{\sqrt{2\pi\theta_2^2}} \exp\left(-\frac{(X_k - \theta_1)^2}{2\theta_2^2}\right) + \frac{1}{2} p_{\text{gauss}}(X_k; \theta_3, \theta_4) \\ &= \frac{1}{2} \frac{1}{\sqrt{2\pi\theta_2^2}} + \frac{1}{2} p_{\text{gauss}}(X_k; \theta_3, \theta_4). \end{aligned}$$

- Taking the limit $\theta_2 \rightarrow +0$ (i.e., the variance θ_2 going to 0), we have

$$\lim_{\theta_2 \rightarrow +0} p_{\theta}(X_k) = \lim_{\theta_2 \rightarrow +0} \left(\frac{1}{2} \frac{1}{\sqrt{2\pi\theta_2^2}} + \frac{1}{2} p_{\text{gauss}}(X_k; \theta_3, \theta_4) \right) = \infty.$$

- The limit $\theta_2 \rightarrow +0$ can be taken, because $\theta_2 \in (0, c]$.

Example where MLE is not Well-Defined

- On the other hand, for all $i \neq k$ we have

$$\begin{aligned} p_{\theta}(X_i) &= \frac{1}{2} p_{\text{gauss}}(X_i; \theta_1, \theta_2) + \frac{1}{2} p_{\text{gauss}}(X_i; \theta_3, \theta_4) \\ &\geq \frac{1}{2} p_{\text{gauss}}(X_i; \theta_3, \theta_4). \end{aligned}$$

- Therefore,

$$\begin{aligned} \lim_{\theta_2 \rightarrow +0} \ell_n(\theta) &= \lim_{\theta_2 \rightarrow +0} \prod_{i=1}^n p_{\theta}(X_i) = \lim_{\theta_2 \rightarrow +0} p_{\theta}(X_k) \prod_{i \neq k}^n p_{\theta}(X_i) \\ &\geq \left(\lim_{\theta_2 \rightarrow +0} p_{\theta}(X_k) \right) \prod_{i \neq k} \frac{1}{2} p_{\text{gauss}}(X_i; \theta_3, \theta_4) = \infty. \end{aligned}$$

This implies that

$$\max_{\theta \in \Theta} \ell_n(\theta) \geq \lim_{\theta_2 \rightarrow +0} \ell_n(\theta) = \infty.$$

Example where MLE is not Well-Defined

- This example shows that MLE is **not always well-defined**.
- We need to be careful about **how the parameter set Θ is defined**.

Exercise

Construct other examples where MLE is not well-defined.

Assumptions for the KL Divergence to be Well-Defined

Assumption 3 (The existence of the KL divergence)

- The true density $p(x)$ satisfies

$$-\infty < \int p(x) \log p(x) dx < \infty.$$

- For the model $p_\theta(x)$, there exists a function $g : \mathcal{X} \rightarrow [0, \infty)$ such that

$$|\log p_\theta(x)| \leq g(x) \quad \text{for all } x \in \mathcal{X} \text{ and } \theta \in \Theta$$

and

$$\int g(x)p(x)dx < \infty.$$

Assumptions for the KL Divergence to be Well-Defined

- The latter condition implies that

$$\begin{aligned} \left| \int p(x) \log p_{\theta}(x) dx \right| &< \int p(x) |\log p_{\theta}(x)| dx \\ &\leq \int p(x) g(x) dx < \infty. \end{aligned}$$

- Therefore, the above conditions imply that the KL divergence

$$\begin{aligned} KL(p \| p_{\theta}) &= \int p(x) \log \frac{p(x)}{p_{\theta}(x)} dx \\ &= \int p(x) \log p(x) dx - \int p(x) \log p_{\theta}(x) dx \end{aligned}$$

is finite and thus well-defined.

Exercise

- Construct examples of p and p_{θ} for which the KL divergence **cannot be defined**.

Assumption for the Identifiability

Assumption 4 (Identifiability)

- Expected log density $\int p(x) \log p_{\theta}(x) dx$ has a **unique** maximizer $\theta^* \in \Theta$: i.e.,

$$\int p(x) \log p_{\theta^*}(x) dx > \int p(x) \log p_{\theta}(x) dx \quad \text{for all } \theta \in \Theta \text{ with } \theta \neq \theta^*.$$

- In other words, θ^* is the **unique** minimizer of the KL-divergence:

$$\begin{aligned} KL(p \| p_{\theta^*}) &= \int p(x) \log p(x) dx - \int p(x) \log p_{\theta^*}(x) dx \\ &< \int p(x) \log p(x) dx - \int p(x) \log p_{\theta}(x) dx = KL(p \| p_{\theta}) \\ &\quad \text{for all } \theta \in \Theta \text{ with } \theta \neq \theta^*. \end{aligned}$$

- In this case, we call the model $P_{\Theta} = \{p_{\theta} \mid \theta \in \Theta\}$ is **identifiable** with respect to p .

Assumption for the Identifiability

- If Assumption 4 (identifiability) is true, the notation

$$\theta^* = \arg \max_{\theta \in \Theta} \int p(x) \log p_{\theta}(x) dx = \arg \min_{\theta \in \Theta} KL(p \| p_{\theta})$$

is justified (because the “argmax” only consists of one element, θ^*).

- Assumption 4 enables us to **define θ^*** as the **quantity of interest** (or the **estimand**) in statistical estimation.
- Thus, we can discuss the “consistency” of the MLE $\hat{\theta}_n \rightarrow \theta^*$ as $n \rightarrow \infty$.
- This will be important in particular
 - when we are interested in the optimal parameter θ^* **itself**; and
 - when we want to perform **hypothesis testing** regarding θ^* .

Interpretation of the Optimal Parameter θ^*

- Let's consider what the optimal parameter θ^* is.
- Assume that the KL divergence between the true unknown density p and the optimal model density p_{θ^*} is zero:

$$KL(p \| p_{\theta^*}) = 0.$$

- In this case,
 - We have $p = p_{\theta^*}$, because $KL(p \| p_{\theta^*}) = 0$ if and only if $p = p_{\theta^*}$.
 - Therefore, $p \in \mathcal{P}_{\Theta} = \{p_{\theta} \mid \theta \in \Theta\}$ i.e., the model \mathcal{P}_{Θ} is correctly specified.
- Thus, we can interpret θ^* as the true parameter in this case.
- The convergence of MLE $\hat{\theta}_n \rightarrow \theta^*$ implies that the MLE is consistent in estimating the true parameter θ^* .

Interpretation of the Optimal Parameter θ^*

Summary

- $KL(p||p_{\theta^*}) = 0$ corresponds to the correctly specified case $p \in \mathcal{P}_{\Theta}$.
- Since $p = p_{\theta^*}$, the optimal parameter θ^* is interpreted as the true parameter.

Interpretation of the Optimal Parameter θ^*




Figure 4: When $KL(p \| p_{\theta^*}) = 0$ (correctly specified case)

Interpretation of the Optimal Parameter θ^*

- Assume the KL divergence between the true density p and the optimal model density p_{θ^*} is **larger than zero**:

$$KL(p \| p_{\theta^*}) = \min_{\theta \in \Theta} KL(p \| p_{\theta}) > 0,$$

- In this case,
 - we have $p \neq p_{\theta^*}$, i.e., the optimal model density p_{θ^*} **does not match** the true density p ;
 - thus $p \notin \mathcal{P}_{\Theta} = \{p_{\theta} \mid \theta \in \Theta\}$, i.e., the **model \mathcal{P}_{Θ} is misspecified**.
- In this case, we can interpret p_{θ^*} as the **best approximation** to the true density p as measured by the KL divergence.
- Thus, we can interpret θ^* as the **parameter that gives the best approximation** of the model \mathcal{P}_{Θ} to the true p .

Interpretation of the Optimal Parameter θ^*

Summary

- $KL(p\|p_{\theta^*}) > 0$ corresponds to the misspecified case $p \notin \mathcal{P}_\Theta$.
- Since $KL(p\|p_{\theta^*}) = \min_{\theta \in \Theta} KL(p\|p_\theta)$, the optimal parameter θ^* is interpreted as the parameter that gives the best approximation p_{θ^*} to the true density p under the KL divergence.

Interpretation of the Optimal Parameter θ^*




Figure 5: When $KL(p \parallel p_{\theta^*}) > 0$ (model misspecification).

Example where the Model is not Identifiable

- Consider a 2-component Gaussian mixture model;

$$p_{\theta}(x) = \frac{1}{2}p_{\text{gauss}}(x; \theta_1, \theta_2) + \frac{1}{2}p_{\text{gauss}}(x; \theta_3, \theta_4)$$

with

$$\theta = (\theta_1, \theta_2, \theta_3, \theta_4) \in \Theta \subset \mathbb{R} \times (0, \infty) \times \mathbb{R} \times (0, \infty).$$

- Define the parameter set Θ by

$$\Theta := [-a, a] \times [b, c] \times [-a, a] \times [b, c]$$

for constants $a, b, c > 0$.

- The model is **not identifiable**, because **switching** (θ_1, θ_2) and (θ_3, θ_4) produces the **same density function**.

Example where the Model is not Identifiable

- To show this, let

$$(\mu_1, \sigma_1^2) \in [-a, a] \times [b, c], \quad (\mu_2, \sigma_2^2) \in [-a, a] \times [b, c]$$

be arbitrary constants such that $\sigma_1^2 \neq \sigma_2^2$.

- Then, for $\theta^* := (\mu_1, \sigma_1^2, \mu_2, \sigma_2^2)$, we have

$$p_{\theta^*}(x) = \frac{1}{2}p_{\text{gauss}}(x; \mu_1, \sigma_1^2) + \frac{1}{2}p_{\text{gauss}}(x; \mu_2, \sigma_2^2)$$

- For $\tilde{\theta}^* := (\mu_2, \sigma_2^2, \mu_1, \sigma_1^2)$

$$p_{\tilde{\theta}^*}(x) = \frac{1}{2}p_{\text{gauss}}(x; \mu_2, \sigma_2^2) + \frac{1}{2}p_{\text{gauss}}(x; \mu_1, \sigma_1^2)$$

- Thus, we have

$$p_{\theta^*} = p_{\tilde{\theta}^*} \quad \text{while} \quad \theta^* \neq \tilde{\theta}^*.$$

- Therefore the mixture model with this parameter set Θ is **not identifiable**.

Example where the Model is not Identifiable

- A simple trick to make this model identifiable is to restrict the parameter set Θ .
- For instance, if we define the parameter set as

$$\Theta := \{(\theta_1, \theta_2, \theta_3, \theta_4) \in [-a, a] \times [b, c] \times [-a, a] \times [b, c] \mid \theta_2 < \theta_4\}$$

then the mixture model becomes identifiable.

- This corresponds to assuming that one mixture component has a smaller variance than the other.

Exercise

Construct other examples where the model is not identifiable.

MLE Consistency Theorem

Theorem: Consistency of MLE (Theorem 2.2 of [White, 1982])

- Suppose that Assumptions 1, 2, 3 and 4 are satisfied.
- Let

$$\hat{\theta}_n \in \arg \max_{\theta \in \Theta} \ell_n(\theta) = \arg \max_{\theta \in \Theta} \prod_{i=1}^n p_{\theta}(X_i)$$

be the MLE with i.i.d. data $X_1, \dots, X_n \sim p$.

- Let $\theta^* \in \Theta$ be the optimal parameter

$$\theta^* = \arg \max_{\theta \in \Theta} \int p(x) \log p_{\theta}(x) dx = \arg \min_{\theta \in \Theta} KL(p \| p_{\theta})$$

- Then $\hat{\theta}_n$ converges to θ^* almost surely: i.e.,

$$\Pr(\lim_{n \rightarrow \infty} \hat{\theta}_n = \theta^*) = 1.$$

MLE Consistency Theorem

The proof idea is that

- 1 First show that

$$\frac{1}{n} \sum_{i=1}^n \log p_{\theta}(X_i) \rightarrow \int p(x) \log p_{\theta}(x) dx \quad \text{as } n \rightarrow \infty$$

uniformly for all $\theta \in \Theta$.

- 2 Then conclude that

$$\hat{\theta}_n \in \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log p_{\theta}(X_i) \rightarrow \theta^* = \arg \max_{\theta \in \Theta^*} \int p(x) \log p_{\theta}(x) dx.$$

as $n \rightarrow \infty$.

Outline

- 1 Estimation in Parametric Models
- 2 Maximum Likelihood Estimation
- 3 MLE as Kullback-Leibler Divergence Minimization
- 4 Consistency of MLE
- 5 Conclusions and Further Readings

Conclusions

- MLE can be understood as searching for a model density that **best approximates** the true density in terms of the **KL divergence**.
- MLE makes sense also in the **misspecified case** where the true density does not belong to the model class.
- MLE is **not always consistent**; we need **conditions = assumptions**.
- These conditions provide a **guideline** for designing your parametric model.

Conclusions

More generic takeaways:

- A role of convergence analysis is to understand **conditions** under which **the method of interest works well**.
- Even the MLE - one of the simplest approaches - requires several conditions.
- So please always try to understand conditions under which your favorite statistical/ML method should work!

Further Readings

- [Fisher, 1922, Section 6].
- [White, 1982]
- [Efron and Hastie, 2016, Chapter 4]



Akaike, H. (1998).

Information theory and an extension of the maximum likelihood principle.

In *Selected Papers of Hirotugu Akaike*, pages 199–213. Springer.



Berger, J. O. (1985).

Statistical Decision Theory and Bayesian Analysis.

Springer Science & Business Media.



Efron, B. and Hastie, T. (2016).

Computer Age Statistical Inference.

Cambridge University Press.



Fisher, R. A. (1922).

On the mathematical foundations of theoretical statistics.

Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character,
222(594-604):309–368.



Gray, R. M. (2011).

Entropy and Information Theory.
Springer Science & Business Media.



Silverman, B. W. (1986).
Density Estimation for Statistics and Data Analysis.
Chapman and Hall.



Van der Vaart, A. W. (1998).
Asymptotic statistics.
Cambridge University Press.



White, H. (1982).
Maximum likelihood estimation of misspecified models.
Econometrica, pages 1–25.