

**Example**


Illustrate the effects of sampling and rectangular windowing on a continuous-time sine wave and its Fourier transform.

**NB** the difference between  $X_C(\omega)$  and  $\hat{X}_C(\omega)$

The discrete Fourier transform

206


*Solution over page*



- (a) In the example  $x(t)$  is a sine wave with frequency  $\omega_0$  rad/s  
It has a Fourier transform with impulses at  $\pm\omega_0$  rad/s

Sampling at a rate of  $\omega_s$  is akin to multiplying by infinite train of impulses  
Multiplication in the time domain is convolution in the  $f$  domain so

- (b) To obtain the FT of  $x_c(t)$  we convolve the transforms of  $x(t)$  &  $\delta_T(t)$



The FT of  $\delta_T(t)$  is a train of impulses  
in the  $f$  domain at  $\pm\omega_s, \pm 2\omega_s, \dots$


- (c)  $\hat{x}_c(t)$  is obtained by multiplying  $x_c(t)$  by a window of width  $T$  seconds  
 $w_T(t)$  has a  $\text{sinc}(x)$  shape - see lecture 3.

- (d) we then convolve the transform of the sampled analogue signal  
with that of the rectangular window.

The result is that  $\hat{X}_c(\omega)$  is a similar but distorted version  
of  $X_c(\omega)$

### Example

Explain the operation of the 8-point DFT by a complex input phasor with an integer number of cycles within the 8-sample input data sequence.



The discrete Fourier transform


217

again notice how the speed of rotation increases going down the matrix.

the phasors rotate clockwise  
input rotates anticlockwise

Can you explain the symmetry about  $w_s$ , i.e. aliasing.

NB that the last row appears to rotate anticlockwise  
- explain aliasing about  $w_s$



The second row finds the component at 1 cycle per block length

eg for a simple input signal a sampled complex phasor with a period of 8 samples

$$x(n) = \exp(j 2\pi n/8) = \cos(2\pi n/8) + j \sin(2\pi n/8)$$

The second row of the matrix represents a filter ~~is~~ matched to a centre frequency corresponding to a component at one rotational cycle of the phasor per block length.

All values of  $x(\cdot)$  are rotated as shown to give a large value to  $X(1)$

What happens for  $X(2) \dots X(7)$ ?

What happens for  $\theta \neq 0$