

**Essential Mathematical Methods for Engineers  
(MathEng)  
EXAM**

December 2023

Duration: 2 hrs, all documents and calculators permitted

ATTEMPT ALL QUESTIONS – ANSWER IN ENGLISH




- 
1. Using Euler's identity (or any other appropriate method), write down an expression for the complex Fourier series of the signal  $x(t)$ :

$$x(t) = 3 \cos(5t) + 4 \sin(10t)$$

[5 marks]

- 
2. Develop an expression for the Fourier Transform of the signal  $x(t)$  illustrated in Figure Q2 below.



**Figure Q2**

[6 marks]

- 
3. A linear, time-invariant system has the following transfer function:

$$H(s) = \frac{10(s + 100)}{s^2 + 2s + 100}$$

- (a) Derive an expression for  $H(s)$  in the usual, normal form.
- (b) Determine the frequency-invariant gain  $K$  and the position of any poles and zeros.
- (c) Sketch a Bode plot of the magnitude-frequency response.
- (d) Sketch a Bode plot of the phase-frequency response.

[8 marks]

- 
4. Sketch magnitude and phase responses for a sampled data system with a pair of complex conjugate zeros and two poles at the origin.




Figure Q4

[4 marks]

- 
5. A random variable  $X$  is uniformly distributed between  $x = 0$  and  $x = 1$ . Via any appropriate method, determine the expected value  $E[Y]$  of  $Y = \exp(X)$ .

[5 marks]

- 
6. Identify the pivots and free variables of the following two matrices  $A$  and  $B$ . Following the method which we studied in class, find the special solution corresponding to each free variable and, by combining the special solutions, describe every solution to  $Ax=0$  and  $Bx=0$ .

$$A = \begin{bmatrix} 1 & 2 & 2 & 4 & 6 \\ 1 & 2 & 3 & 6 & 9 \\ 0 & 0 & 1 & 2 & 3 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 2 & 4 & 2 \\ 0 & 4 & 4 \\ 0 & 8 & 8 \end{bmatrix}$$

[7 marks]

- 
7. For a projection matrix  $P = A(A^T A)^{-1} A^T$ , show that  $P^2 = P$  and then explain, in terms of the column space of  $P$ , why projections  $Pb$  and  $P(Pb)$  give identical results.

[5 marks]

---

---

**Table of selected Laplace transforms**

$$X(s) = \int_{-\infty}^{\infty} x(t) \exp(-st) dt$$

N.B.: lower limit is 0 for one-sided Laplace transform

$$x(t) = \frac{1}{2\pi j} \int_{\sigma-j\infty}^{\sigma+j\infty} X(s) \exp(st) ds$$

$x(t) \ (t \geq 0)$	$X(s)$
$\delta(t)$	1
$\delta(t - \alpha)$	$\exp(-\alpha s)$
1 (unit step)	$\frac{1}{s}$
$t$ (unit ramp)	$\frac{1}{s^2}$
$\exp(-\alpha t)$	$\frac{1}{s + \alpha}$
$t \exp(-\alpha t)$	$\frac{1}{(s + \alpha)^2}$
$\sin(\alpha t)$	$\frac{\alpha}{s^2 + \alpha^2}$
$\cos(\alpha t)$	$\frac{s}{s^2 + \alpha^2}$
$e^{-\alpha t} \sin(\alpha t)$	$\frac{\omega}{(s + \alpha)^2 + \omega^2}$
$e^{-\alpha t} \cos(\alpha t)$	$\frac{s + \alpha}{(s + \alpha)^2 + \omega^2}$

---



**Bode plots**

Poles or zeros on the real axis:

$$(s + a) = a \left( \frac{s}{a} + 1 \right) = \frac{1}{\tau} (\tau s + 1)$$

Complex conjugate poles (or zeros):

$$(s^2 + As + B) = (s^2 + 2\zeta\omega_0 s + \omega_0^2) = \omega_0^2 ((s/\omega_0)^2 + 2\zeta(s/\omega_0) + 1)$$



---

**Table of selected z-transforms**

$$x_c(t) = \sum_{n=0}^{\infty} x(n\Delta t)\delta(t - n\Delta t)$$

$$X_c(s) = \sum_{n=0}^{\infty} x(n\Delta t)\exp(-n\Delta ts)$$

$$X_c(s) = X(z)|_{z=e^{\Delta ts}}$$

$$X_c(\omega) = X_s(s)|_{s=j\omega}$$

$$X(\omega) = X(z)|_{z=\exp(\Delta t j\omega)}$$

$$X(z) = \sum_{n=0}^{\infty} x(n\Delta t)z^{-n}$$

$x(n) \ (n \geq 0)$	$X(z)$
$\delta(n)$ unit pulse	1
$\delta(n-m)$	$z^{-m}$
1 (unit step)	$\frac{z}{z-1}$
$n$ (unit ramp)	$\frac{z}{(z-1)^2}$
$\exp(-\alpha n)$	$\frac{z}{(z-e^{-\alpha})}$
$n \exp(-\alpha n)$	$\frac{e^{-\alpha} z}{(z-e^{-\alpha})^2}$
$\sin(\beta n)$	$\frac{z \sin(\beta)}{z^2 - 2z \cos(\beta) + 1}$
$\cos(\beta n)$	$\frac{z^2 - z \cos(\beta)}{z^2 - 2z \cos(\beta) + 1}$
$e^{-\alpha n} \sin(\beta n)$	$\frac{ze^{-\alpha} \sin(\beta)}{z^2 - 2ze^{-\alpha} \cos(\beta) + e^{-2\alpha}}$
$e^{-\alpha n} \cos(\beta n)$	$\frac{z^2 - ze^{-\alpha} \cos(\beta)}{z^2 - 2ze^{-\alpha} \cos(\beta) + e^{-2\alpha}}$

---

**Table of selected Fourier transform pairs**

Function	$x(t)$	$X(\omega)$
Rectangular function of width $\tau$	$\Pi(t/\tau)$	$\tau \operatorname{sinc}(\omega\tau/2)$
Triangular function of width $2\tau$	$\Lambda(t/\tau)$	$\tau \operatorname{sinc}^2(\omega\tau/2)$
Train of impulses every $\Delta t$	$\delta_T(t)$	$2\pi/\Delta t \sum_n \delta(\omega - 2\pi n/\Delta t)$

NB:  $\operatorname{sinc}(x) = \sin(\pi x)/\pi x$

NB:  $\operatorname{sa}(x) = \sin(x)/x$

---

**Euler's identity**

$$\exp(j\theta) = \cos \theta + j \sin \theta$$

$$\cos(n\omega_0 t) = \frac{\exp(jn\omega_0 t) + \exp(-jn\omega_0 t)}{2}$$

$$\sin(n\omega_0 t) = \frac{\exp(jn\omega_0 t) - \exp(-jn\omega_0 t)}{2j}$$

---

**Fourier series and transforms**

Trigonometric Fourier series	
$x(t) = \frac{A_0}{2} + \sum_{n=1}^{\infty} A_n \cos(n\omega_0 t) + B_n \sin(n\omega_0 t)$ $A_n = \frac{2}{T} \int_{-T/2}^{T/2} x(t) \cos(n\omega_0 t) dt$ $B_n = \frac{2}{T} \int_{-T/2}^{T/2} x(t) \sin(n\omega_0 t) dt$	
Complex Fourier series – periodic and continuous in time, discrete in frequency	
$x(t) = \sum_{n=-\infty}^{+\infty} X_n \exp(jn\omega_0 t)$	$X_n = \frac{1}{T} \int_{-T/2}^{T/2} x(t) \exp(-jn\omega_0 t) dt$
Fourier transform – continuous in time, continuous in frequency	
$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(\omega) \exp(j\omega t) d\omega$	$X(\omega) = \int_{-\infty}^{\infty} x(t) \exp(-j\omega t) dt$
Discrete-time Fourier transform – discrete in time, continuous and periodic in frequency	
$x(n\Delta t) = \frac{\Delta t}{2\pi} \int_{-\pi/\Delta t}^{\pi/\Delta t} X(\omega) \exp(jn\Delta t\omega) d\omega$	$X(\omega) = \sum_{n=-\infty}^{\infty} x(n\Delta t) \exp(-j\omega n\Delta t)$
Discrete Fourier transform – discrete and periodic in time and in frequency	
$x(n) = \frac{1}{N} \sum_{k=0}^{N-1} X(k) \exp\left(\frac{jnk2\pi}{N}\right)$	$X(k) = \sum_{n=0}^{N-1} x(n) \exp\left(\frac{-jnk2\pi}{N}\right)$

---

**Transformation of random variables**

$$f_Y(y) = \sum_{i=1}^N f_X(x_i) \left| \frac{dx_i}{dy} \right|_{x_i=g_i^{-1}(y)}$$

$$f_{UV}(u, v) = f_{XY}(x, y) \left| \frac{\partial(x, y)}{\partial(u, v)} \right|_{\substack{x=g_1^{-1}(u, v) \\ y=g_2^{-1}(u, v)}}$$

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix}$$