

C v3 Vertical Slice v1 - Stage 3 Human Audit

Automated validation: PASS (89 checks / 0 errors). Human audit: PENDING. Stage 4 not started.


Pilot coordinates are non-deployable and are not the production 361-point coordinate dataset.

Outcome summary

{"logical_points": 20, "generated_points": 0, "generated_coordinates": 0, "unresolved_stage3": 4, "blocked_by_stage2": 16, "legacy_coordinate_inputs": 0}

Suppressed-provenance watchpoint

{"suppressed_executable_operand_count": 0, "suppressed_status_aggregation_influence_count": 0, "suppressed_only_resolved_path_count": 0}



LI4: UNRESOLVED

Stage 2 primary: RESOLVED | blocker: QUALITATIVE_DIRECTION_HAS_NO_FROZEN_PHYSICAL_DISPLACEMENT_RULE

WHO source

location | S:LI4:location | page None | On the dorsum of the hand, radial to the midpoint of the second metacarpal bone.

Stage 3 trace

{ "directional_relation_ids": ["RL:S:LI4:location:00:relative-to"], "location_statement_id": "S:LI4:location", "rule": "Stage3 refuses nearest/distance heuristics for unquantified directional language." }

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

GB23: UNRESOLVED

Stage 2 primary: RESOLVED | blocker: APPROVED_CALIBRATION_HAS_NO_EXPLICIT_PHYSICAL_CUN_SCALE

WHO source

location | S:GB23:location | page None | In the lateral thoracic region, in the fourth intercostal space, 1 B-cun anterior to the midaxillary line.

Stage 3 trace

{"location_statement_id": "S:GB23:location", "measurement_ids": ["PM:S:GB23:location:00"], "non_metric_calibration_ids": ["PM:S:GB23:location:00"], "rule": "Stage3 does not infer metres-per-cun from atlas joint distances."}

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

BL40: UNRESOLVED

Stage 2 primary: RESOLVED | blocker: LEFT_ASPECT_RAY_ELIGIBLE_GEOMETRIC_HIT_CARDINALITY_2

WHO source

location | S:BL40:location | page None | On the posterior aspect of the knee, at the midpoint of the popliteal crease.

Stage 3 trace

{ "location_statement_id": "S:BL40:location", "primary_relation_ids": ["RL:S:BL40:location:00:midpoint-of-entity", "RL:S:BL40:location:01:surface-landmark"] }

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

TE6: UNRESOLVED

Stage 2 primary: RESOLVED | blocker: APPROVED_CALIBRATION_HAS_NO_EXPLICIT_PHYSICAL_CUN_SCALE

WHO source

location | S:TE6:location | page None | On the posterior aspect of the forearm, midpoint of the interosseous space between the radius and the ulna, 3 B-cun proximal to the dorsal wrist crease.

note | S:TE6:note:1 | page None | Note: 1 B-cun proximal to TE5, between the radius and the ulna, at the same level as tE7.

Stage 3 trace

{ "location_statement_id": "S:TE6:location", "measurement_ids": ["PM:S:TE6:location:00"], "non_metric_calibration_ids": ["PM:S:TE6:location:00"], "rule": "Stage3 does not infer metres-per-cun from atlas joint distances." }

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

HT7: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:HT7:location | page None | On the anteromedial aspect of the wrist, radial to the flexor carpi ulnaris tendon, on the palmar wrist crease.

note | S:HT7:note:1 | page None | Note: In the depression radial to the proxi-mal border of the pisiform bone, on the palmar wrist crease.

Stage 3 trace

```
{"location_statement_id": "S:HT7:location"}  
  
[ ] WHO semantics compatible with synthesis family  
[ ] hard constraints executable  
[ ] cardinality/laterality correct  
[ ] anatomy plausible  
[ ] aspect-directed ray projection correct  
[ ] skin triangle reconstruction correct  
[ ] projection budget acceptable  
[ ] unresolved cases not force-solved  
[ ] ACCEPT  
[ ] REJECT
```

ST1: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:ST1:location | page None | On the face, between the eyeball and the infraorbital margin, directly inferior to the pupil.

Stage 3 trace

```
{"location_statement_id": "S:ST1:location"}
```

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

GB14: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:GB14:location | page None | On the head, 1 B-cun superior to the eye-brow, directly superior to the centre of the pupil.

Stage 3 trace

```
{"location_statement_id": "S:GB14:location"}
```

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

LU6: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:LU6:location | page None | On the anterolateral aspect of the forearm, on the line connecting LU5 with LU9, 7 B-cun superior to the palmar wrist crease.

note | S:LU6:note:1 | page None | Note: LU6 is 5 B-cun inferior to LU5, 1 B-cun superior to the midpoint of the line connecting LU5 with LU9.

Stage 3 trace

```
{"location_statement_id": "S:LU6:location"}

[ ] WHO semantics compatible with synthesis family
[ ] hard constraints executable
[ ] cardinality/laterality correct
[ ] anatomy plausible
[ ] aspect-directed ray projection correct
[ ] skin triangle reconstruction correct
[ ] projection budget acceptable
[ ] unresolved cases not force-solved
[ ] ACCEPT
[ ] REJECT
```

LI7: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:LI7:location | page None | On the posterolateral aspect of the forearm, on the line connecting LI5 with LI11, 5 B-cun superior to the dorsal wrist crease.

Stage 3 trace

```
{"location_statement_id": "S:LI7:location"}
```

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

GB26: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:GB26:location | page None | On the lateral abdomen, inferior to the free extremity of the 11th rib, at the same level as the centre of umbilicus.

note | S:GB26:note:1 | page None | Note 1: GB26 can be located by first locat-ing the tenth rib, then searching for the free extremity of the 11th rib that is situated immediately below the inferior border of the costal arch.

note | S:GB26:note:2 | page None | Note 2: GB26 is inferior to LR13, at the same level as CV8.

Stage 3 trace

- { "location_statement_id": "S:GB26:location" }
- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

ST2: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:ST2:location | page None | On the face, in the infraorbital foramen.

Stage 3 trace

```
{"location_statement_id": "S:ST2:location"}
```

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

ST10: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:ST10:location | page None | In the anterior region of the neck, at the same level as the cricoid cartilage, just anterior to the border of the sternocleido-mastoid muscle.

Stage 3 trace

```
{"location_statement_id": "S:ST10:location"}
```

[] WHO semantics compatible with synthesis family

[] hard constraints executable

[] cardinality/laterality correct

[] anatomy plausible

[] aspect-directed ray projection correct

[] skin triangle reconstruction correct

[] projection budget acceptable

[] unresolved cases not force-solved

[] ACCEPT

[] REJECT

LI18: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:LI18:location | page None | On the anterior aspect of the neck, at the same level as the superior border of the thyroid cartilage, between the anterior and posterior borders of the sternocleidomas-toid muscle.

Stage 3 trace

```
{"location_statement_id": "S:LI18:location"}
```

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

LI17: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:LI17:location | page None | On the anterior aspect of the neck, at the same level as the cricoid cartilage, just posterior to the border of the sternocleido-mastoid muscle.

note | S:LI17:note:1 | page None | Note: Directly inferior to LI18, at the same level as St10.

Stage 3 trace

{ "location_statement_id": "S:LI17:location" }

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

BL17: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:BL17:location | page None | In the upper back region, at the same level as the inferior border of the spinous process of the seventh thoracic vertebra (T7), 1.5 B-cun lateral to the posterior median line.

note | S:BL17:note:1 | page None | Note: The inferior angle of the scapula is at the same level as the spinous process of the seventh thoracic vertebra.

Stage 3 trace

{ "location_statement_id": "S:BL17:location" }

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

BL23: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:BL23:location | page None | In the lumbar region, at the same level as the inferior border of the spinous process of the second lumbar vertebra (L2), 1.5 B-cun lateral to the posterior median line.

Stage 3 trace

- { "location_statement_id": "S:BL23:location" }
- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

BL25: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:BL25:location | page None | In the lumbar region, at the same level as the inferior border of the spinous process of the fourth lumbar vertebra (L4), 1.5 B-cun lateral to the posterior median line.

Stage 3 trace

- { "location_statement_id": "S:BL25:location" }
- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

TE20: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:TE20:location | page None | On the head, just superior to the auricular apex.

note | S:TE20:note:1 | page None | Note: With the auricle folded forward and pressed against the head, the point is located where the auricular apex touches the head.

Stage 3 trace

```
{"location_statement_id": "S:TE20:location"}
```

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

ST4: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:ST4:location | page None | On the face, 0.4 f-cun lateral to the angle of the mouth.

note | S:ST4:note:1 | page None | Note: Lateral to the angle of the mouth, the point is located in the nasolabial sulcus or on the continuation of the nasolabial sulcus.

Stage 3 trace

```
{"location_statement_id": "S:ST4:location"}
```

- [] WHO semantics compatible with synthesis family
- [] hard constraints executable
- [] cardinality/laterality correct
- [] anatomy plausible
- [] aspect-directed ray projection correct
- [] skin triangle reconstruction correct
- [] projection budget acceptable
- [] unresolved cases not force-solved
- [] ACCEPT
- [] REJECT

ST9: BLOCKED_BY_STAGE2

Stage 2 primary: UNRESOLVED | blocker: Stage2 primary location is not RESOLVED

WHO source

location | S:ST9:location | page None | In the anterior region of the neck, at the same level as the superior border of the thyroid cartilage, anterior to the sterno-cleidomastoid muscle, over the common carotid artery.

note | S:ST9:note:1 | page None | Note 1: The sternocleidomastoid muscle is more distinct when the head is turned to the opposite side against resistance.

note | S:ST9:note:2 | page None | Note 2: ST9 is located at the same level as LI18, SI16 and the superior border of the thyroid cartilage. St9 is located anterior to the sternocleidomastoid muscle, and SI16, posterior to the sternocleidomastoid muscle and LI18, between the anterior and pos-terior borders of the sternocleidomastoid muscle.

Stage 3 trace

```
{"location_statement_id": "S:ST9:location"}
```

[] WHO semantics compatible with synthesis family
[] hard constraints executable
[] cardinality/laterality correct
[] anatomy plausible
[] aspect-directed ray projection correct
[] skin triangle reconstruction correct
[] projection budget acceptable
[] unresolved cases not force-solved
[] ACCEPT
[] REJECT