

2/3 13

$$\begin{aligned} \text{№1. а)} x + 2x^2 + 3x^3 + 4x^4 + \dots + nx^n &= x(1 + 2x + 3x^2 + \dots + nx^{n-1}) = \\ &= x(x + x^2 + x^3 + \dots + x^n)' = x \left(\frac{1 - x^{n+1}}{1 - x} \right)' = \frac{x}{(1-x)^2} \cdot \\ &\cdot \left(-(n+1)x^n + 1 - x^{n+1} \right) \underset{n \rightarrow \infty}{\overset{|x| < 1}{=}} \frac{x}{(1-x)^2} \end{aligned}$$

б) Ряд Тейлора \Rightarrow числовая - e^x

№2.

$$\text{а)} \sum_{k \geq 0} \binom{n+k}{k} x^k$$

$$\text{Заметим, что } (1+x)^{-(n+1)} = \sum_{k=0}^{\infty} C_{-(n+1)}^k (-x)^k \equiv$$

$$C_{-(n+1)}^k = \frac{(-n)(-n-1)\dots(-n-k)}{k!} = (-1)^k \frac{n(n+1)\dots(n+k)}{k!} =$$

$$= (-1)^k C_{n+k}^k$$

$$\equiv \sum_{k=0}^{\infty} C_{n+k}^k x^k = \sum_{k=0}^{\infty} \binom{n+k}{k} x^k \Rightarrow \boxed{(1+x)^{-(n+1)}}$$

$$\text{б)} \sum_{k \geq 0} \binom{k}{n} x^k = \sum_{k \geq n} \binom{k}{n} x^k \stackrel{i=k-n}{=} \sum_{i \geq 0} C_{i+n}^n$$

$$x^{i+n} = x^n \sum_{i \geq 0} C_{i+n}^n x^i = \boxed{x^n (1+x)^{-(n+1)}} \quad (\text{см. пункт а)}$$

$$\text{в)} \sum_{k=1}^{\infty} (2k+1) x^k = 2 \sum_{k=1}^{\infty} k x^k + \sum_{k=1}^{\infty} x^k = \frac{2x}{(1-x)^2} + \frac{1}{1-x} -$$

$$-1 = \boxed{\frac{3x - x^2}{(1-x)^2}}$$

Донатик

№3. а) Рассм. $f(x) = \sum_{k=1}^n \binom{n}{k+1} x^{k+1}$

$$\int f(x) dx = \sum_{k=1}^n \binom{n}{k+1} \frac{1}{k+2} x^{k+2} + C_1 = g(x)$$

$$\int g(x) dx = \sum_{k=1}^n \binom{n}{k+1} \frac{1}{(k+2)(k+3)} x^{k+3} + C_1 x + C_2 = h(x)$$

$$\text{т.е. } h(1) = \sum_{k=1}^n \frac{\binom{n}{k+1}}{(k+2)(k+3)}, \text{ т.е. } C_1 = -C_2$$

$$f(x) = (1+x)^n - 1 - nx$$

$$\int f(x) dx = \frac{1}{n+1} (1+x)^{n+1} - x - n \frac{x^2}{2} + C_1 = g(x)$$

$$\int g(x) dx = \frac{(1+x)^{n+2}}{(n+1)(n+2)} - \frac{x^2}{2} - n \frac{x^3}{6} + C_1 x + C_2 = h(x)$$

$$h(1) = \frac{2^{n+2}}{(n+1)(n+2)} - \frac{1}{2} - \frac{n}{6}$$

б) $\sum_{k=0}^n k \binom{n}{k} 2^k$

$$f(x) = (1+x)^n = \sum_{k=0}^n C_n^k x^k = (1+x)^n$$

$$f'(x) = \sum_{k=1}^n k C_n^k x^{k-1} = n(1+x)^{n-1}$$

$$f'(2) = \sum_{k=1}^n k C_n^k 2^{k-1}$$

$$2f'(2) = 2n \cdot 3^{n-1} = \sum_{k=1}^n k C_n^k 2^k = \sum_{k=0}^n k C_n^k 2^k$$

Искомое - $2n \cdot 3^{n-1}$

в) $\sum_{k=0}^{99} \binom{99}{k} \binom{100}{k}$

$$(1+x)^{2n+1} = (1+x)^n (1+x)^{n+1} = \sum_{k=0}^n \binom{n}{k} x^k \sum_{k=0}^{n+1} \binom{n+1}{k} x^k$$

$$\sum_{k=0}^{2n+1} \binom{2n+1}{k} x^k = \sum_{k=0}^n \binom{n}{k} x^k \sum_{k=0}^{n+1} \binom{n+1}{k} x^k$$

коэф. при x^n : $\binom{2n+1}{n} = \sum_{k=0}^n \binom{n}{n-k} \binom{n+1}{k} =$

$$= \sum_{k=0}^n \binom{n}{k} \binom{n+1}{k} \Rightarrow$$

$$\Rightarrow \sum_{k=0}^{99} \binom{99}{k} \binom{100}{k} = \boxed{\binom{199}{99}}$$

№4. $f(x) = (1+x)^n = \sum_{k=0}^n \binom{n}{k} x^k$

$$f'(x) = n(1+x)^{n-1} = \sum_{k=0}^{n-1} k \binom{n}{k} x^{k-1}$$

$$f''(x) = n(n-1)(1+x)^{n-2} = \sum_{k=0}^n k(k-1) \binom{n}{k} x^{k-2}$$

$$f''(1) = n(n-1)2^{n-2} = \sum_{k=2}^n k(k-1) \binom{n}{k} =$$

$$= \sum_{k=2}^n k(k-1) \binom{n}{k} \blacksquare$$

№5. $g(x) = a_0 + (a_0 + a_1)x + (a_0 + a_1 + a_2)x^2 + \dots =$

$$= a_0(1+x+\dots+x^n) + a_1x(1+\dots+x^n) + a_2x^2(1+\dots+x^n) + \dots =$$

$$= a_0 \frac{1-x^{n+1}}{1-x} + a_1x \frac{1-x^{n+1}}{1-x} + \dots + a_n x^n \frac{1-x^{n+1}}{1-x} + \dots =$$

$$\frac{1}{1-x} (a_0 + a_1x + \dots + a_n x^n) \Rightarrow$$

\Rightarrow условие $a_0 + a_1x + \dots + a_n x^n = \boxed{g(x)(1-x)}$