

Prove the following propositions. Format your proof so each step of the proof is on its own line; each line should still be a complete sentence. Below, I have entered a nonsensical proof as a model.

Proposition 1. Suppose $a, b, c \in \mathbb{Z}$. If $a^2 + b^2 = c^2$, then a or b is even.

Proof.

Suppose $a^2 + b^2 = c^2$ and it's not the case that a or b is even.

Therefore, a and b are both odd.

So $a = 2x + 1$ and $b = 2y + 1$ for some integers x and y .

Either c is even or odd.

Case 1: c is even.

Then $c = 2z$ for some integer x .

The expression $a^2 + b^2 = c^2$ becomes $(2x + 1)^2 + (2y + 1)^2 = 4z^2$.

Expanding the expression yields $4x^2 + 4x + 1 + 4y^2 + 4y + 1 = 4z^2$.

Factoring yields $4(x^2 + y^2 + x + y) + 2 = 4z^2$.

Simplifying the expression shows $4k + 2 = 4j$ where k and j are the integers $x^2 + y^2 + x + y$ and $4z^2$ respectfully.

Observe that $2 = 4(k - j)$, thus $4|2$.

We have arrived at contradiction.

Case 2: c is odd.

Then $c = 2z + 1$ for some integer x .

The expression $a^2 + b^2 = c^2$ becomes $(2x + 1)^2 + (2y + 1)^2 = (2z + 1)^2$.

Expanded and factored yields $2(2x^2 + 2x + 2y^2 + 2y + 1) = 2(2z^2 + 2z) + 1$

Simplifying the expression shows that $2k = 2j + 1$ for integers k and j .

Therefore, an even number equals an odd number.

We have arrived at contradiction.

□

Proposition 2. Suppose $x, y \in \mathbb{Z}$. If $x + y$ is even, then x and y have the same parity.

Proof.

Suppose x and y do not have the same parity.

Then x is even and y odd without loss of generality.

So $x = 2a$ and $y = 2b + 1$ for some integers a and b .

The expression $x + y$ becomes $2a + 2b + 1 = 2(a + b) + 1$.

Therefore $x + y = 2m + 1$ where m is the integer $a + b$.

So, $x + y$ is odd. □

Proposition 3. If $a \equiv b \pmod{n}$, then $\gcd(a, n) = \gcd(b, n)$.

Proof.

Suppose $a \equiv b \pmod{n}$.

Then $n \mid (a - b)$.

Then $nx = a - b$ for some integer x .

The division algorithm states that:

$a = nq_1 + r_1$ where $q_1, r_1 \in \mathbb{Z}$ where $0 \leq r_1 < n$.

$b = nq_2 + r_2$ where $q_2, r_2 \in \mathbb{Z}$ where $0 \leq r_2 < n$.

So, $nx = nq_1 + r_1 - nq_2 - r_2$

$= n(q_1 - q_2) + r_1 - r_2$

Observe that $r_1 - r_2$ must be some multiple of n in order for the equality to hold.

So, $ny = r_1 - r_2$.

However, $-n < r_1 - r_2 < n$, so the only possible multiple is 0.

So, $n(0) = 0 = r_1 - r_2 \implies r_1 = r_2$.

Therefore they share the same remainder.

Moving forward: $r = r_1 = r_2$.

Continuing,

$$a = nq_1 + r$$

$$b = nq_2 + r$$

n and r may share prime factors, we can factor these out and replace them with their product p :

$$a = p(n'q_1 + r')$$

$$b = p(n'q_2 + r')$$

Where $n = n'p$ and $r = r'p$.

$\gcd(a, n) =$ greatest x where $x|a$ and $x|n$.

$\gcd(b, n) =$ greatest x where $x|b$ and $x|n$.

So, there exists some x and y where $x \neq y$, $x|a$, $x|n$, $y|b$, $y|n$.

□