

Cryopreservation of HD11 cells

Chicken macrophage-like HD11 cells are suspended in complete Roswell Park Memorial Institute (RPMI)-1640 medium with

1. 50% FBS
2. 10% DMSO

and stored at -140°C .

Thawing and Propagation

- a. The chicken macrophage-like cell line HD11, transformed by the avian leukemia virus strain MC29 was maintained in RPMI-1640 containing
 1. 25 mM HEPES
 2. 300 mg/l L-glutamine
 3. 2% chicken serum
 4. 8% fetal bovine serum
 5. antibiotics (10,000 U/ml penicillin and streptomycin)

at 41°C , 5% CO_2 , and 95% humidity.

- b. The cells should be passaged twice weekly by washing the cells in Dulbecco's phosphate-buffered saline without calcium and magnesium and detaching adherent cells using a 0.25% trypsin/EDTA solution supplemented with phenol red.
- c. The cells can be harvested for experiments after 3 to 20 passages – using trypsin/ EDTA
- d. Resuspend the cells to a concentration of 200,000 cells/mL in complete RPMI-1640 medium.

Harvesting and experimental use

- a. The cells are detached using trypsin/EDTA, harvested and washed with PBS.
- b. Aliquots of cell suspensions (2×10^5 cells/ml) are seeded into 96-well culture plates and cultured in RPMI-1640 supplemented with 8% FBS but without antibiotics.
- c. The cells were allowed to grow to about 85% confluence during an overnight incubation at 41°C , 5% CO_2 , and 95% humidity.
- d. The cells are exposed to antigens at a fixed concentration of $0.5\mu\text{L/mL}$ or a concentration gradient of $0.2 - 30\mu\text{L/mL}$ to create dose response curves.

- e. NO production by the cells can be measured after 48 hours of stimulation by Griess test.

Griess Test

- a. Griess reagent:
3g N-(1-naphthyl) ethylenediamine in 1L 2.5% phosphoric acid - Solution A
10g sulfanilamide in 1L 2.5% phosphoric acid - Solution B

Mix sol A and sol B in 1:1 ratio
- b. Transfer 50 μ L of supernatant from stimulated HD11 culture to a 96-well flat bottom cell culture plate.
- c. Add 50 μ L Griess reagent to each well.
The mixture will turn purple on reaction with nitrite ions in the cell culture supernatant.
- d. Read at an optical density of 540 nm

Relative Expression of iNOS and Cytokines

- a. HD11 cells are harvested, either 8 h after stimulation or at different time points between 0–48 h, using 200 μ L PBS+5 mM EDTA
- b. Centrifuge at 400 \times g.
- c. Discard the supernatant
- d. Resuspend the pellet in RLT buffer and stored at -20°C until further processing.
- e. After thawing, RNA is isolated
- f. Prepared cDNA
- g. RT-qPCR reactions were performed using 400 nM primers and SYBR Green Master Mix.
- h. Changes in gene expression over time upon stimulation are assessed using $t=0$ as a reference time point and expressed as $2^{-\Delta\Delta\text{Ct}}$ -values, according to the Livak method with Ct being the number of cycles before a signal above the threshold (background) level was reached.
- i. The results are normalized to gene expression levels of the housekeeping genes 28S and GAPDH.