

# **Numbers, Sequences and Series**

**Lecture Notes**

**Dr. Silvio Fanzon**

**19 Oct 2023**

**Academic Year 2023/24**

**Department of Mathematics**

**University of Hull**

# Table of contents

<b>Welcome</b>	<b>3</b>
Digital Notes . . . . .	3
Readings . . . . .	3
<b>1 Introduction</b>	<b>4</b>
<b>2 Preliminaries</b>	<b>12</b>
2.1 Sets . . . . .	12
2.2 Logic . . . . .	13
2.3 Operations on sets . . . . .	14
2.3.1 Union and intersection . . . . .	14
2.3.2 Inclusion and equality . . . . .	15
2.3.3 Infinite unions and intersections . . . . .	17
2.3.4 Complement . . . . .	18
2.3.5 Power set . . . . .	19
2.3.6 Product of sets . . . . .	20
2.4 Equivalence relation . . . . .	20
2.5 Order relation . . . . .	23
2.6 Intervals . . . . .	25
2.7 Functions . . . . .	26
2.8 Absolute value or Modulus . . . . .	30
2.9 Triangle inequality . . . . .	34
2.10 Proofs in Mathematics . . . . .	37
2.11 Induction . . . . .	40
<b>3 Real Numbers</b>	<b>44</b>
3.1 Fields . . . . .	44
3.2 Ordered fields . . . . .	51
3.3 Cut Property . . . . .	52
3.4 Supremum and infimum . . . . .	59
3.4.1 Upper bound, supremum, maximum . . . . .	60
3.4.2 Lower bound, infimum, minimum . . . . .	63
3.5 Completeness . . . . .	66
3.6 Equivalence of Completeness and Cut Property . . . . .	71
3.7 Axioms of Real Numbers . . . . .	75

<b>4</b>	<b>Properties of <math>\mathbb{R}</math></b>	<b>78</b>
4.1	Special subsets of $\mathbb{R}$ . . . . .	78
4.2	Archimedean Property . . . . .	78
4.3	Nested Interval Property . . . . .	82
4.4	Revisiting Sup and inf . . . . .	85
4.5	Density of $\mathbb{Q}$ in $\mathbb{R}$ . . . . .	89
4.6	Existence of Square Roots . . . . .	91
4.7	Cardinality . . . . .	91
<b>5</b>	<b>Complex Numbers</b>	<b>92</b>
<b>6</b>	<b>Sequences</b>	<b>93</b>
6.1	Example: Heron's Method . . . . .	93
6.1.1	Coding the Algorithm . . . . .	97
6.2	Fibonacci Sequence . . . . .	97
<b>7</b>	<b>Series</b>	<b>98</b>
	<b>License</b>	<b>99</b>
	Reuse . . . . .	99
	Citation . . . . .	99
	<b>References</b>	<b>100</b>

# Welcome

These are the Lecture Notes of **Numbers, Sequences and Series 400297** for T1 2023/24 at the University of Hull. I will follow these lecture notes during the course. If you have any question or find any typo, please email me at

[S.Fanzon@hull.ac.uk](mailto:S.Fanzon@hull.ac.uk)

Up to date information about the course, Tutorials and Homework will be published on the University of Hull **Canvas Website**

[canvas.hull.ac.uk/courses/67551](https://canvas.hull.ac.uk/courses/67551)

and on the **Course Webpage** hosted on my website

[silviofanzon.com/blog/2023/NSS](https://silviofanzon.com/blog/2023/NSS)

## Digital Notes

Digital version of these notes available at

[silviofanzon.com/2023-NSS-Notes](https://silviofanzon.com/2023-NSS-Notes)

## Readings

We will study the set of real numbers  $\mathbb{R}$ , and then sequences and series in  $\mathbb{R}$ . I will follow mainly the textbook by Bartle and Sherbert [2]. Another good reading is the book by Abbott [1]. I also point out the classic book by Rudin [3], although this is more difficult to understand.

**!** You are not expected to purchase any of the above books. These lecture notes will cover 100% of the topics you are expected to know in order to excel in the final exam.

# 1 Introduction

The first aim of this lecture notes is to rigorously introduce the set of **real numbers**, which is denoted by  $\mathbb{R}$ . But what do we mean by real numbers? To start our discussion, introduce the set of natural numbers (or non-negative integers)

$$\mathbb{N} = \{0, 1, 2, 3, 4, 5, \dots\}$$

On this set we have a notion of **sum** of two numbers, denoted as usual by

$$n + m$$

for  $n, m \in \mathbb{N}$ . Here the symbol  $\in$  denotes that  $m$  and  $n$  belong to  $\mathbb{N}$ . For example  $3 + 7$  results in 10.

## Question 1.1

Can the sum be inverted? That is, given any  $n, m \in \mathbb{N}$ , can you always find  $x \in \mathbb{N}$  such that

$$n + x = m? \tag{1.1}$$

Of course to invert (1.1) we can just perform a **subtraction**, implying that

$$x = m - n.$$

But there is a catch. In general  $x$  does not need to be in  $\mathbb{N}$ . For example, take  $n = 10$  and  $m = 1$ . Then  $x = -9$ , which does not belong to  $\mathbb{N}$ . Therefore the answer to Question 1.1 is **NO**.

To make sure that we can always invert the sum, we need to **extend** the set  $\mathbb{N}$ . This is done simply by introducing the set of **integers**

$$\mathbb{Z} := \{-n, n : n \in \mathbb{N}\},$$

that is, the set

$$\mathbb{Z} := \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}.$$

The sum can be extended to  $\mathbb{Z}$ , by defining

$$(-n) + (-m) := -(m + n) \tag{1.2}$$

for all  $m, n \in \mathbb{N}$ . Now every element of  $\mathbb{Z}$  possesses an **inverse**, that is, for each  $n \in \mathbb{Z}$ , there exists  $m \in \mathbb{Z}$ , such that

$$n + m = 0.$$

Can we characterize  $m$  explicitly? Of course! Seeing the definition at (1.2), we simply have

$$m = -n.$$

On the set  $\mathbb{Z}$  we can also define the operation of **multiplication**, in the usual way we learnt in school. For  $n, m \in \mathbb{Z}$ , we denote the multiplication by  $nm$  or  $n \cdot m$ . For example  $7 \cdot 2 = 14$  and  $1 \cdot (-1) = -1$ .

### Question 1.2

Can the multiplication in  $\mathbb{Z}$  be inverted? That is, given any  $n, m \in \mathbb{Z}$ , can you always find  $x \in \mathbb{Z}$  such that

$$nx = m? \tag{1.3}$$

To invert (1.3) if  $n \neq 0$ , we can just perform a **division**, to obtain

$$x = \frac{m}{n}.$$

But again there is a catch. Indeed taking  $n = 2$  and  $m = 1$  yields  $x = 1/2$ , which does not belong to  $\mathbb{Z}$ . The answer to Question 1.2 is therefore **NO**.

Thus, in order to invert the multiplication, we need to **extend** the set of integers  $\mathbb{Z}$ . This extension is called the set of **rational numbers**, defined by

$$\mathbb{Q} := \left\{ \frac{m}{n} : m, n \in \mathbb{Z}, n \neq 0 \right\}.$$

We then extend the operations of sum and multiplication to  $\mathbb{Q}$  by defining

$$\frac{m}{n} + \frac{p}{q} := \frac{mq + np}{nq}$$

and

$$\frac{m}{n} \cdot \frac{p}{q} := \frac{mp}{nq}$$

Now the multiplication is invertible in  $\mathbb{Q}$ . Specifically, each non-zero element has an inverse: the inverse of  $m/n$  is given by  $n/m$ .

To summarize, we have extended  $\mathbb{N}$  to  $\mathbb{Z}$ , and  $\mathbb{Z}$  to  $\mathbb{Q}$ . By construction we have

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q}.$$

Moreover **sum** and **product** are **invertible** in  $\mathbb{Q}$ . Now we are happy right? So and so.

**Question 1.3**

Can we draw the set  $\mathbb{Q}$ ?

It is clear how to draw  $\mathbb{Z}$ , as seen below.




Figure 1.1: Representation of integers  $\mathbb{Z}$

However  $\mathbb{Q}$  is much **larger** than the set  $\mathbb{Z}$  represented by the ticks in Figure 1.1. What do we mean by **larger**? For example, consider  $0 \in \mathbb{Q}$ .

**Question 1.4**

What is the number  $x \in \mathbb{Q}$  which is closest to 0?

There is no right answer to the above question, since whichever rational number  $m/n$  you consider, you can always squeeze the rational number  $m/(2n)$  in between:

$$0 < \frac{m}{2n} < \frac{m}{n}.$$

For example think about the case of the numbers

$$\frac{1}{n} \text{ for } n \in \mathbb{N}, n \neq 0.$$

Such numbers get arbitrarily close to 0, as depicted below.




Figure 1.2: Fractions  $\frac{1}{n}$  can get arbitrarily close to 0

Maybe if we do the same reasoning with other progressively smaller rational numbers, we manage to fill out the interval  $[0, 1]$ . In other words, we might conjecture the following.

**Conjecture 1.5**

Maybe  $\mathbb{Q}$  can be represented by a continuous line.

Do you think the above conjecture is true? If it was, mathematics would be quite boring. Indeed Conjecture 1.5 is **false**, as shown by the Theorem below.

**Theorem 1.6**

The number  $\sqrt{2}$  does not belong to  $\mathbb{Q}$ .

Theorem 1.6 is the reason why  $\sqrt{2}$  is called an **irrational number**. For reference, a few digits of  $\sqrt{2}$  are given by

$$\sqrt{2} = 1.414213562373095048 \dots$$

and the situation is as in the picture below.




Figure 1.3: Representing  $\sqrt{2}$  on the numbers line.

We can therefore see that Conjecture 1.5 is **false**, and  $\mathbb{Q}$  is not a line: indeed  $\mathbb{Q}$  has a **gap** at  $\sqrt{2}$ . Let us see why Theorem 1.6 is true.

**Proof:** Proof of Theorem 1.6

We prove that

$$\sqrt{2} \notin \mathbb{Q}$$

by **contradiction**.

Wait, what does this mean? Proving the claim by contradiction means assuming that the claim is **false**.

This means we **assume** that

$$\sqrt{2} \in \mathbb{Q}. \tag{1.4}$$

From this assumption we then start deducing other statements, hoping to encounter a statement which is **FALSE**. But if (1.4) leads to a false statement, then it must be that (1.4) is **FALSE**. Thus the contrary of (1.4) must hold, meaning that

$$\sqrt{2} \notin \mathbb{Q}$$

as we wanted to show. This would conclude the proof.

Now we need to actually show that (1.4) will lead to a contradiction. Since this is our first proof, let us take it slowly, step-by-step.



1. Assuming (1.4) just means that there exists  $q \in \mathbb{Q}$  such that

$$q = \sqrt{2}. \quad (1.5)$$

2. Since  $q \in \mathbb{Q}$ , by definition we have

$$q = \frac{m}{n}$$

for some  $m, n \in \mathbb{N}$  with  $n \neq 0$ .

3. Recalling (1.5), we then have

$$\frac{m}{n} = \sqrt{2}.$$

4. We can square the above equation to get

$$\frac{m^2}{n^2} = 2. \quad (1.6)$$

5. **Without loss of generality**, we can **assume** that  $m$  and  $n$  have no common factors.

Wait. What does Step 5 mean? You will encounter the sentence *without loss of generality* many times in mathematics. It is often abbreviated in **WLOG**. WLOG means that the assumption that follows is chosen arbitrarily, but does not affect the validity of the proof in general.

For example in our case we can assume that  $m$  and  $n$  have no common factor. This is because if  $m$  and  $n$  had common factors, then it would mean

$$m = a\tilde{m}, \quad n = a\tilde{n}$$

for some  $a \in \mathbb{N}$  with  $a \neq 0$ . Then

$$\frac{m}{n} = \frac{a\tilde{m}}{a\tilde{n}} = \frac{\tilde{m}}{\tilde{n}}.$$

Therefore by (1.6)

$$\frac{\tilde{m}^2}{\tilde{n}^2} = 2.$$

The proof can now proceed in the same way we would have proceeded from Step 4, but in addition we have the hypothesis that  $\tilde{m}$  and  $\tilde{n}$  have no common factors.

6. Equation (1.6) implies

$$m^2 = 2n^2. \quad (1.7)$$

Therefore the integer  $m^2$  is an even number.

Why is  $m^2$  even? As you already know, **even** numbers are

$$0, 2, 4, 6, 8, 10, 12, \dots$$

All these numbers have in common that they can be divided by 2, and so they can be written as

$$2p$$

for some  $p \in \mathbb{N}$ . For example 52 is even, because

$$52 = 2 \cdot 26.$$

Instead, **odd** numbers are

$$1, 3, 5, 7, 9, 11, \dots$$

These can be all written as

$$2p + 1$$

for some  $p \in \mathbb{N}$ . For example 53 is odd, because

$$53 = 2 \cdot 26 + 1.$$

7. Thus  $m$  is an even number, and so there exists  $p \in \mathbb{N}$  such that

$$m = 2p. \tag{1.8}$$

Why is (1.8) true? Let us see what happens if we take the square of an even number  $m = 2p$

$$m^2 = (2p)^2 = 4p^2 = 2(2p^2) = 2q.$$

Thus  $m^2 = 2q$  for some  $q \in \mathbb{N}$ , and so  $m^2$  is an even number. If instead  $m$  is odd, then  $m = 2p + 1$  and

$$m^2 = (2p + 1)^2 = 4p^2 + 4p + 1 = 2(2p^2 + 2p) + 1$$

showing that also  $m^2$  is odd.

This justifies Step 7: Indeed we know that  $m^2$  is an even number from Step 6. If  $m$  was odd, then  $m^2$  would be odd. Hence  $m$  must be even as well.

8. If we substitute (1.8) in (1.7) we get

$$m^2 = 2n^2 \implies (2p)^2 = 2n^2 \implies 4p^2 = 2n^2$$

Dividing both terms by 2, we obtain

$$n^2 = 2p^2. \tag{1.9}$$

9. We now make a series of observations:

- Equation (1.9) says that  $n^2$  is even.
- Step 6 says that  $m^2$  is even.
- Therefore  $n$  and  $m$  are also even.
- Hence  $n$  and  $m$  have 2 as common factor.
- But Step 5 says that  $n$  and  $m$  have no common factors.
- **CONTRADICTION**

10. Our reasoning has run into a **contradiction**, starting from assumption (1.4), which says that

$$\sqrt{2} \in \mathbb{Q}.$$

Hence the above must be **FALSE**, and so

$$\sqrt{2} \notin \mathbb{Q}$$

ending the proof.

Seeing that  $\sqrt{2} \notin \mathbb{Q}$ , we might be tempted to just fill in the gap by adding  $\sqrt{2}$  to  $\mathbb{Q}$ . However, with analogous proof to Theorem 1.6, we can prove that

$$\sqrt{p} \notin \mathbb{Q}$$

for each prime number  $p$ . As there are infinite prime numbers, this means that  $\mathbb{Q}$  has infinite gaps. Then we might attempt to fill in these gaps via the extension

$$\tilde{\mathbb{Q}} := \mathbb{Q} \cup \{\sqrt{p} : p \text{ prime}\}.$$

However even this is not enough, as we would still have numbers which are not contained in  $\tilde{\mathbb{Q}}$ , for example

$$\sqrt{2} + \sqrt{3}, \pi, \pi + \sqrt{2} \notin \tilde{\mathbb{Q}}.$$

### Remark 1.7

Proving that

$$\sqrt{2} + \sqrt{3} \notin \mathbb{Q}$$

is relatively easy, and will be left as an **exercise**. Instead, proving that

$$\pi \notin \mathbb{Q}$$

is way more complicated. There are several proof of the fact, all requiring mathematics which is more advanced of the one presented in this course. For some proofs, see this [Wikipedia page](#).

The reality of things is that to **complete**  $\mathbb{Q}$  and make it into a **continuous line** we have to add a lot of points. Indeed, we need to add way more points than the ones already contained in  $\mathbb{Q}$ . Such extension of  $\mathbb{Q}$  will be called  $\mathbb{R}$ , the set of **real numbers**. The inclusions will therefore be

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}.$$

The set  $\mathbb{R}$  is not at all trivial to construct. In fact, at first we will not construct it, but just do the following:

- We will assume that  $\mathbb{R}$  **exists** and satisfies some basic **axioms**.
- One of the axioms is that  $\mathbb{R}$  fills **all** the **gaps** that  $\mathbb{Q}$  has. Therefore  $\mathbb{R}$  can be thought as a **continuous** line.
- We will study the **properties** of  $\mathbb{R}$  which descend from such **axioms**.

For example one of the properties of  $\mathbb{R}$  will be the following:

**Theorem 1.8:** We will prove this in the future

$\mathbb{R}$  contains all the square roots. This means that for every  $x \in \mathbb{R}$  with  $x \geq 0$ , we have

$$\sqrt{x} \in \mathbb{R}.$$

At the end of this chapter we will provide a concrete **model** for the real numbers  $\mathbb{R}$ , to prove once and for all that such set indeed exists.

**Theorem 1.9:** We will prove this in the future

There exists a set  $\mathbb{R}$ , called the set of real numbers, which has the following properties:

- $\mathbb{R}$  extends  $\mathbb{Q}$ , that is,
$$\mathbb{Q} \subset \mathbb{R}.$$
- $\mathbb{R}$  satisfies certain **axioms**.
- $\mathbb{R}$  fills **all** the **gaps** that  $\mathbb{Q}$  has. In particular  $\mathbb{R}$  can be represented by a **continuous** line.

## 2 Preliminaries

Before introducing  $\mathbb{R}$  we want to make sure that we cover all the basics needed for the task.

### 2.1 Sets

A set is a **collection** of objects. These objects are called **elements** of the set. For example in the previous section we mentioned the following sets:

- $\mathbb{N}$  the set of natural numbers
- $\mathbb{Z}$  the set of integers
- $\mathbb{Q}$  the set of rational numbers
- $\mathbb{R}$  the set of real numbers

Given an arbitrary set  $A$ , we write

$$x \in A$$

if the element  $x$  belongs to the set  $A$ . If an element  $x$  is not contained in  $A$ , we say that

$$x \notin A.$$

#### Remark 2.1

A set can contain all sorts of elements. For example the students in a classroom can be modelled by a set  $S$ . The elements of the set are the students. For example

$$S = \{\text{Alice, Olivia, Jake, Sahab}\}$$

In this case we have

$$\text{Alice} \in S$$

but instead

$$\text{Silvio} \notin S.$$

## 2.2 Logic

In this section we introduce some basic logic symbols. Suppose that you are given two statements, say  $\alpha$  and  $\beta$ . The formula

$$\alpha \implies \beta$$

means that  $\alpha$  **implies**  $\beta$ . In other words, if  $\alpha$  is true then also  $\beta$  is true.

The formula

$$\alpha \longleftarrow \beta$$

means that  $\alpha$  is implied by  $\beta$ : if  $\beta$  is true then also  $\alpha$  is true.

When we write

$$\alpha \iff \beta \tag{2.1}$$

we mean that  $\alpha$  and  $\beta$  are **equivalent**. Note that (2.1) is equivalent to

$$\alpha \implies \beta \text{ and } \beta \implies \alpha.$$

Such equivalence is very useful in proofs.

### Example 2.2

We have that

$$x > 0 \implies x > -100,$$

and

$$\text{contradiction} \longleftarrow \sqrt{2} \in \mathbb{Q}.$$

Concerning  $\iff$  we have

$$x^2 < 2 \iff -\sqrt{2} < x < \sqrt{2}.$$

We now introduce logic **quantifiers**. These are

- $\forall$  which reads **for all**
- $\exists$  which reads **exists**
- $\exists!$  which reads **exists unique**
- $\nexists$  which reads **does not exist**

These work in the following way. Suppose that you are given a statement  $\alpha(x)$  which depends on the point  $x \in \mathbb{R}$ . Then we say

- $\alpha(x)$  is satisfied for all  $x \in A$  with  $A$  some collection of numbers. This translates to the symbols

$$\alpha(x) \text{ is true } \forall x \in A,$$

- There exists some  $x$  in  $\mathbb{R}$  such that  $\alpha(x)$  is satisfied: in symbols

$$\exists x \in \mathbb{R} \text{ such that } \alpha(x) \text{ is true,}$$

- There exists a unique  $x_0$  in  $\mathbb{R}$  such that  $\alpha(x)$  is satisfied: in symbols

$$\exists! x_0 \in \mathbb{R} \text{ such that } \alpha(x_0) \text{ is true,}$$

- $\alpha(x)$  is never satisfied:

$$\nexists x \in \mathbb{R} \text{ such that } \alpha(x) \text{ is true.}$$

### Example 2.3

Let us make concrete examples:

- The expression  $x^2$  is always non-negative. Thus we can say

$$x^2 \geq 0 \text{ for all } x \in \mathbb{R}.$$

- The equation  $x^2 = 1$  has two solutions  $x = 1$  and  $x = -1$ . Therefore we can say

$$\exists x \in \mathbb{R} \text{ such that } x^2 = 1.$$

- The equation  $x^3 = 1$  has a unique solution  $x = 1$ . Thus

$$\exists! x \in \mathbb{R} \text{ such that } x^3 = 1.$$

- We know that the equation  $x^2 = 2$  has no solutions in  $\mathbb{Q}$ . Then

$$\nexists x \in \mathbb{Q} \text{ such that } x^2 = 2.$$

## 2.3 Operations on sets

### 2.3.1 Union and intersection

For two sets  $A$  and  $B$  we define their **union** as the set

$$A \cup B := \{x : x \in A \text{ or } x \in B\}.$$

The **intersection** of  $A$  and  $B$  is defined by

$$A \cap B := \{x : x \in A \text{ and } x \in B\}.$$

We denote the **empty set** by the symbol  $\emptyset$ . Two sets are **disjoint** if

$$A \cap B = \emptyset.$$

**Example 2.4**

Define the subset of rational numbers

$$S := \left\{ x \in \mathbb{Q} : 0 < x < \frac{5}{2} \right\}.$$

Then we have

$$\mathbb{N} \cap S = \{1, 2\}.$$

We can also define the sets of **even** and **odd** numbers by

$$E := \{2n : n \in \mathbb{N}\}, \quad (2.2)$$

$$O := \{2n + 1 : n \in \mathbb{N}\}. \quad (2.3)$$

Then we have

$$\mathbb{N} \cap E = E, \quad \mathbb{N} \cap O = O, \quad (2.4)$$

$$O \cup E = \mathbb{N}, \quad O \cap D = \emptyset. \quad (2.5)$$

**2.3.2 Inclusion and equality**

Given two sets  $A$  and  $B$ , we say that  $A$  is **contained** in  $B$  if all the elements of  $A$  are also contained in  $B$ . This will be denoted with the **inclusion** symbol  $\subset$ , that is,

$$A \subset B.$$

In this case we say that

- $A$  is a **subset** of  $B$ ,
- $B$  is a **superset** of  $A$ .

The inclusion  $A \subset B$  is equivalent to the implication:

$$x \in A \implies x \in B$$

for all  $x \in A$ . The symbol  $\implies$  reads **implies**, and denotes the fact that the first condition implies the second.



**Example 2.5**

Given two sets  $A$  and  $B$  we always have

$$(A \cap B) \subset A, (A \cap B) \subset B, \quad (2.6)$$

$$A \subset (A \cup B), B \subset (A \cup B). \quad (2.7)$$

We say that two sets  $A$  and  $B$  are equal if they contain the **same** elements. We denote equality by the symbol

$$A = B.$$

**Example 2.6**

The sets

$$A = \{1, 2, 3\}$$

and

$$B = \{3, 1, 2\}$$

are equal. This is because they contain exactly the same elements: **order** does not matter when talking about sets.

**Proposition 2.7**

Let  $A$  and  $B$  be sets. Then

$$A = B$$

**if and only if**

$$A \subset B \text{ and } B \subset A.$$

**Proof**

The proof is almost trivial. However it is a good exercise in basic logic, so let us do it.

1. First implication  $\implies$  :

Suppose that  $A = B$ . Let us show that  $A \subset B$ . Since  $A = B$ , this means that all the elements of  $A$  are also contained in  $B$ . Therefore if we take  $x \in A$  we have

$$x \in A \implies x \in B.$$

This shows  $A \subset B$ . The proof of  $B \subset A$  is similar.

2. Second implication  $\impliedby$  :

Suppose that  $A \subset B$  and  $B \subset A$ . We need to show  $A = B$ , that is,  $A$  and  $B$  have the same elements. To this end let  $x \in A$ . Since  $A \subset B$  then we have  $x \in B$ . Thus  $B$  contains all the elements of  $A$ .

Since we are also assuming  $B \subset A$ , this means that  $A$  contains all the elements of  $B$ . Hence  $A$  and  $B$  contain the same elements, and  $A = B$ .

The above proposition is very useful when we need to **prove** that two sets are equal: rather than showing directly that  $A = B$ , we can prove that  $A \subset B$  and  $B \subset A$ .

### 2.3.3 Infinite unions and intersections

Suppose given a set  $\Omega$ , and a family of sets  $A_n \subset \Omega$ , where  $n \in \mathbb{N}$ . Then we can define the **infinite union**

$$\bigcup_{n \in \mathbb{N}} A_n := \{x \in \Omega : x \in A_n \text{ for at least one } n \in \mathbb{N}\}.$$

The **infinite intersection** is defined as

$$\bigcap_{n \in \mathbb{N}} A_n := \{x \in \Omega : x \in A_n \text{ for all } n \in \mathbb{N}\}.$$

#### Example 2.8

Let the ambient set be  $\Omega := \mathbb{N}$  and define the family  $A_n$  by

$$A_1 := \{1, 2, 3, 4, \dots\} \quad (2.8)$$

$$A_2 := \{2, 3, 4, 5, \dots\} \quad (2.9)$$

$$A_3 := \{3, 4, 5, 6, \dots\} \quad (2.10)$$

$$\dots \dots \quad (2.11)$$

$$A_n := \{n, n+1, n+2, n+3, \dots\}, \quad (2.12)$$

for arbitrary  $n \in \mathbb{N}$ . Then

$$\bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}. \quad (2.13)$$

The above equality can be easily proven using Proposition 2.7. Indeed, assume that  $m \in \bigcup_{n \in \mathbb{N}} A_n$ . Then  $m \in A_n$  for at least one  $n \in \mathbb{N}$ . Since  $A_n \subset \mathbb{N}$ , we conclude that  $m \in \mathbb{N}$ . This shows

$$\bigcup_{n \in \mathbb{N}} A_n \subset \mathbb{N}.$$

Conversely, suppose that  $m \in \mathbb{N}$ . By definition  $m \in A_m$ . Hence there exists at least one index  $n$ ,  $n = m$  in this case, such that  $m \in A_n$ . Then by definition  $m \in \bigcup_{n \in \mathbb{N}} A_n$ , showing that

$$\mathbb{N} \subset \bigcup_{n \in \mathbb{N}} A_n.$$

Hence we conclude (2.13) by Proposition 2.7.

We also have that

$$\bigcap_{n \in \mathbb{N}} A_n = \emptyset. \quad (2.14)$$

We prove the above by **contradiction**. Indeed, suppose that (2.14) is false, i.e.,

$$\bigcap_{n \in \mathbb{N}} A_n \neq \emptyset.$$

This means there exists some  $m \in \mathbb{N}$  such that  $m \in \bigcap_{n \in \mathbb{N}} A_n$ . Hence, by definition,  $m \in A_n$  for all  $n \in \mathbb{N}$ . However  $m \notin A_{m+1}$ , yielding a contradiction. Thus (2.14) holds.

### 2.3.4 Complement

Suppose that  $A$  and  $B$  are subsets of a larger set  $\Omega$ . The **complement** of  $A$  with respect to  $B$  is the set of elements of  $B$  which do not belong to  $A$ , that is

$$B \setminus A := \{x \in \Omega : x \in B \text{ and } x \notin A\}.$$

In particular, the complement of  $A$  with respect to  $\Omega$  is denoted by

$$A^c := \Omega \setminus A := \{x \in \Omega : x \notin A\}.$$

#### Remark 2.9

Suppose that  $A \subset \Omega$ . Then  $A$  and  $A^c$  form a **partition** of  $\Omega$ , in the sense that

$$A \cup A^c = \Omega \text{ and } A \cap A^c = \emptyset.$$

#### Example 2.10

Suppose  $A, B \subset \Omega$ . Then

$$A \subset B \iff B^c \subset A^c.$$

Let us prove the above claim:

- First implication  $\implies$  :  
Suppose that  $A \subset B$ . We need to show that  $B^c \subset A^c$ . Hence, assume  $x \in B^c$ . By definition this means that  $x \notin B$ . Now notice that we cannot have that  $x \in A$ . Indeed, assume  $x \in A$ . By assumption we have  $A \subset B$ , hence  $x \in B$ . But we had assumed  $x \notin B$ , contradiction. Therefore it must be that  $x \notin A$ . Thus  $B^c \subset A^c$ .

- Second implication  $\Leftarrow$  :  
Essentially the same proof, hence we omit it.

We conclude by stating the De Morgan's Laws. The proof will be left as an exercise.

**Proposition 2.11:** De Morgan's Laws

Suppose  $A, B \subset \Omega$ . Then

$$(A \cap B)^c = A^c \cup B^c$$

and

$$(A \cup B)^c = A^c \cap B^c.$$

### 2.3.5 Power set

Let  $\Omega$  be an arbitrary set. We define the **power set** of  $\Omega$  as

$$\mathcal{P}(\Omega) := \{A : A \subset \Omega\},$$

that is, the power set of  $\Omega$  is the set of all subsets of  $\Omega$ .

**Remark 2.12**

It holds that:

1.  $\mathcal{P}(\Omega)$  is always non-empty, since we have that

$$\emptyset \in \mathcal{P}(\Omega), \quad \Omega \in \mathcal{P}(\Omega).$$

2. Given  $A, B \in \mathcal{P}(\Omega)$ , then the sets

$$A \cup B, \quad A \cap B, \quad A^c, \quad B \setminus A$$

are all elements of  $\mathcal{P}(\Omega)$ .

3. Suppose  $\Omega$  is **discrete** and **finite**, that is,

$$\Omega = \{x_1, \dots, x_m\}$$

for some  $m \in \mathbb{N}$ . Then  $\mathcal{P}(\Omega)$  contains  $2^m$  elements.

This is because for each  $x_i \in \Omega$  we have just two choices: either include  $x_i$  in a subset, or do not include  $x_i$  in a subset.

**Example 2.13**

Define the set

$$\Omega = \{x, y, z\}.$$

Then  $\mathcal{P}(\Omega)$  has  $2^3 = 8$  elements. These are

- $\emptyset$
- $\{x\}$
- $\{y\}$
- $\{z\}$
- $\{x, y\}$
- $\{x, z\}$
- $\{y, z\}$
- $\{x, y, z\}$

We therefore write

$$\mathcal{P}(\Omega) = \{\emptyset, \{x\}, \{y\}, \{z\}, \{x, y\} \tag{2.15}$$

$$\{x, z\}, \{y, z\}, \{x, y, z\}\}. \tag{2.16}$$

**2.3.6 Product of sets**

Suppose  $A$  and  $B$  are two sets. The **product** of  $A$  and  $B$  is the set of pairs

$$A \times B := \{(a, b) : a \in A, b \in B\}.$$

By definition two elements in  $A \times B$  are the same, in symbols

$$(a, b) = (\tilde{a}, \tilde{b})$$

if and only if they are equal component-by-component, that is

$$a = \tilde{a}, \quad b = \tilde{b}.$$

**2.4 Equivalence relation**

Suppose  $A$  is a set. A **binary relation**  $R$  on  $A$  is a subset

$$R \subset A \times A.$$

**Definition 2.14:** Equivalence relation

A binary relation  $R$  is called an **equivalence relation** if it satisfies the following properties:

1. **Reflexive:** For each  $x \in A$  one has

$$(x, x) \in R,$$

This is saying that all the elements in  $A$  must be related to themselves

2. **Symmetric:** We have

$$(x, y) \in R \implies (y, x) \in R$$

If  $x$  is related to  $y$ , then  $y$  is related to  $x$

3. **Transitive:** We have

$$(x, y) \in R, (y, z) \in R \implies (x, z) \in R$$

If  $x$  is related to  $y$ , and  $y$  is related to  $z$ , then  $x$  must be related to  $z$

If  $(x, y) \in R$  we write

$$x \sim y$$

and we say that  $x$  and  $y$  are **equivalent**.

**Definition 2.15:** Equivalence classes

Suppose  $R$  is an **equivalence relation** on  $A$ . The **equivalence class** of an element  $x \in A$  is the set

$$[x] := \{y \in A : y \sim x\}.$$

The set of equivalence classes of elements of  $A$  with respect to the equivalence relation  $R$  is denoted by

$$A/R := \{[x] : x \in A\}.$$

Let us immediately clarify the above definitions by considering the prototypical example of equivalence relation: the **equality**.

**Example 2.16:** Equality is an equivalence relation

Consider the set of natural numbers  $\mathbb{N}$ . The equality defines a **binary relation** on  $\mathbb{N} \times \mathbb{N}$ , via

$$R := \{(x, y) \in \mathbb{N} \times \mathbb{N} : x = y\}.$$

Let us check that  $R$  is an **equivalence relation**:

1. **Reflexive:** It holds, since  $x = x$  for all  $x \in \mathbb{N}$ ,

2. Symmetric: Again  $x = y$  if and only if  $y = x$ ,

3. Transitive: If  $x = y$  and  $y = z$  then  $x = z$ .

The class of equivalence of  $x \in \mathbb{N}$  is given by

$$[x] = \{x\},$$

that is, this relation is quite trivial, given that each element of  $\mathbb{N}$  can only be related to itself. The quotient space is then

$$\mathbb{N}/R = \{[x] : x \in \mathbb{N}\} = \{\{x\} : x \in \mathbb{N}\}.$$

### Example 2.17

Suppose that  $R$  is a binary relation on the set  $\mathbb{Q}$  of rational numbers defined by

$$x \sim y \iff x - y \in \mathbb{Z}.$$

Then  $R$  is an equivalence relation on  $\mathbb{Q}$ . Indeed:

1. Reflexive: Let  $x \in \mathbb{Q}$ . Then  $x - x = 0$  and  $0 \in \mathbb{Z}$ . Thus  $x \sim x$ .
2. Symmetric: If  $x \sim y$  then  $x - y \in \mathbb{Z}$ . But then also

$$-(x - y) = y - x \in \mathbb{Z}$$

and so  $y \sim x$ .

3. Transitive: Suppose  $x \sim y$  and  $y \sim z$ . Then

$$x - y \in \mathbb{Z} \text{ and } y - z \in \mathbb{Z}.$$

Thus we have

$$x - z = (x - y) + (y - z) \in \mathbb{Z}$$

showing that  $x \sim z$ . This shows that  $R$  is an equivalence relation on  $\mathbb{Q}$ .

Now note that

$$y \sim x \iff y - x \in \mathbb{Z}$$

and the above is equivalent to

$$\exists n \in \mathbb{Z} \text{ s.t. } y - x = n$$

which again is equivalent to

$$\exists n \in \mathbb{Z} \text{ s.t. } y = x + n.$$

Therefore all the elements of  $\mathbb{Q}$  related to  $x$  by  $R$  are of the form

$$x + n, \forall n \in \mathbb{Z}.$$

The equivalence classes with respect to  $R$  are then

$$[x] = \{x + n : n \in \mathbb{Z}\}.$$

Each equivalence class has exactly one element in  $[0, 1) \cap \mathbb{Q}$ , meaning that:

$$\forall x \in \mathbb{Q}, \exists! q \in \mathbb{Q} \text{ s.t. } 0 \leq q < 1 \text{ and } q \in [x].$$

Therefore

$$\mathbb{Q}/R = \{[x] : x \in \mathbb{Q}\} = \{q \in \mathbb{Q} : 0 \leq q < 1\}.$$

## 2.5 Order relation

Similarly, we define **order relations**.

### Definition 2.18: Partial order

A binary relation  $R$  on  $A$  is called a **partial order** if it satisfies the following properties:

1. **Reflexive:** For each  $x \in A$  one has

$$(x, x) \in R,$$

2. **Transitive:** We have

$$(x, y) \in R, (y, z) \in R \implies (x, z) \in R$$

3. **Antisymmetric:** We have

$$(x, y) \in R \text{ and } (y, x) \in R \implies x = y$$

This is the only new condition with respect to the definition of equivalence relation, and it replaces symmetry.

### Definition 2.19: Total order

A binary relation  $R$  on  $A$  is called a **total order relation** if it satisfies the following properties:

1. **Partial order:**  $R$  is a partial order on  $A$ .
2. **Total:** For each  $x, y \in A$  we have

$$(x, y) \in R \text{ or } (y, x) \in R.$$

This is saying that all elements in  $A$  are related.



An example of partial order is the operation of **set inclusion**.

**Example 2.20:** Set inclusion is a partial order

Consider an arbitrary non-empty set  $\Omega$  and consider its **power set**

$$\mathcal{P}(\Omega) = \{A : A \subset \Omega\}.$$

The inclusion defines **binary relation** on  $\mathcal{P}(\Omega) \times \mathcal{P}(\Omega)$ , via

$$R := \{(A, B) \in \mathcal{P}(\Omega) \times \mathcal{P}(\Omega) : A \subset B\}.$$

Let us check that  $R$  is an **order relation**:

1. Reflexive: It holds, since  $A \subset A$  for all  $A \in \mathcal{P}(\Omega)$ ,
2. Transitive: If  $A \subset B$  and  $B \subset C$ , then by definition of inclusion  $A \subset C$ .
3. Antisymmetric: If  $A \subset B$  and  $B \subset A$ , then  $A = B$  by Proposition 2.7.

Therefore  $R$  is a **partial order** on  $\mathcal{P}(\Omega)$ . Note that in general  $R$  is **not** a total order. For example if we consider

$$\Omega = \{x, y\}.$$

Thus

$$\mathcal{P}(\Omega) = \{\emptyset, \{x\}, \{y\}, \{x, y\}\}.$$

If we pick  $A = \{x\}$  and  $B = \{y\}$  then  $A \cap B = \emptyset$ , meaning that

$$A \not\subset B, \quad B \not\subset A,$$

showing that  $R$  is not a total order.

A very important example of total order is the **inequality** on  $\mathbb{Q}$ .

**Example 2.21:** Inequality is a total order

Consider the set of rationals  $\mathbb{Q}$ . The usual inequality defines a **binary relation** on  $\mathbb{Q} \times \mathbb{Q}$ , via

$$R := \{(x, y) \in \mathbb{Q} \times \mathbb{Q} : x \leq y\}.$$

Let us check that  $R$  is an **order relation**:

1. Reflexive: It holds, since  $x \leq x$  for all  $x \in \mathbb{Q}$ ,
2. Transitive: If  $x \leq y$  and  $y \leq z$  then  $x \leq z$ .

3. Antisymmetric: If  $x \leq y$  and  $y \leq x$  then  $x = y$ .

Finally, we also have that  $R$  is a **total order** on  $\mathbb{Q}$ , since for all  $x, y \in \mathbb{Q}$  we have

$$x \leq y \text{ or } y \leq x.$$

### Notation 2.22

If  $\Omega$  is a set and  $R$  is a total order on  $\Omega$ , we write

$$(x, y) \in R \iff x \leq y.$$

Therefore the symbol  $\leq$  will always denote a total order relation.

## 2.6 Intervals

In this section we assume to have available the set  $\mathbb{R}$  of **real numbers**, which we recall is an extension of  $\mathbb{Q}$ . We now introduce the concept of **interval**.

### Definition 2.23

Let  $a, b \in \mathbb{R}$  with  $a < b$ . We define the **open interval**  $(a, b)$  as the set

$$(a, b) := \{x \in \mathbb{R} : a < x < b\}.$$

We define the **close interval**  $[a, b]$  as the set

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}.$$

In general we also define the intervals

$$[a, b) := \{x \in \mathbb{R} : a \leq x < b\}, \quad (2.17)$$

$$(a, b] := \{x \in \mathbb{R} : a < x \leq b\}, \quad (2.18)$$

$$(a, \infty) := \{x \in \mathbb{R} : x > a\}, \quad (2.19)$$

$$[a, \infty) := \{x \in \mathbb{R} : x \geq a\}, \quad (2.20)$$

$$(-\infty, b) := \{x \in \mathbb{R} : x < b\}, \quad (2.21)$$

$$(-\infty, b] := \{x \in \mathbb{R} : x \leq b\}. \quad (2.22)$$

Some of the above intervals are depicted in Figure 2.1, Figure 2.2, Figure 2.3, Figure 2.4 below.

Figure 2.1: Interval  $(a, b)$ Figure 2.2: Interval  $[a, b]$ 

## 2.7 Functions

### Definition 2.24: Functions

Let  $A$  and  $B$  be sets. A **function** from  $A$  to  $B$  is a rule which associates **at each** element  $x \in A$  a **single** element  $y \in B$ . Notations:

- We write

$$f : A \rightarrow B$$

to indicate such rule,

- For  $x \in A$ , we denote by

$$y := f(x) \in B$$

the element associated with  $x$  by  $f$ .

- We will often denote the map  $f$  also by

$$x \mapsto f(x).$$

In addition:

- The set  $A$  is called the **domain** of  $f$ ,
- The **range** of  $f$  is the set

$$\{y \in B : y = f(x) \text{ for some } x \in A\} \subset B.$$

### Warning

We want to stress the importance of the first two sentences in Definition 2.24. Assume that  $f : A \rightarrow B$  is a function. Then:

Figure 2.3: Interval  $(a, \infty)$

Figure 2.4: Interval  $(-\infty, b]$ 

- To each element  $x \in A$  we can **only** associate **one** element  $f(x) \in B$ ,
- Every element  $x \in A$  has to be associated to an element  $f(x) \in B$ .

**Example 2.25**

Assume given the two sets

$$A = \{a_1, a_2\}, \quad B = \{b_1, b_2, b_3\}.$$

Let us see a few examples:

- Define  $f : A \rightarrow B$  by setting

$$f(a_1) = b_1, \quad f(a_2) = b_1.$$

In this way  $f$  is a function, with domain  $A$  and range

$$f(A) = \{b_1\} \subset B.$$

- Define  $g : A \rightarrow B$  by setting

$$g(a_1) = b_2, \quad g(a_1) = b_3, \quad g(a_2) = b_3$$

Then  $g$  is **NOT** a function, since the element  $a_1$  has two elements associated.

- Define  $h : A \rightarrow B$  by setting

$$h(a_1) = b_1.$$

Then  $h$  is **NOT** a function, since the element  $a_2$  has no element associated.

**Example 2.26**

Let us make two examples of functions on  $\mathbb{R}$ :

- Define  $f : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f(x) = x^2.$$

Note that the domain of  $f$  is given by  $\mathbb{R}$ , while the range is

$$f(\mathbb{R}) = [0, \infty).$$

- Define  $g : \mathbb{R} \rightarrow \mathbb{R}$  as the logarithm:

$$g(x) = \log(x).$$

This time the domain is  $(0, \infty)$ , while the range is  $g(\mathbb{R}) = \mathbb{R}$ .




Figure 2.5: Schematic picture of the function  $f$







Figure 2.6: Schematic picture of the function  $g$

Figure 2.7: Schematic picture of the function  $h$ Figure 2.8: Plot of function  $f(x) = x^2$

Figure 2.9: Plot of function  $g(x) = \log(x)$ 

## 2.8 Absolute value or Modulus

In this section we assume to have available the set  $\mathbb{R}$  of **real numbers**, which we recall is an extension of  $\mathbb{Q}$ .

### Definition 2.27: Absolute value

For  $x \in \mathbb{R}$  we define its **absolute value** as the quantity

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

### Example 2.28

By definition one has  $|x| = x$  if  $x \geq 0$ . For example

$$|\pi| = \pi, \quad |\sqrt{2}| = \sqrt{2}, \quad |0| = 0.$$

Instead  $|x| = -x$  if  $x < 0$ . For example

$$|-\pi| = \pi, \quad |-\sqrt{2}| = \sqrt{2}, \quad |-10| = 10.$$

Let us also make the following basic remark, whose proof will be left as an exercise.

**Remark 2.29**

For all  $x \in \mathbb{R}$  one has

$$|x| \geq 0.$$

Moreover

$$|x| = 0 \iff x = 0.$$

Another basic remark (proof by exercise).

**Remark 2.30**

For all  $x \in \mathbb{R}$  one has

$$|x| = |-x|.$$

We can use the definition of absolute value to define the **absolute value function**. This is the function

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) := |x|.$$

You might be familiar with the graph associated to  $f$ , as seen below.





Figure 2.10: Plot of the absolute value function  $f(x) = |x|$

It is also useful to understand the absolute value in a geometric way.




**Remark 2.31:** Geometric interpretation of  $|x|$ 

A number  $x \in \mathbb{R}$  can be represented with a point on the real line  $\mathbb{R}$ . The non-negative number  $|x|$  represents the **distance** of  $x$  from the origin 0. Notice that this works for both positive and negative numbers  $x_1$  and  $x_2$  respectively, as shown in Figure 2.11 below.

Figure 2.11: Geometric interpretation of  $|x|$ **Remark 2.32:** Geometric interpretation of  $|x - y|$ 

If  $x, y \in \mathbb{R}$  then the number  $|x - y|$  represents the distance between  $x$  and  $y$  on the real line, as shown in Figure 2.12 below. Note that by Remark 2.30 we have

$$|x - y| = |y - x|.$$

Figure 2.12: Geometric interpretation of  $|x - y|$ 

In the next Lemma we show a fundamental equivalence regarding the absolute value.

**Lemma 2.33**

Let  $x, y \in \mathbb{R}$ . Then

$$|x| \leq y \iff -y \leq x \leq y.$$

The geometric meaning of the above statement is clear: the distance of  $x$  from the origin is less than  $y$ , in formulae

$$|x| \leq y,$$

if and only if  $x$  belongs to the interval  $[-y, y]$ , in formulae

$$-y \leq x \leq y.$$

A sketch of this explanation is seen in Figure 2.13 below.




Figure 2.13: Geometric meaning of Lemma 2.33

**Proof:** Proof of Lemma 2.33

*Step 1: First implication.*

Suppose first that

$$|x| \leq y. \quad (2.23)$$

Recalling that the absolute value is non-negative, from (2.23) we deduce that  $0 \leq |x| \leq y$ . In particular it holds

$$y \geq 0. \quad (2.24)$$

We make separate arguments for the cases  $x \geq 0$  and  $x < 0$ :

- Case 1:  $x \geq 0$ . From (2.23), (2.24) and from  $x \geq 0$  we have

$$-y \leq 0 \leq x = |x| \leq y$$

which shows

$$-y \leq x \leq y.$$

- Case 2:  $x < 0$ . From (2.23), (2.24) and from  $x < 0$  we have

$$-y \leq 0 < -x = |x| \leq y$$

which shows

$$-y \leq -x \leq y.$$

Multiplying the above inequalities by  $-1$  yields

$$-y \leq x \leq y.$$

*Step 2: Second implication.*

Suppose now that

$$-y \leq x \leq y. \quad (2.25)$$

We make separate arguments for the cases  $x \geq 0$  and  $x < 0$ :

- Case 1:  $x \geq 0$ . Since  $x \geq 0$ , from (2.25) we get

$$|x| = x \leq y$$

showing that

$$|x| \leq y.$$

- Case 2:  $x < 0$ . Since  $x < 0$ , from (2.25) we have

$$-y \leq x = -|x|.$$

Multiplying the above inequality by  $-1$  yields

$$|x| \leq y.$$

With the same arguments, just replacing  $\leq$  with  $<$ , one can also show the following.

#### Lemma 2.34

Let  $x, y \in \mathbb{R}$ . Then

$$|x| < y \iff -y < x < y.$$

## 2.9 Triangle inequality

The triangle inequality relates the absolute value to the sum operation. It is a very important inequality, which we will use a lot in the future.

#### Theorem 2.35: Triangle inequality

For every  $x, y \in \mathbb{R}$  we have

$$||x| - |y|| \leq |x + y| \leq |x| + |y|. \quad (2.26)$$

Before proceeding with the proof, let us discuss the geometric meaning of the triangle inequality.

#### Remark 2.36: Geometric meaning of triangle inequality

The notion of absolute value can be extended also to vectors in the plane. Suppose that  $x$  and  $y$  are two vectors in the plane, as in Figure 2.14 below. Then  $|x|$  and  $|y|$  can be interpreted as the **lengths** of these

vectors.

Using the rule of sum of vectors, we can draw  $x + y$ , as shown in Figure 2.15 below. From the picture it is evident that

$$|x + y| \leq |x| + |y|, \quad (2.27)$$

that is, *the length of each side of a triangle does not exceed the sum of the lengths of the two remaining sides*. Note that (2.27) is exactly the second inequality in (2.26). This is why (2.26) is called triangle inequality.



Figure 2.14: Vectors  $x$  and  $y$



Figure 2.15: Summing the vectors  $x$  and  $y$ . The triangle inequality relates the length of  $x + y$  to the length of  $x$  and  $y$

**Proof:** Proof of Theorem 2.32

Assume that  $x, y \in \mathbb{R}$ . We prove the two inequalities in (2.26) individually.

*Step 1. Proof of the second inequality in (2.26).*

Trivially we have

$$|x| \leq |x|.$$

Therefore we can apply Lemma 2.33 and infer

$$-|x| \leq x \leq |x|. \quad (2.28)$$

Similarly we have that  $|y| \leq |y|$ , and so Lemma 2.33 implies

$$-|y| \leq y \leq |y|. \quad (2.29)$$

Summing (2.28) and (2.29) we get

$$-(|x| + |y|) \leq x + y \leq |x| + |y|.$$

We can now again apply Lemma 2.33 to get

$$|x + y| \leq |x| + |y|, \quad (2.30)$$

which is the second inequality in (2.26).

*Step 2. Proof of the first inequality in (2.26).*

Note that the trivial identity

$$x = x + y - y$$

always holds. We then have

$$|x| = |x + y - y| \quad (2.31)$$

$$= |(x + y) + (-y)| \quad (2.32)$$

$$= |a + b| \quad (2.33)$$

with  $a = x + y$  and  $b = -y$ . We can now apply (2.30) to  $a$  and  $b$  to obtain

$$|x| = |a + b| \quad (2.34)$$

$$\leq |a| + |b| \quad (2.35)$$

$$= |x + y| + |-y| \quad (2.36)$$

$$= |x + y| + |y| \quad (2.37)$$

Therefore

$$|x| - |y| \leq |x + y|. \quad (2.38)$$

We can now swap  $x$  and  $y$  in (2.38) to get

$$|y| - |x| \leq |x + y|.$$

By rearranging the above inequality we obtain

$$-|x + y| \leq |x| - |y|. \quad (2.39)$$

Putting together (2.38) and (2.39) yields

$$-|x + y| \leq |x| - |y| \leq |x + y|.$$

By Lemma 2.33 the above is equivalent to

$$||x| - |y|| \leq |x + y|,$$

which is the first inequality in (2.26).

An immediate consequence of the triangle inequality are the following inequalities, which are left as an exercise.

#### Remark 2.37

For any  $x, y \in \mathbb{R}$  it holds

$$||x| - |y|| \leq |x - y| \leq |x| + |y|.$$

Moreover for any  $x, y, z \in \mathbb{R}$  it holds

$$|x - y| \leq |x - z| + |z - y|.$$

## 2.10 Proofs in Mathematics

In a mathematical proof one needs to show that

$$\alpha \implies \beta \quad (2.40)$$

where

- $\alpha$  is a given set of assumptions, or **Hypothesis**
- $\beta$  is a conclusion, or **Thesis**

Proving (2.40) means convincing ourselves that  $\beta$  follows from  $\alpha$ . Common strategies to prove (2.40) are:

1. **Contradiction:** Assume that the thesis is **false**, and hope to reach a contradiction: that is, prove that

$$\neg\beta \implies \text{contradiction}$$

where  $\neg\beta$  is the **negation** of  $\beta$ .

For example we already proved by contradiction that

$$\text{Definition of } \mathbb{Q} \implies \sqrt{2} \notin \mathbb{Q},$$

In the above statement

$$\alpha = (\text{Definition of } \mathbb{Q}) .$$

$$\beta = (\sqrt{2} \notin \mathbb{Q}) .$$

Therefore

$$\neg\beta = (\sqrt{2} \in \mathbb{Q}) .$$

2. **Direct:** Sometimes proofs will also need **direct** arguments, meaning that one need to show directly that (2.40) holds.
3. **Contrapositive:** The statement (2.40) is equivalent to

$$\neg\beta \implies \neg\alpha . \quad (2.41)$$

Thus, instead of proving (2.40), one could show (2.41). The statement (2.41) is called the **contrapositive** of (2.40).

Let us make an example.

### Proposition 2.38

Two real numbers  $a, b$  are equal if and only if for every real number  $\varepsilon > 0$  it follows that  $|a - b| < \varepsilon$ .

Before proceeding with the proof, note that the above statement is just saying that:

Two numbers are equal if and only if they are **arbitrarily** close

By *arbitrarily close* we mean that they are *as close as you want the to be*.

### Proof: of Proposition 2.38

Let us first rephrase the statement using mathematical symbols:

Let  $a, b \in \mathbb{R}$ . Then it holds:

$$a = b \iff |a - b| < \varepsilon, \forall \varepsilon > 0 .$$

Setting

$$\alpha = (a = b) \quad (2.42)$$

$$\beta = (|a - b| < \varepsilon, \forall \varepsilon > 0) \quad (2.43)$$

the statement is equivalent to

$$\alpha \iff \beta.$$

To show the above, it is sufficient to show that

$$\alpha \implies \beta \quad \text{and} \quad \beta \implies \alpha.$$

*Step 1. Proof that  $\alpha \implies \beta$ .*

This proof can be carried out by a **direct** argument. Since we are assuming  $\alpha$ , this means

$$a = b.$$

We want to see that  $\beta$  holds. Therefore fix an arbitrary  $\varepsilon > 0$ . This means that  $\varepsilon$  can be **any** positive number, as long as you fix it. Clearly

$$|a - b| = |0| = 0 < \varepsilon$$

since  $a = b$ ,  $|0| = 0$ , and  $\varepsilon > 0$ . The above shows that

$$|a - b| < \varepsilon.$$

As  $\varepsilon > 0$  was arbitrary, we have just proven that

$$|a - b| < \varepsilon, \quad \forall \varepsilon > 0,$$

meaning that  $\beta$  holds and the proof is concluded.

*Step 2. Proof that  $\beta \implies \alpha$ .*

Let us prove this implication by showing the **contrapositive**

$$\neg\alpha \implies \neg\beta.$$

So let us assume  $\neg\alpha$  is true. This means that

$$a \neq b.$$

We have to see that  $\neg\beta$  holds. But  $\neg\beta$  means that

$$\exists \varepsilon_0 > 0 \text{ s.t. } |a - b| \geq \varepsilon_0.$$

The above is satisfied by choosing

$$\varepsilon_0 := |a - b|,$$

since  $\varepsilon_0 > 0$  given that  $a \neq b$ .



## 2.11 Induction

Another technique for carrying out proofs is **induction**, which we take as an axiom.

### Axiom 2.39: Principle of Induction

Let  $S \subset \mathbb{N}$ . Suppose that

1. We have  $1 \in S$ , and
2. Whenever  $n \in S$ , then  $(n + 1) \in S$ .

Then we have

$$S = \mathbb{N}.$$

### Important

The above is an **axiom**, meaning that we do not prove it, but rather we just **assume it holds**.

### Remark 2.40

It would be possible to prove the Principle of Induction starting from elementary axioms for  $\mathbb{N}$ , called the **Peano Axioms**, see the [Wikipedia page](#).

However, in justifying basic principles of mathematics, one at some point needs to draw a line. This means that something which looks elementary needs to be assumed to hold, in order to have a starting point for proving deeper statements.

In the case of the Principle of Induction, the intuition is clear:

The Principle of Induction is just describing the **domino effect**: *If one tile falls, then the next one will fall as well. Therefore if the first tile falls, all the tiles will fall.*

It seems reasonable to assume such evident principle.

The Principle of Induction can be used to prove statements which depend on some index  $n \in \mathbb{N}$ . Precisely, the following statement holds.

### Corollary 2.41: Principle of Induction - Alternative formulation

Let  $\alpha(n)$  be a statement which depends on  $n \in \mathbb{N}$ . Suppose that

1.  $\alpha(1)$  is true, and

2. Whenever  $\alpha(n)$  is true, then  $\alpha(n + 1)$  is true.

Then  $\alpha(n)$  is true for all  $n \in \mathbb{N}$ .

### Proof

Define the set

$$S := \{n \in \mathbb{N} \text{ s.t. } \alpha(n) \text{ is true}\}.$$

Then

1. We have  $1 \in S$ , since  $\alpha(1)$  is true.
2. If  $n \in S$  then  $\alpha(n)$  is true. By assumption this implies that  $\alpha(n + 1)$  is true. Therefore  $(n + 1) \in S$ .

Therefore  $S$  satisfies the assumptions of the Induction Principle and we conclude that

$$S = \mathbb{N}.$$

By definition this means that  $\alpha(n)$  is true for all  $n \in \mathbb{N}$ .

### Example 2.42: Formula for summing first $n$ natural numbers

Using the Principle of Induction we can prove that

$$1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n + 1)}{2} \quad (2.44)$$

holds for all  $n \in \mathbb{N}$ .

*Proof.* To be really precise, consider the statement

$$\alpha(n) := \text{the above formula is true for } n.$$

In order to apply induction, we need to show that

1.  $\alpha(1)$  is true,
2. If  $\alpha(n)$  is true then  $\alpha(n + 1)$  is true.

Let us proceed:

1. It is immediate to check that (2.44) holds for  $n = 1$ .

2. Suppose (2.44) holds for  $n$ . Then

$$1 + \dots + n + (n + 1) = \frac{n(n + 1)}{2} + (n + 1) \quad (2.45)$$

$$= \frac{n(n + 1) + 2(n + 1)}{2} \quad (2.46)$$

$$= \frac{(n + 1)(n + 2)}{2} \quad (2.47)$$

where in the first equality we used that (2.44) holds for  $n$ . We then have

$$1 + \dots + n + (n + 1) = \frac{(n + 1)(n + 2)}{2},$$

which shows that (2.44) holds for  $n + 1$ .

By the Principle of Induction we then conclude that  $\alpha(n)$  is true for all  $n \in \mathbb{N}$ , which means that (2.44) holds for all  $n \in \mathbb{N}$ .

### Example 2.43: Statements about sequences of numbers

Suppose you are given a collection of numbers

$$\{x_n \text{ s.t. } n \in \mathbb{N}\}.$$

Such collection of numbers is called **sequence**. Assume that

$$x_1 := 1 \quad (2.48)$$

$$x_{n+1} := \frac{x_n}{2} + 1. \quad (2.49)$$

A sequence defined as above is called **recurrence sequence**. Using the above rule we can compute all the terms of  $x_n$ .

For example

$$x_2 = \frac{x_1}{2} + 1 = \frac{1}{2} + 1 = \frac{3}{2} \quad (2.50)$$

$$x_3 = \frac{x_2}{2} + 1 = \frac{3}{4} + 1 = \frac{7}{4}. \quad (2.51)$$

By computing these terms, we suspect that the sequence might be increasing, meaning that

$$x_{n+1} \geq x_n \quad (2.52)$$

for all  $n \in \mathbb{N}$ .

*Claim.* (2.52) holds for all  $n \in \mathbb{N}$ .

*Proof of Claim.*

We argue by induction:

1. We have seen that  $x_1 = 1$  and  $x_2 = 3/2$ . Thus

$$x_2 \geq x_1.$$

2. Suppose now that

$$x_{n+1} \geq x_n. \tag{2.53}$$

We need to prove that

$$x_{n+2} \geq x_{n+1}. \tag{2.54}$$

Indeed, we can multiply the inequality (2.53) by  $1/2$  and add 1 to get

$$\frac{x_{n+1}}{2} + 1 \geq \frac{x_n}{2} + 1.$$

The above is equivalent, by definition, to (2.54).

Therefore the assumptions of the Induction Principle are satisfied, and (2.52) follows.

## 3 Real Numbers

In this chapter we introduce the system of Real Numbers  $\mathbb{R}$  and study some of its properties.

### 3.1 Fields

In order to introduce  $\mathbb{R}$ , we need the concepts of **binary operation** and **field**. We proceed in a general setting, starting from a set  $K$ .

#### Definition 3.1: Binary operation

A binary operation on a set  $K$  is a function

$$\circ : K \times K \rightarrow K$$

which maps the ordered pair  $(x, y)$  into  $x \circ y$ .

#### Notation 3.2

There are two main binary operations we are interested in:

- **Addition:** denoted by  $+$ . The addition, or **sum** of  $x, y \in K$  is denoted by

$$x + y.$$

- **Multiplication:** denoted by  $\cdot$ . The multiplication, or **product** of  $x, y \in K$  is denoted by

$$x \cdot y \text{ or } xy.$$

**Example 3.3:** of binary operation

Let  $K = \{0, 1\}$ . We can for example define operations of sum and product on  $K$  according to the tables

$+$	$0$	$1$
$0$	$0$	$1$
$1$	$1$	$0$

$\cdot$	$0$	$1$
$0$	$0$	$0$
$1$	$0$	$1$

The above mean that

$$0 + 0 = 1 + 1 = 0, \quad 0 + 1 = 1 + 0 = 1,$$

$$0 \cdot 0 = 0 \cdot 1 = 1 \cdot 0 = 0, \quad 1 \cdot 1 = 1.$$

This is just one option. Note that we could not have defined

$$1 + 1 = 2,$$

since  $2 \notin K$ .

Binary operations take ordered pairs of elements of  $K$  as input. Therefore the operation

$$x \circ y \circ z$$

does not make sense, since we do not know which one between

$$x \circ y \quad \text{or} \quad y \circ z$$

has to be performed first. Moreover the outcome of an operation depends on order:

$$x \circ y \neq y \circ x.$$

This motivates the following definition.

**Definition 3.4**

Let  $K$  be a set and  $\circ : K \times K \rightarrow K$  be a binary operation on  $K$ . We say that:

1.  $\circ$  is **commutative** if

$$x \circ y = y \circ x, \quad \forall x, y \in K$$

2.  $\circ$  is **associative** if

$$(x \circ y) \circ z = x \circ (y \circ z), \quad \forall x, y, z \in K$$

3. An element  $e \in K$  is called **neutral element** of  $\circ$  if

$$x \circ e = e \circ x = x, \quad \forall x \in K$$

4. Let  $e$  be a neutral element of  $\circ$  and let  $x \in K$ . An element  $y \in K$  is called an **inverse** of  $x$  with respect to  $\circ$  if

$$x \circ y = y \circ x = e.$$

**Example 3.5**

Let  $K$  with  $+$  and  $\cdot$  be as in Example 3.1. The sum satisfies:

- $+$  is commutative, since

$$0 + 1 = 1 + 0 = 0.$$

- $+$  is associative, since for example

$$(0 + 1) + 1 = 1 + 1 = 0, \quad 0 + (1 + 1) = 0 + 0 = 0,$$

and therefore

$$(0 + 1) + 1 = 0 + (1 + 1).$$

In general one can show that  $+$  is associative by checking all the other permutations.

- The neutral element of  $+$  is 0, since

$$0 + 0 = 0, \quad 1 + 0 = 0 + 1 = 1.$$

- Every element has an inverse. Indeed, the inverse of 0 is 0, since

$$0 + 0 = 0,$$

while the inverse of 1 is 1, since

$$1 + 1 = 1 + 1 = 0.$$

The multiplication satisfies:

- $\cdot$  is commutative, since

$$1 \cdot 0 = 0 \cdot 1 = 0.$$

- $\cdot$  is associative, since for example

$$(0 \cdot 1) \cdot 1 = 0 \cdot 1 = 0, \quad 0 \cdot (1 \cdot 1) = 0 \cdot 1 = 0,$$

and therefore

$$(0 \cdot 1) \cdot 1 = 0 \cdot (1 \cdot 1).$$

By checking all the other permutations one can show that  $\cdot$  is associative.

- The neutral element of  $\cdot$  is 1, since

$$0 \cdot 1 = 1 \cdot 0 = 0, \quad 1 \cdot 1 = 1.$$

- The element 0 has no inverse, since

$$0 \cdot 0 = 0 \cdot 1 = 1 \cdot 0 = 0,$$

and thus we never obtain the neutral element 1. The inverse of 1 is given by 1, since

$$1 \cdot 1 = 1.$$

**Example 3.6**

Let  $K = \{0, 1\}$  be a set with binary relation  $\circ$  defined by the table

$\circ$	0	1
0	1	1
1	0	0

In this case  $\circ$  is not commutative since

$$0 \circ 1 = 1, \quad 1 \circ 0 = 0$$

and therefore

$$0 \circ 1 \neq 1 \circ 0.$$

Moreover  $\circ$  is not associative, since

$$(0 \circ 1) \circ 1 = 1 \circ 1 = 0,$$

while

$$0 \circ (1 \circ 1) = 0 \circ 0 = 1,$$

so that

$$(0 \circ 1) \circ 1 \neq 0 \circ (1 \circ 1).$$

We are ready to define fields.

**Definition 3.7: Field**

Let  $K$  be a set with binary operations of **addition**

$$+ : K \times K \rightarrow K, \quad (x, y) \mapsto x + y$$

and **multiplication**

$$\cdot : K \times K \rightarrow K, \quad (x, y) \mapsto x \cdot y = xy.$$

We call the triple  $(K, +, \cdot)$  a **field** if:

1. The addition  $+$  satisfies:  $\forall x, y, z \in K$

- (A1) **Commutativity and Associativity:**

$$x + y = y + x$$

$$(x + y) + z = x + (y + z)$$

- (A2) **Additive Identity:** There exists a **neutral element** in  $K$  for  $+$ , which we call 0. It holds:

$$x + 0 = 0 + x = x$$



- (A3) **Additive Inverse:** There exists an **inverse** of  $x$  with respect to  $+$ . We call this element the **additive inverse** of  $x$  and denote it by  $-x$ . It holds

$$x + (-x) = (-x) + x = 0$$

2. The multiplication  $\cdot$  satisfies:  $\forall x, y, z \in K$

- (M1) **Commutativity and Associativity:**

$$x \cdot y = y \cdot x$$

$$(x \cdot y) \cdot z = x \cdot (y \cdot z)$$

- (M2) **Multiplicative Identity:** There exists a **neutral element** in  $K$  for  $\cdot$ , which we call 1. It holds:

$$x \cdot 1 = 1 \cdot x = x$$

- (M3) **Multiplicative Inverse:** If  $x \neq 0$  there exists an **inverse** of  $x$  with respect to  $\cdot$ . We call this element the **multiplicative inverse** of  $x$  and denote it by  $x^{-1}$ . It holds

$$x \cdot x^{-1} = x^{-1} \cdot x = 1$$

3. The operations  $+$  and  $\cdot$  are related by

- (AM) **Distributive Property:**  $\forall x, y, z \in K$

$$x \cdot (y + z) = (x \cdot y) + (x \cdot z).$$

### Example 3.8

Let  $K$  with  $+$  and  $\cdot$  be as in Example 3.1. We can show that  $(K, +, \cdot)$  is a field. Indeed we have already shown in Example 3.5 that:

- (A1) and (M1) hold,
- (A2) holds with neutral element 0,
- (M2) holds with neutral element 1,
- (A3) every element has an additive inverse, with

$$-0 = 0, \quad -1 = 1,$$

- (M3) every element which is not 0 a multiplicative inverse, with

$$1^{-1} = 1.$$

We are left to show the Distributive Property (AM). Indeed:

- (AM) For all  $y, z \in K$  we have

$$0 \cdot (y + z) = 0, \quad (0 \cdot y) + (0 \cdot z) = 0 + 0 = 0,$$

and also

$$1 \cdot (y + z) = y + z, \quad (1 \cdot y) + (1 \cdot z) = y + z.$$

Thus (AM) holds.

### Definition 3.9: Subtraction and division

Let  $(K, +, \cdot)$  be a field. We define:

- **Subtraction** as the operation  $-$  defined by

$$x - y := x + (-y), \quad \forall x, y \in K,$$

where  $-y$  is the additive inverse of  $y$ .

- **Division** as the operation  $/$  defined by

$$x/y := x \cdot y^{-1}, \quad \forall x, y \in K, \quad y \neq 0,$$

where  $y^{-1}$  is the multiplicative inverse of  $y$ .

### Proposition 3.10: Uniqueness of neutral elements and inverses

Let  $(K, +, \cdot)$  be a field. Then

1. There is a unique element in  $K$  with the property of 0,
2. There is a unique element in  $K$  with the property of 1,
3. For all  $x \in K$  there is a unique additive inverse  $-x$ ,
4. For all  $x \in K, x \neq 0$ , there is a unique multiplicative inverse  $x^{-1}$ .

### Proof

1. Suppose that  $0 \in K$  and  $\tilde{0} \in K$  are both neutral element of  $+$ , that is, they both satisfy (A2). Then

$$0 + \tilde{0} = 0$$

since  $\tilde{0}$  is a neutral element for  $+$ . Moreover

$$\tilde{0} + 0 = \tilde{0}$$

since  $0$  is a neutral element for  $+$ . By commutativity of  $+$ , see property (A1), we have

$$0 = 0 + \tilde{0} = \tilde{0} + 0 = \tilde{0},$$

showing that  $0 = \tilde{0}$ . Hence the neutral element for  $+$  is unique.

2. Exercise.

3. Let  $x \in K$  and suppose that  $y, \tilde{y} \in K$  are both additive inverses of  $x$ , that is, they both satisfy (A<sub>3</sub>). Therefore

$$x + y = 0$$

since  $y$  is an additive inverse of  $x$  and

$$x + \tilde{y} = 0$$

since  $\tilde{y}$  is an additive inverse of  $x$ . Therefore we can use commutativity and associativity of  $+$ , see property (A<sub>1</sub>), and the fact that  $0$  is the neutral element of  $+$ , to infer

$$y = y + 0 = y + (x + \tilde{y}) \quad (3.1)$$

$$= (y + x) + \tilde{y} = (x + y) + \tilde{y} \quad (3.2)$$

$$= 0 + \tilde{y} = \tilde{y}, \quad (3.3)$$

concluding that  $y = \tilde{y}$ . Thus there is a unique additive inverse of  $x$ , and

$$y = \tilde{y} = -x,$$

with  $-x$  the element from property (A<sub>3</sub>).

4. Exercise.

Using the properties of field we can also show that the usual properties of sum, subtraction, multiplication and division still hold in any field. We list such properties in the following proposition.

**Proposition 3.11:** Properties of field operations

Let  $(K, +, \cdot)$  be a field. Then for all  $x, y, z \in K$ ,

- $x + y = x + z \implies y = z$
- $x \cdot y = x \cdot z$  and  $x \neq 0 \implies y = z$
- $-0 = 0$
- $1^{-1} = 1$
- $x \cdot 0 = 0$
- $-1 \cdot x = -x$
- $-(-x) = x$
- $(x^{-1})^{-1} = x$  if  $x \neq 0$
- $(x \cdot y)^{-1} = x^{-1} \cdot y^{-1}$

The above properties can be all proven with elementary use of the field properties (A<sub>1</sub>)-(A<sub>3</sub>), (M<sub>1</sub>)-(M<sub>3</sub>) and (AM). This is an exercise in patience, and is left to the reader.

Let us conclude with examining the sets of numbers introduced in Chapter 1.

**Theorem 3.12**

Consider the sets  $\mathbb{N}$ ,  $\mathbb{Z}$ ,  $\mathbb{Q}$  with the usual operations  $+$  and  $\cdot$ . We have:

- $(\mathbb{N}, +, \cdot)$  is **not a field**:  
It satisfies properties  $(A_1)$ ,  $(A_2)$ ,  $(M_1)$ ,  $(M_2)$ ,  $(AM)$  of fields. It is missing properties  $(A_3)$  and  $(M_3)$ , the additive and multiplicative inverse properties, respectively.
- $(\mathbb{Z}, +, \cdot)$  is **not a field**:  
It satisfies properties  $(A_1)$ ,  $(A_2)$ ,  $(A_3)$ ,  $(M_1)$ ,  $(M_2)$ ,  $(AM)$  of fields. Thus it is only missing  $(M_3)$ , the multiplicative inverse property.
- $(\mathbb{Q}, +, \cdot)$  is **a field**.

The proof is omitted.

## 3.2 Ordered fields

**Definition 3.13**

Let  $K$  be a set with binary operations  $+$  and  $\cdot$ , and with an order relation  $\leq$ . We call  $(K, +, \cdot, \leq)$  an **ordered field** if:

1.  $(K, +, \cdot)$  is a field
2. There  $\leq$  is of **total order** on  $K$ :  $\forall x, y, z \in K$

- (O<sub>1</sub>) **Reflexivity**:

$$x \leq x$$

- (O<sub>2</sub>) **Antisymmetry**:

$$x \leq y \text{ and } y \leq x \implies x = y$$

- (O<sub>3</sub>) **Transitivity**:

$$x \leq y \text{ and } y \leq z \implies x \leq z$$

- (O<sub>4</sub>) **Total order**:

$$x \leq y \text{ or } y \leq x$$

3. The operations  $+$  and  $\cdot$ , and the total order  $\leq$ , are related by the following properties:  $\forall x, y, z \in K$

- (AM) **Distributive**: Relates addition and multiplication via

$$x \cdot (y + z) = x \cdot y + x \cdot z$$

- (AO) Relates addition and order with the requirement:

$$x \leq y \implies x + z \leq y + z$$

- (MO) Relates multiplication and order with the requirement:

$$x \geq 0, y \geq 0 \implies x \cdot y \geq 0$$

**Example 3.14**

$(\mathbb{Q}, +, \cdot, \leq)$  is an **ordered field**.

### 3.3 Cut Property

We have just introduced the notion of **field**, and noted that the set of rational numbers with the usual operations

$$(\mathbb{Q}, +, \cdot)$$

is a **field**.

We now need to address the key issue we proved in Chapter 1, that is, that

$$\sqrt{2} \notin \mathbb{Q}.$$

This means that  $\mathbb{Q}$  has **gaps**, and cannot be represented as a **continuous** line. The rigorous definition of **lack of gaps** needs the concept of **cut** of a set.

**Definition 3.15:** Partition of a set

Let  $S$  be a non-empty set. The pair  $(A, B)$  is a **partition** of  $S$  if

$$A, B \subset S, \quad A \neq \emptyset, \quad B \neq \emptyset,$$

and

$$S = A \cup B, \quad A \cap B = \emptyset.$$




Figure 3.1: Schematic picture of a partition  $(A, B)$  of the set  $K$ .

**Definition 3.16:** Cut of a set

Let  $S$  be a non-empty set with a total order relation  $\leq$ . The pair  $(A, B)$  is a **cut** of  $S$  if

1.  $(A, B)$  is a **partition** of  $S$ ,
2. We have

$$a \leq b, \quad \forall a \in A, \forall b \in B.$$

The **cut** of a set is often called **Dedekind cut**, named after **Richard Dedekind**, who used cuts to give an explicit construction of the real numbers  $\mathbb{R}$ , see [Wikipedia page](#).

**Definition 3.17:** Cut property

Let  $S$  be a non-empty set with a total order relation  $\leq$ . We say that  $S$  has the **cut property** if for every cut  $(A, B)$  of  $S$  there exists some  $s \in S$  such that

$$a \leq s \leq b, \quad \forall a \in A, \forall b \in B.$$

We call  $s$  the **separator** of the cut  $(A, B)$ .

**Example 3.18**

Let  $S = \mathbb{Q}$  and consider the sets

$$A = (-\infty, s] \cap \mathbb{Q}, \quad B = (s, \infty) \cap \mathbb{Q}.$$

for some  $s \in \mathbb{Q}$ . Then the pair  $(A, B)$  is a cut of  $\mathbb{Q}$ , and  $s$  is the separator.




Figure 3.2:  $(A, B)$  is a cut of  $\mathbb{Q}$  with separator  $s$ .

**Question 3.19**

Do all ordered fields have the Cut Property? Does  $\mathbb{Q}$  have the Cut Property?

The answer to the above question is **NO**. For example the pair

$$A = (-\infty, \sqrt{2}) \cap \mathbb{Q}, \quad B = (\sqrt{2}, \infty) \cap \mathbb{Q}. \quad (3.4)$$

is a cut of  $\mathbb{Q}$ , since  $\sqrt{2} \notin \mathbb{Q}$ . However what is the separator? It should be  $s = \sqrt{2}$ , given that clearly

$$a \leq \sqrt{2} \leq b, \quad \forall a \in A, \quad \forall b \in B.$$

However  $\sqrt{2} \notin \mathbb{Q}$ , so we are **NOT ALLOWED** to take it as separator. Indeed, we can show that  $(A, B)$  defined as in (3.4) has no separator.




Figure 3.3:  $(A, B)$  is a cut of  $\mathbb{Q}$  which has no separator

**Theorem 3.20:**  $\mathbb{Q}$  does not have the cut property.

$\mathbb{Q}$  does not have the cut property. More explicitly, there exist a cut  $(A, B)$  of  $\mathbb{Q}$  which has no separator.

**Remark 3.21**

Before proceeding with the proof, let us summarize the ideas behind it:

We will consider the cut  $(A, B)$  in (3.4). We then assume by contradiction that  $(A, B)$  admits a separator  $L \in \mathbb{Q}$ , so that

$$a \leq L \leq b, \quad \forall a \in A, \forall b \in B. \quad (3.5)$$

Since  $(A, B)$  is a partition of  $\mathbb{Q}$ , then either  $L \in A$  or  $L \in B$ . These will both lead to a contradiction:

- If  $L \in A$ , then by definition we will have

$$L < \sqrt{2}.$$

Now the idea is that  $\sqrt{2} \notin \mathbb{Q}$ , and hence we can always find some  $\tilde{L} \in \mathbb{Q}$  such that

$$L < \tilde{L} < \sqrt{2}. \quad (3.6)$$

Therefore  $\tilde{L} \in A$ , and (3.5) implies

$$\tilde{L} \leq L,$$

which contradicts (3.6).

- If  $L \in B$ , then by definition we will have

$$\sqrt{2} < L.$$

Similarly, as  $\sqrt{2} \notin \mathbb{Q}$ , we can find  $\tilde{L} \in \mathbb{Q}$  such that

$$\sqrt{2} < \tilde{L} < L.$$

Arguing as above we find a contradiction.

Therefore both  $L \in A$  and  $L \in B$  lead to a contradiction. We conclude that  $(A, B)$  has no separator in  $\mathbb{Q}$ .

Time to make the ideas in the above remark rigorous.

**Proof:** Proof of Theorem 3.20

Let  $A$  and  $B$  be the sets defined in (3.4). It is useful to rewrite  $A$  and  $B$  in the form

$$A = A_1 \cup A_2,$$



where

$$A_1 = \{q \in \mathbb{Q} : q < 0\},$$

$$A_2 = \{q \in \mathbb{Q} : q \geq 0, q^2 < 2\},$$

and

$$B = \{q \in \mathbb{Q} : q > 0, q^2 > 2\}.$$

*Step 1.  $(A, B)$  is a cut of  $\mathbb{Q}$ :*

We need to prove the following:

1.  $(A, B)$  is a partition of  $\mathbb{Q}$ . This is because  $A, B \subset \mathbb{Q}$  with  $A \neq \emptyset$  and  $B \neq \emptyset$ . Moreover  $A \cap B = \emptyset$  and

$$A \cup B = \mathbb{Q},$$

given that  $\sqrt{2} \notin \mathbb{Q}$ , and so there is no element  $q \in \mathbb{Q}$  such that  $q^2 = 2$ .

2. It holds

$$a \leq b, \quad \forall a \in A, \forall b \in B.$$

Indeed, suppose that  $a \in A$  and  $b \in B$ . We have two cases:

- $a \in A_1$ : Therefore  $a < 0$ . In particular

$$a < 0 < b,$$

given that  $b > 0$  for all  $b \in B$ . Thus  $a < b$ .

- $a \in A_2$ : Therefore  $a \geq 0$  and  $a^2 < 2$ . In particular

$$a^2 < 2 < b^2,$$

since  $b^2 > 2$  for all  $b \in B$ . In particular

$$a^2 < b^2.$$

Since  $b > 0$  for all  $b \in B$ , from the above inequality we infer  $a < b$ , concluding.

*Step 2.  $(A, B)$  has no separator:*

Suppose by contradiction that  $(A, B)$  admits a separator

$$L \in \mathbb{Q}.$$

By definition this means

$$a \leq L \leq b, \quad \forall a \in A, \forall b \in B. \quad (3.7)$$

Since

$$L \in \mathbb{Q}, \quad \mathbb{Q} = A \cup B, \quad A \cap B = \emptyset,$$

then either  $L \in A$  or  $L \in B$ . We will see that both these possibilities lead to a contradiction:

Case 1:  $L \in A$ .

By (3.7) we know that

$$a \leq L, \quad \forall a \in A. \quad (3.8)$$

In particular the above implies

$$L \geq 0 \quad (3.9)$$

since  $0 \in A$ . Therefore we must have  $L \in A_2$ , that is,

$$L \geq 0 \quad \text{and} \quad L^2 < 2. \quad (3.10)$$

Set

$$\tilde{L} := L + \frac{1}{n}$$

for  $n \in \mathbb{N}$ ,  $n \neq 0$  to be chosen later. Clearly we have

$$\tilde{L} \in \mathbb{Q} \quad \text{and} \quad L < \tilde{L}. \quad (3.11)$$

From (3.9) and (3.11) we have also

$$\tilde{L} > 0. \quad (3.12)$$

We now want to show that there is a choice of  $n$  such that  $\tilde{L}^2 < 2$ , which will lead to a contradiction. Indeed, we can estimate

$$\tilde{L}^2 = \left(L + \frac{1}{n}\right)^2 \quad (3.13)$$

$$= L^2 + \frac{1}{n^2} + 2\frac{L}{n} \quad (3.14)$$

$$< L^2 + \frac{1}{n} + 2\frac{L}{n} \quad \left(\text{using } \frac{1}{n} < \frac{1}{n^2}\right) \quad (3.15)$$

$$= L^2 + \frac{2L+1}{n}. \quad (3.16)$$

If we now impose that

$$L^2 + \frac{2L+1}{n} < 2,$$

we can rearrange the above and obtain

$$n(2 - L^2) > 2L + 1.$$

Now note that  $L^2 < 2$  by assumption (3.10). Thus we can divided by  $(2 - L^2)$  and obtain

$$n > \frac{2L+1}{2-L^2}.$$

Therefore we have just shown that

$$n > \frac{2L+1}{2-L^2} \implies \tilde{L}^2 < 2.$$

Together with (3.12) this implies  $\tilde{L} \in A$ . Therefore we have

$$\tilde{L} \leq L$$

by (3.8). On the other hand it also holds

$$\tilde{L} > L$$

by (3.11), and therefore we have a contradiction. Thus  $L \notin A$ .

Case 2:  $L \in B$ .

As  $L \in B$ , we have by definition

$$L > 0, \quad L^2 > 2. \quad (3.17)$$

Moreover since  $L$  is a separator, see (3.7), in particular

$$L \leq b, \quad \forall b \in B. \quad (3.18)$$

Define now

$$\tilde{L} := L - \frac{1}{n}$$

with  $n \in \mathbb{N}$ ,  $n \neq 0$  to be chosen later. Clearly we have

$$\tilde{L} \in \mathbb{Q}, \quad \tilde{L} < L. \quad (3.19)$$

We now show that  $n$  can be chosen so that  $\tilde{L} \in B$ . Indeed

$$\tilde{L}^2 = \left(L - \frac{1}{n}\right)^2 \quad (3.20)$$

$$= L^2 + \frac{1}{n^2} - 2\frac{L}{n} \quad (3.21)$$

$$> L^2 - \frac{1}{n^2} - 2\frac{L}{n} \quad \left(\text{using } \frac{1}{n^2} > -\frac{1}{n^2}\right) \quad (3.22)$$

$$> L^2 - \frac{1}{n} - 2\frac{L}{n} \quad \left(\text{using } -\frac{1}{n^2} > -\frac{1}{n}\right) \quad (3.23)$$

$$= L^2 - \frac{1+2L}{n}. \quad (3.24)$$

Now we impose

$$L^2 - \frac{1+2L}{n} > 2$$

which is equivalent to

$$n(L^2 - 2) > 1 + 2L.$$

Since we are assuming  $L \in B$ , then  $L^2 > 2$ , see (3.17). Therefore we can divide by  $(L^2 - 2)$  and get

$$n > \frac{1+2L}{L^2 - 2}.$$

In total, we have just shown that

$$n > \frac{1 + 2L}{L^2 - 2} \implies \tilde{L}^2 > 2,$$

proving that  $\tilde{L} \in B$ . Therefore by (3.18) we get

$$L \leq \tilde{L}.$$

This contradicts (3.19).

*Conclusion:*

We have seen that assuming that  $(A, B)$  has a separator  $L \in \mathbb{Q}$  leads to a contradiction. Thus the cut  $(A, B)$  has no separator.

### Remark 3.22

The above proof can be summarized by saying that the set

$$A = (-\infty, \sqrt{2}) \cap \mathbb{Q}$$

does not admit a **largest** element in  $\mathbb{Q}$ , and that the set

$$B = (\sqrt{2}, \infty) \cap \mathbb{Q}$$

does not admit a **lowest element** in  $\mathbb{Q}$ . We will clarify this remark in the next section.

## 3.4 Supremum and infimum

A crucial definition in Analysis is the one of supremum or infimum of a set. This is also another way of studying the **gaps** of  $\mathbb{Q}$ .

### Example 3.23: Intuition about supremum and infimum

Consider the set

$$A = [0, 1) \cap \mathbb{Q}.$$

Intuitively, we understand that  $A$  is bounded, i.e. not infinite. We also see that

- 0 is the lowest element of  $A$
- 1 is the highest element of  $A$

However we see that  $0 \in A$  while  $1 \notin A$ . We will see that

- 0 can be defined as the **infimum** and **minimum** of  $A$ .
- 1 can be defined as the **supremum**, but not **maximum**, of  $A$ .

### 3.4.1 Upper bound, supremum, maximum

We start by defining the supremum. First we need the notion of upper bound of a set.

#### Definition 3.24: Upper bound and bounded above

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ :

1. We say that  $b \in K$  is an **upper bound** for  $A$  if

$$a \leq b, \quad \forall a \in A.$$

2. We say that  $A$  is **bounded above** if there exists an upper bound  $b \in K$  for  $A$ .

#### Definition 3.25: Supremum

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ . A number  $s \in K$  is called **least upper bound** or **supremum** of  $A$  if:

- i.  $s$  is an upper bound for  $A$ ,
- ii.  $s$  is the smallest upper bound of  $A$ , that is,

$$\text{If } b \in K \text{ is upper bound for } A \text{ then } s \leq b.$$

#### Notation 3.26

We will almost always prefer the name supremum to least upper bound. For  $A \subset K$  the supremum is denoted by

$$s := \sup A.$$

#### Remark 3.27

Note that if a set  $A \subset K$  is **NOT** bounded above, then the supremum does not exist, as there are no upper bounds of  $A$ .

**Proposition 3.28**

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ . If

$$\sup A$$

exists, then it is unique.

**Proof**

Suppose there exist  $s_1, s_2 \in K$  such that

$$s_1 = \sup A, \quad s_2 = \sup A.$$

Then:

- Since  $s_2 = \sup A$ , in particular  $s_2$  is an upper bound for  $A$ . Since  $s_1 = \sup A$  then  $s_1$  is the lowest upper bound. Thus we get

$$s_1 \leq s_2.$$

- Exchanging the roles  $s_1$  and  $s_2$  in the above reasoning we also get

$$s_2 \leq s_1.$$

This shows  $s_1 = s_2$ .

**Warning**

In general:

- A set can have infinite upper bounds,
- The supremum does not belong to the set.

For example

$$A = [0, 1) \cap \mathbb{Q}$$

has for upper bounds all the numbers  $b \in \mathbb{Q}$  with  $b > 1$ . Moreover one can show that

$$\sup A = 1,$$

and so

$$\sup A \notin A.$$

**Warning**

The supremum does not exist in general. For example let

$$A = [0, \sqrt{2}) \cap \mathbb{Q}.$$

We will show that  $\sup A$  does not exist in  $\mathbb{Q}$ . Indeed we will have that

$$\sup A = \sqrt{2} \in \mathbb{R}.$$




Figure 3.4: Supremum and upper bounds of a set  $A$  in the field  $K$

**Definition 3.29: Maximum**

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ . A number  $M \in K$  is called the **maximum** of  $A$  if:

$$M \in A \text{ and } a \leq M, \forall a \in A.$$

We denote the maximum by

$$M = \max A.$$

**Proposition 3.30**

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ . If the maximum of  $A$  exists, then also the supremum exists, and

$$\sup A = \max A.$$

**Proof**

Let

$$M = \max A.$$

Then:

- By definition we have  $M \in A$  and

$$a \leq M, \quad \forall a \in A.$$

In particular the above tells us that  $M$  is an upper bound of  $A$ .

- We claim that  $M$  is the least upper bound. Indeed, suppose  $b$  is an upper bound of  $A$ , that is,

$$a \leq b, \quad \forall a \in A.$$

In particular, since  $M \in A$ , by the above condition we have

$$M \leq b.$$

Therefore  $M$  is the least upper bound of  $A$ , meaning that  $M = \sup A$ .

### Warning

The converse of the above statement is not true: In general the sup might exist while the max does not. For example

$$A = [0, 1) \cap \mathbb{Q}$$

is such that

$$\sup A = 1$$

but  $\max A$  does not exist. Instead for the set

$$B = [0, 1] \cap \mathbb{Q}$$

we have that

$$\max A = \sup A = 1.$$

### 3.4.2 Lower bound, infimum, minimum

We now introduce the definitions of lower bound, infimum, minimum. These are the counterpart of upper bound, supremum and maximum, respectively.

#### Definition 3.31: Upper bound, bounded below, infimum, minimum

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ :

1. We say that  $l \in K$  is a **lower bound** for  $A$  if

$$l \leq a, \quad \forall a \in A.$$

2. We say that  $A$  is **bounded below** if there exists a lower bound  $l \in K$  for  $A$ .



3. We say that  $i \in K$  is the **greatest lower bound** or **infimum** of  $A$  if:

- $i$  is a lower bound for  $A$ ,
- $i$  is the largest lower bound of  $A$ , that is,

If  $l \in K$  is a lower bound for  $A$  then  $l \leq i$ .

If it exists, the infimum is denoted by

$$i = \inf A.$$

4. We say that  $m \in K$  is the **minimum** of  $A$  if:

$$m \in A \text{ and } m \leq a, \forall a \in A.$$

If it exists, we denote the minimum by

$$m = \min A.$$




Figure 3.5: Infimum and lower bounds of a set  $A$  in the field  $K$

### Proposition 3.32

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ :

1. If  $\inf A$  exists, then it is unique.
2. If the minimum of  $A$  exists, then also the infimum exists, and

$$\inf A = \min A.$$

The proof uses similar arguments to the one employed in the previous section, and is left to the reader as an exercise.

**Warning**

We have

- A set can have infinite lower bounds,
- The infimum does not belong to the set.

For example

$$A = (0, 1) \cap \mathbb{Q}$$

has for lower bounds all the numbers  $b \in \mathbb{Q}$  with  $b < 1$ . Moreover we will show that

$$\inf A = 0,$$

and so

$$\inf A \notin A.$$

**Warning**

The infimum does not exist in general. For example let

$$A = (\sqrt{2}, 5] \cap \mathbb{Q}.$$

We will show that  $\inf A$  does not exist in  $\mathbb{Q}$ . Indeed we will have that

$$\inf A = \sqrt{2} \in \mathbb{R}.$$

**Warning**

In general the inf might exist while the min does not. For example

$$A = (0, 1) \cap \mathbb{Q}$$

is such that

$$\inf A = 0$$

but  $\min A$  does not exist. Instead for the set

$$B = [0, 1] \cap \mathbb{Q}$$

we have that

$$\inf A = \min A = 0.$$

**Proposition 3.33**

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ . If  $\inf A$  and  $\sup A$  exist, then

$$\inf A \leq a \leq \sup A, \quad \forall a \in A.$$

The proof is simple, and is left as an exercise. We now have a complete picture about supremum and infimum, see figure below.




Figure 3.6: Supremum, upper bounds, infimum and lower bounds of a set  $A$  in  $K$

We conclude with another simple proposition. The proof is again left to the reader.

**Proposition 3.34: Relationship between sup and inf**

Let  $(K, +, \cdot, \leq)$  be an ordered field and  $A \subset K$ . Define

$$-A := \{-a : a \in A\}.$$

It holds:

- If  $\sup A$  exists, then  $\inf A$  exists and

$$\inf(-A) = -\sup A.$$

- If  $\inf A$  exists, then  $\sup A$  exists and

$$\sup(-A) = -\inf A.$$

## 3.5 Completeness

We have introduced the concepts of supremum and infimum on an ordered field  $K$ .

**Question 3.35**

Suppose  $(K, +, \cdot, \leq)$  is an ordered field, and that  $A \subset K$  is non-empty and bounded above. Does

$$\sup A$$

always exist?

The answer to the above question is **NO**. Like we did with the Cut Property, the counterexample can be found in the set of rational numbers  $\mathbb{Q}$ . A set bounded above for which the supremum does not exist is, for example,

$$A = [0, \sqrt{2}) \cap \mathbb{Q}. \quad (3.25)$$

**Theorem 3.36**

There exists a set  $A \subset \mathbb{Q}$  such that

- $A$  is non-empty,
- $A$  is bounded above,
- $\sup A$  does not exist in  $\mathbb{Q}$ .

The proof uses the same ideas we used for showing that  $\mathbb{Q}$  does not have the Cut Property.

**Proof**

Define the set  $A$  as in (3.25). Equivalently, this can be written as

$$A = \{q \in \mathbb{Q} : q \geq 0, q^2 < 2\}.$$

*Step 1.  $A$  is bounded above.*

Take  $b := 9$ . Then  $b$  is an upper bound for  $A$ . Indeed by definition

$$q^2 < 2, q \geq 0, \forall q \in A.$$

Therefore

$$q^2 < 2 < 9 \implies q^2 < 9 \implies q < 3 = b.$$

*Step 2.  $\sup A$  does not exist.*

Assume by contradiction that

$$s = \sup A \in \mathbb{Q}$$

exists. By definition it holds

$$s \geq q, \quad \forall q \in A \quad (3.26)$$

$$b \geq q, \forall q \in A \implies s \leq b \quad (3.27)$$

There are two possibilities:  $s \in A$  or  $s \notin A$ :

- *Case 1.*  $s \in A$ .

If  $s \in A$  by definition

$$s \geq 0, \quad s^2 < 2. \quad (3.28)$$

Define

$$\tilde{s} := s + \frac{1}{n}$$

with  $n \in \mathbb{N}$ ,  $n \neq 0$  to be chosen later. Then

$$\tilde{s}^2 = \left(s + \frac{1}{n}\right)^2 \quad (3.29)$$

$$= s^2 + \frac{1}{n^2} + 2\frac{s}{n} \quad (3.30)$$

$$< s^2 + \frac{1}{n} + 2\frac{s}{n} \quad \left(\text{using } \frac{1}{n} < \frac{1}{n^2}\right) \quad (3.31)$$

$$= s^2 + \frac{2s+1}{n}. \quad (3.32)$$

If we now impose that

$$s^2 + \frac{2s+1}{n} < 2,$$

we can rearrange the above and obtain

$$n(2 - s^2) > 2s + 1.$$

Now note that  $s^2 < 2$  by assumption (3.28). Thus we can divide by  $(2 - s^2)$  and obtain

$$n > \frac{2s+1}{2-s^2}.$$

To summarize, we have just shown that

$$n > \frac{2s+1}{2-s^2} \implies \tilde{s}^2 < 2.$$

Moreover  $\tilde{s} := (s + 1/n) \in \mathbb{Q}$ . Therefore

$$\tilde{s} \in A.$$

Since  $s = \sup A$ , we then have

$$\tilde{s} \leq s.$$

However

$$\tilde{s} := s + \frac{1}{n} > s,$$

yielding a contradiction. Thus  $s \in A$  is not possible.

- *Case 2.*  $s \notin A$ .

If  $s \notin A$ , by the fact that  $s = \sup A$  and by definition of  $A$  we get

$$s > 0, \quad s^2 > 2. \quad (3.33)$$

Define

$$\tilde{s} := s - \frac{1}{n}.$$

We have

$$\tilde{s}^2 = \left(s - \frac{1}{n}\right)^2 \quad (3.34)$$

$$= s^2 + \frac{1}{n^2} - 2\frac{s}{n} \quad (3.35)$$

$$> s^2 - \frac{1}{n^2} - 2\frac{s}{n} \quad \left(\text{using } \frac{1}{n^2} > -\frac{1}{n^2}\right) \quad (3.36)$$

$$> s^2 - \frac{1}{n} - 2\frac{s}{n} \quad \left(\text{using } -\frac{1}{n^2} > -\frac{1}{n}\right) \quad (3.37)$$

$$= s^2 - \frac{1+2s}{n}. \quad (3.38)$$

Now we impose

$$s^2 - \frac{1+2s}{n} > 2$$

which is equivalent to

$$n(s^2 - 2) > 1 + 2s.$$

By (3.33) we have  $s^2 > 2$ . Therefore we can divide by  $(s^2 - 2)$  and get

$$n > \frac{1+2s}{s^2-2}.$$

In total, we have just shown that

$$n > \frac{1+2s}{s^2-2} \implies \tilde{s}^2 > 2.$$

Therefore  $\tilde{s} \notin A$ , and by definition of  $A$  we have

$$\tilde{s} \geq q, \quad \forall q \in A.$$

Moreover  $\tilde{s} := (s - 1/n) \in \mathbb{Q}$ . Therefore  $\tilde{s}$  is an upper bound of  $A$  in  $\mathbb{Q}$ . Since  $s = \sup A$  is the smallest upper bound, see (3.27), it follows

$$s \leq \tilde{s}.$$

However

$$\tilde{s} := s - \frac{1}{n} < s,$$

obtaining a contradiction. Then  $s \notin A$ .

*Conclusion.*

We have assumed by contradiction that  $s = \sup A$  exists in  $\mathbb{Q}$ . In this case either  $s \in A$  or  $s \notin A$ . In both cases we found a contradiction. Therefore  $\sup A$  does not exist.

The above theorem shows that the supremum does not necessarily exist. What about the infimum?

**Question 3.37**

Suppose  $(K, +, \cdot, \leq)$  is an ordered field, and that  $A \subset K$  is non-empty and bounded below. Does

$$\inf A$$

always exist?

The answer to the above question is again **NO**. A set bounded below for which the infimum does not exist is, for example,

$$A = (\sqrt{2}, 10] \cap \mathbb{Q}.$$

The proof of this fact is, of course, very similar to the one of Theorem 3.36, and is therefore omitted.

Thus infimum and supremum do not exist in general. The fields for which all the bounded sets admit supremum or infimum are called **complete**.

**Definition 3.38: Completeness**

Let  $(K, +, \cdot, \leq)$  be an ordered field. We say that  $K$  is **complete** if it holds the property:

- (AC) For every  $A \subset K$  non-empty and bounded above

$$\sup A \in K.$$

**Notation 3.39**

We have that:

- Property (AC) is called **Axiom of Completeness**
- If  $K$  is an ordered field in which (AC) holds, then  $K$  is called a **complete ordered field**

Notice that if the **Axiom of Completeness** holds, then also the infimum exists. This is shown in the following proposition.

**Proposition 3.40**

Let  $(K, +, \cdot, \leq)$  be a complete ordered field. Suppose that  $A \subset K$  is non-empty and bounded below. Then

$$\inf A \in K.$$

**Proof**

Suppose that  $A \subset K$  is non-empty and bounded below. Then

$$-A := \{-a : a \in A\}$$

is non-empty and bounded above. By completeness we have that  $\sup(-A)$  exists in  $K$ . But then Proposition 3.34 implies that  $\inf A$  exists in  $K$ , with

$$\inf A = -\sup(-A).$$

## 3.6 Equivalence of Completeness and Cut Property

We can show that **Completeness** is equivalent to the **Cut Property**. Such result is not essential, but its proof is very instructive.

**Theorem 3.41:** Equivalence of Cut Property and Completeness

Let  $(K, +, \cdot, \leq)$  be an ordered field. Then they are equivalent:

1.  $K$  has the **Cut Property**
2.  $K$  is **Complete**

**Remark 3.42:** Ideas for proving Theorem 3.41

The proof of Theorem 3.41 is rather long, but the ideas are simple:

*Step 1. Cut Property  $\implies$  Completeness.* Suppose  $K$  has the Cut Property. To prove that  $K$  is Complete, we need to:

- Consider an arbitrary set  $A \subset K$  such that  $A \neq \emptyset$  and  $A$  is bounded above.
- Show that  $A$  has a supremum.

To achieve this, consider the set

$$B := \{b \in K : b \geq a, \forall a \in A\},$$



which is the set of Upper Bounds of  $A$ . We can show that the pair

$$(B^c, B)$$

is a Cut of  $K$ . As  $K$  has the Cut Property, then there exists  $s \in K$  separator of  $(B^c, B)$ . We will show that the separator  $s$  is the supremum of  $A$

$$s = \sup A.$$

Thus  $K$  is complete. See Figure 3.7 for a schematic picture of the above construction.

*Step 2. Completeness  $\implies$  Cut Property.* Conversely, suppose that  $K$  is Complete. To prove that  $K$  has the Cut Property, we need to:

- Consider a cut  $(A, B)$  of  $K$ .
- Show that  $(A, B)$  has a separator  $s \in K$ .

This implication is easier. Indeed, since  $A$  is non-empty and bounded above, by Completeness there exists

$$\sup A \in K.$$

We will show that

$$s := \sup A$$

is a separator for the cut  $(A, B)$ . See Figure 3.8 for a schematic picture of the above construction.




Figure 3.7: Let  $s$  be the separator of the cut  $(B^c, B)$ , with  $B$  the set of upper bounds of  $A$ . Then  $s = \sup A$ .




Figure 3.8: Let  $(A, B)$  be a cut of  $K$  and let  $s = \sup A$ . Then  $s$  is the separator of the cut  $(A, B)$ .

Keeping the above ideas in mind, let us proceed with the proof.

**Proof:** Proof of Theorem 3.41

*Step 1. Cut Property  $\implies$  Completeness.*

We need to prove that  $K$  is complete. To this end, consider  $A \subset K$  non-empty and bounded above. Define the set of upper bounds of  $A$ :

$$B := \{b \in K : b \geq a, \forall a \in A\}.$$

*Claim. The pair  $(B^c, B)$  is a cut of  $K$ .*

*Proof of Claim.* We have to prove two points:

- $(B^c, B)$  forms a partition of  $K$ .

Indeed, we have  $B \neq \emptyset$ , since  $A$  is bounded above. Further, we have  $B^c \neq \emptyset$ , since  $A$  is non-empty. Thus

$$K = B^c \cup B, \quad B^c \cap B = \emptyset.$$

Then  $(B^c, B)$  is a partition of  $K$ .

- We have

$$x \leq y, \quad \forall x \in B^c, \forall y \in B. \quad (3.39)$$

To show the above, let  $x \in B^c$  and  $y \in B$ . By definition of  $B$  we have that elements of  $B^c$  are not upper bounds of  $A$ . Therefore  $x$  is not an upper bound. This means there exists  $\tilde{a} \in A$  which is larger than  $x$ , that is,

$$x \leq \tilde{a}.$$

Since  $y \in B$ , then  $y$  is an upper bound for  $A$ , so that

$$\tilde{a} \leq y, \quad \forall \tilde{a} \in A.$$

Therefore

$$x \leq \tilde{a} \leq y,$$

concluding (3.39).

Thus  $(B^c, B)$  is a cut of  $K$  and the claim is proven.

Since  $(B^c, B)$  is a cut of  $K$ , by the Cut Property there exists a separator  $s \in K$  such that

$$x \leq s \leq y, \quad \forall x \in B^c, \forall y \in B. \quad (3.40)$$

*Claim.  $s$  is an upper bound for  $A$ .*

*Proof of Claim.*

Suppose by contradiction that  $s$  is not an upper bound for  $A$ . Therefore by definition of upper bound, there exists  $\tilde{a} \in A$  such that

$$s < \tilde{a}.$$

Consider the mid-point between  $s$  and  $\tilde{a}$ , that is,

$$m := \frac{s + \tilde{a}}{2} \in K.$$

Since  $m$  is the mid-point between  $s$  and  $\tilde{a}$ , and  $s < \tilde{a}$ , it holds

$$s < m < \tilde{a}.$$

Indeed, since  $s < \tilde{a}$  then

$$s = \frac{2s}{2} < \frac{s + \tilde{a}}{2} < \frac{2\tilde{a}}{2} = \tilde{a}.$$

In particular the above tells us that  $m$  is not an upper bound for  $A$ , given that  $\tilde{a} \in A$  and  $m < \tilde{a}$ . Therefore  $m \in B^c$ , by definition of  $B^c$ . Therefore (3.40) implies

$$m \leq s,$$

which contradicts  $s < m$ . Hence  $s$  is an upper bound of  $A$ , concluding the proof of Claim.

*Conclusion.* We have shown that  $s$  is an upper bound of  $A$ . Condition (3.40) tells us that

$$s \leq y, \quad \forall y \in B.$$

Recalling that  $B$  is the set of upper bounds of  $A$ , this means that  $s$  is the smallest upper bound of  $A$ , that is,

$$s = \sup A \in K.$$

*Step 2. Completeness  $\implies$  Cut Property.*

Suppose  $K$  is complete. We need to show that  $K$  has the Cut Property. Therefore assume  $(A, B)$  is a cut of  $K$ , that is,

$$\begin{aligned} A &\neq \emptyset, \quad B \neq \emptyset, \\ K &= A \cup B, \quad A \cap B = \emptyset, \\ a &\leq b, \quad \forall a \in A, \quad \forall b \in B. \end{aligned} \tag{3.41}$$

Since  $B \neq \emptyset$ , from (3.41) it follows that  $A$  is bounded above: indeed, every element of  $B$  is an upper bound for  $A$ , thanks to (3.41). Since  $A \neq \emptyset$ , by the Axiom of Completeness we have

$$s = \sup A \in K.$$

In particular, by definition of supremum, we have

$$a \leq s, \quad \forall a \in A.$$

Let now  $b \in B$  be arbitrary. From (3.41) we have that

$$a \leq b, \quad \forall a \in A. \tag{3.42}$$

Therefore  $b$  is an upper bound of  $A$ . Since  $s = \sup A$ , we have that  $s$  is the smallest upper bound, and so

$$s \leq b.$$

Given that  $b \in B$  was arbitrary, it actually holds

$$s \leq b, \forall b \in B. \quad (3.43)$$

From (3.42) and (3.43) we therefore have

$$a \leq s \leq b, \forall a \in A, \forall b \in B,$$

showing that  $s$  is a separator of  $(A, B)$ . Thus  $K$  has the Cut Property.

## 3.7 Axioms of Real Numbers

We now have all the key elements to introduce the Real Numbers  $\mathbb{R}$ . These ingredients are:

- Definition of **ordered field**,
- The **Cut Property** or **Axiom of Completeness**.

The definition of  $\mathbb{R}$  is given in an axiomatic way.

### Definition 3.43: System of Real Numbers $\mathbb{R}$

A system of Real Numbers is a set  $\mathbb{R}$  satisfying the following properties:

1. There is an operation  $+$  of **addition** on  $\mathbb{R}$

$$+ : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}, \quad (x, y) \mapsto x + y$$

The addition satisfies:  $\forall x, y, z \in \mathbb{R}$

- (A1) **Commutativity and Associativity:**

$$x + y = y + x$$

$$(x + y) + z = x + (y + z)$$

- (A2) **Additive Identity:**  $\exists 0 \in \mathbb{R}$  s.t.

$$x + 0 = 0 + x = x$$

- (A3) **Additive Inverse:**  $\exists (-x) \in \mathbb{R}$  s.t.

$$x + (-x) = (-x) + x = 0$$

2. There is an operation  $\cdot$  of **multiplication** on  $\mathbb{R}$

$$\cdot : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}, \quad (x, y) \mapsto x \cdot y = xy$$

The multiplication satisfies:  $\forall x, y, z \in \mathbb{R}$

- (M1) **Commutativity and Associativity:**

$$x \cdot y = y \cdot x$$

$$(x \cdot y) \cdot z = x \cdot (y \cdot z)$$

- (M2) **Multiplicative Identity:**  $\exists 1 \in \mathbb{R}$  s.t.

$$x \cdot 1 = 1 \cdot x = x$$

- (M3) **Multiplicative Inverse:** If  $x \neq 0$ ,  $\exists x^{-1} \in \mathbb{R}$  s.t.

$$x \cdot x^{-1} = x^{-1} \cdot x = 1$$

3. There is a relation  $\leq$  of **total order** on  $\mathbb{R}$ . The order satisfies:  $\forall x, y, z \in \mathbb{R}$

- (O1) **Reflexivity:**

$$x \leq x$$

- (O2) **Antisymmetry:**

$$x \leq y \text{ and } y \leq x \implies x = y$$

- (O3) **Transitivity:**

$$x \leq y \text{ and } y \leq z \implies x \leq z$$

- (O4) **Total order:**

$$x \leq y \text{ or } y \leq x$$

4. The operations  $+$  and  $\cdot$ , and the total order  $\leq$ , are related by the following properties:  $\forall x, y, z \in \mathbb{R}$

- (AM) **Distributive:** Relates addition and multiplication via

$$x \cdot (y + z) = x \cdot y + x \cdot z$$

- (AO) Relates addition and order with the requirement:

$$x \leq y \implies x + z \leq y + z$$

- (MO) Relates multiplication and order with the requirement:

$$x \geq 0, y \geq 0 \implies x \cdot y \geq 0$$

5. **Cut Property** holds:

- (CP) Every cut  $(A, B)$  of  $\mathbb{R}$  admits a **separator**  $s \in \mathbb{R}$  s.t.

$$a \leq s \leq b, \quad \forall a \in A, \forall b \in B$$

**Remark 3.44**

Since **Cut Property** and **Axiom of Completeness** are equivalent by Theorem 3.41, one can replace the Cut Property in Definition 3.43 Point 5 with:

5. **Axiom of Completeness** holds:

- (AC) For every  $A \subset \mathbb{R}$  non-empty and bounded above

$$\sup A \in \mathbb{R}$$

**Notation 3.45**

For  $x \in \mathbb{R}$ ,  $x \neq 0$ , the multiplicative inverse is also denoted by

$$x^{-1} = \frac{1}{x}.$$

**Remark 3.46**

Recall that

- $(K, +, \cdot)$  satisfying

$$(A_1)-(A_3), (M_1)-(M_3), (AM)$$

is a **field**

- $(K, +, \cdot, \geq)$  satisfying

$$(A_1)-(A_3), (M_1)-(M_3), (O_1)-(O_4), (AM), (AO), (MO)$$

is an **ordered field**

In particular we have:

- $(\mathbb{N}, +, \cdot)$  is **not a field**
- $(\mathbb{Z}, +, \cdot)$  is **not a field**
- $(\mathbb{Q}, +, \cdot)$  is a **field**
- $(\mathbb{Q}, +, \cdot, \geq)$  is an **ordered field**
- $(\mathbb{R}, +, \cdot, \geq)$  is a **complete ordered field**: that is, an ordered field in which the **Cut Property** (CP) or **Axiom of Completeness** (AC) hold

**Important**

It can be shown that  $(\mathbb{R}, +, \cdot, \geq)$  is the **only** complete ordered field.

The above has to be intended in the following sense: if  $(K, +, \cdot, \geq)$  is another complete ordered field, then  $K$  *looks like*  $\mathbb{R}$ . Mathematically this means that there exists an invertible map  $\Psi : \mathbb{R} \rightarrow K$ , called isomorphism of fields, which preserves  $+$ ,  $\cdot$  and  $\leq$ .

**Question 3.47**

We have only postulated the existence of  $\mathbb{R}$ . Does such complete ordered field actually exist?

The answer is **YES**, and there are several models for such  $\mathbb{R}$ . If time allows, we will look into one of these models at the end of the module.

## 4 Properties of $\mathbb{R}$

Now that we established the axiomatic definition of the Real Numbers  $\mathbb{R}$  as a **complete ordered field**, let us investigate some of the properties of  $\mathbb{R}$ . These will be consequence of the axioms of the real numbers, particularly of the **Axiom of Completeness**.

### 4.1 Special subsets of $\mathbb{R}$

### 4.2 Archimedean Property

The Archimedean property is one of the most useful properties of  $\mathbb{R}$ , and it essentially states that the set of natural numbers  $\mathbb{N}$  is not bounded above in  $\mathbb{R}$ .

More precisely, the Archimedean Property says two things:

1. For any  $x \in \mathbb{R}$  we can always find a natural number  $n \in \mathbb{N}$  such that

$$n > x.$$

2. For any  $x \in \mathbb{R}$  with  $x > 0$ , we can always find a natural number  $m \in \mathbb{N}$  such that

$$0 < \frac{1}{m} < x.$$

The situation is depicted in Figure 4.2.




Figure 4.1: For any  $x > 0$  we can find  $n, m \in \mathbb{N}$  such that  $1/m < x < n$ .

#### Remark 4.1

The Archimedean property might sound trivial. However it far from trivial. Indeed there are examples of ordered fields  $K$  that satisfy:

1.  $\mathbb{N} \subset K$ .



2.  $K$  does not have the Archimedean property.
3. In particular,  $\mathbb{N}$  is bounded above in  $K$ .

Let us proceed with the precise statement of the Archimedean property in  $\mathbb{R}$ .

### Theorem 4.2: Archimedean Property

Let  $x \in \mathbb{R}$  be given. Then:

1. There exists  $n \in \mathbb{N}$  such that

$$n > x.$$

2. Suppose in addition that  $x > 0$ . There exists  $n \in \mathbb{N}$  such that

$$\frac{1}{n} < x.$$

### Proof

*Part 1.* Let  $x \in \mathbb{R}$ . Suppose by contradiction that there is no  $n \in \mathbb{N}$  such that

$$n > x.$$

This means that

$$n \leq x \quad \forall n \in \mathbb{N}. \quad (4.1)$$

The above is saying that the set  $\mathbb{N}$  is bounded above. Since  $\mathbb{N}$  is not empty, by the Axiom of Completeness there exists

$$\alpha = \sup \mathbb{N}.$$

*Claim:*  $(\alpha - 1)$  is not an upper bound for  $\mathbb{N}$ .

*Proof of Claim.* Indeed, we have

$$(\alpha - 1) < \alpha. \quad (4.2)$$

Therefore  $\alpha - 1$  cannot be an upper bound for  $\mathbb{N}$ . Indeed, if by contradiction  $\alpha - 1$  was an upper bound for  $\mathbb{N}$ , then we would have

$$\alpha \leq (\alpha - 1),$$

since  $\alpha$  is the smallest upper bound for  $\mathbb{N}$ . This contradicts (4.2). Therefore  $\alpha - 1$  is not an upper bound for  $\mathbb{N}$ .

*Conclusion.* Since  $\alpha - 1$  is not an upper bound for  $\mathbb{N}$ , there exists  $n_0 \in \mathbb{N}$  such that

$$\alpha - 1 < n_0.$$

The above implies

$$\alpha < n_0 + 1.$$

Since

$$(n_0 + 1) \in \mathbb{N},$$

we have obtained a contradiction, given that  $\alpha$  was the supremum of  $\mathbb{N}$ . Thus (4.1) is false, meaning that there exists  $n \in \mathbb{N}$  such that

$$n > x.$$

*Part 2.* Suppose  $x \in \mathbb{R}$  with  $x > 0$ . We can define

$$y := \frac{1}{x}.$$

By Part 1 there exists  $n \in \mathbb{N}$  such that

$$n > y = \frac{1}{x}.$$

Using that  $x > 0$ , we can rearrange the above inequality to obtain

$$\frac{1}{n} < x,$$

which is the desired thesis.

There is another formulation of the Archimedean Property which, depending on the situation, might be more useful. This formulation says the following: If  $x, y \in \mathbb{R}$  are such that

$$0 < x < y,$$

then there exists  $n \in \mathbb{N}$  such that

$$nx > y.$$

In other words, if one does  $n$  steps of size  $x$  in the positive numbers direction, then the resulting number  $nx$  will be larger than  $y$ . The situation is depicted in Figure 4.2.




Figure 4.2: For  $0 < x < y$  there exists  $n \in \mathbb{N}$  such that  $nx > y$ . In the picture  $n = 3$ .

**Theorem 4.3:** Archimedean Property (Alternative formulation)

Let  $x, y \in \mathbb{R}$ , with  $0 < x < y$ . There exists  $n \in \mathbb{N}$  such that

$$nx > y.$$

**Proof**

Suppose by contradiction that there does not exist some  $n \in \mathbb{N}$  such that

$$nx > y.$$

This means that

$$nx \leq y, \quad \forall n \in \mathbb{N}. \quad (4.3)$$

Define the set

$$A := \{nx : n \in \mathbb{N}\}.$$

Condition (4.3) is saying that  $A$  is bounded above by  $y$ . Moreover  $A$  is trivially non-empty. By the Axiom of Completeness there exists

$$\alpha = \sup A.$$

Since  $\alpha$  is the supremum of  $A$ , by definition of supremum and of the set  $A$ , we have

$$nx \leq \alpha, \quad \forall n \in \mathbb{N}. \quad (4.4)$$

As (4.4) holds for every  $n \in \mathbb{N}$ , then it also holds for  $(n+1)$ , meaning that

$$(n+1)x \leq \alpha.$$

The above implies

$$nx \leq \alpha - x.$$

As  $n$  was arbitrary, we conclude that

$$nx \leq \alpha - x, \quad \forall n \in \mathbb{N}.$$

The above is saying that  $(\alpha - x)$  is an upper bound for  $A$ . Since  $\alpha$  is the supremum of  $A$ , in particular  $\alpha$  is the smallest upper bound. Thus it must hold

$$\alpha \leq \alpha - x.$$

The above is equivalent to

$$x \leq 0,$$

which contradicts our assumption of  $x > 0$ . Therefore (4.3) is false, and there exists  $n \in \mathbb{N}$  such that

$$nx > y,$$

concluding the proof.

### 4.3 Nested Interval Property

Another consequence of the axiom of completeness is the **Nested Interval Property**. This is yet another way of saying the same thing:  $\mathbb{R}$  does not have **gaps**.

Let us look at a construction. Suppose given some closed intervals

$$I_n := [a_n, b_n] = \{x \in \mathbb{R} : a_n \leq x \leq b_n\},$$

where the end points are ordered in the following way:

$$a_1 \leq a_2 \leq \dots \leq a_n \leq \dots \leq b_n \leq \dots \leq b_2 \leq b_1,$$

as shown in Figure 4.3.




Figure 4.3: Nested intervals  $I_n = [a_n, b_n]$ .

The intervals  $I_n$  are **nested**, meaning that

$$I_1 \supset I_2 \supset I_3 \supset \dots \supset I_n \supset \dots$$

For finite intersections we clearly have

$$\bigcap_{n=1}^k I_n = I_k,$$

that is, intersecting the first  $k$  intervals yields  $I_k$ , the smallest interval in the sequence.

#### Question 4.4

Consider the infinite intersection

$$\bigcap_{n=1}^{\infty} I_n := \{x \in \mathbb{R} : x \in I_n, \forall n \in \mathbb{N}\}.$$

What can we say about it? Is it empty? Is it not empty?

The answer is that the infinite intersection is not empty, because  $\mathbb{R}$  was constructed in a way that it does not have gaps.

**Theorem 4.5:** Nested Interval Property

For each  $n \in \mathbb{N}$  assume given a closed interval

$$I_n := [a_n, b_n] = \{x \in \mathbb{R} : a_n \leq x \leq b_n\}.$$

Suppose that the intervals are nested, that is,

$$I_n \supset I_{n+1}, \quad \forall n \in \mathbb{N}.$$

Then

$$\bigcap_{n=1}^{\infty} I_n \neq \emptyset. \quad (4.5)$$

**Proof**

By definition we have

$$\bigcap_{n=1}^{\infty} I_n := \{x \in \mathbb{R} : x \in I_n, \forall n \in \mathbb{N}\}.$$

We want to prove (4.5). This means we need to find a real number  $x$  such that

$$x \in I_n, \quad \forall n \in \mathbb{N}. \quad (4.6)$$

In particular (4.6) is saying that it should hold

$$x \geq a_n, \quad \forall n \in \mathbb{N}.$$

We might be tempted to choose  $x$  to be any of the  $b_n$ . This choice would indeed satisfy the above. However (4.6) also implies that

$$x \leq b_n, \quad \forall n \in \mathbb{N}.$$

Therefore  $x$  has to be larger than all the  $a_n$ , but not too large. This suggests that  $x$  should be defined as a supremum. Define the set

$$A := \{a_n : n \in \mathbb{N}\}.$$

The set  $A$  is non-empty and is bounded above by any of the  $b_n$ . Therefore there exists

$$x = \sup A.$$

By definition of supremum and definition of the set  $A$ , we have

$$a_n \leq x, \quad \forall n \in \mathbb{N}.$$

On the other hand, consider an arbitrary number  $b_n$ . By construction we have

$$a_i \leq b_n, \quad \forall i \in \mathbb{N}.$$

Therefore  $b_n$  is an upper bound for  $A$ . Since the supremum is the smallest upper bound, we conclude that

$$x \leq b_n.$$

The index  $n$  was chosen arbitrarily, and therefore

$$x \leq b_n, \quad \forall n \in \mathbb{N}.$$

In total we have

$$a_n \leq x \leq b_n, \quad \forall n \in \mathbb{N},$$

showing that  $x$  satisfies (4.6). Therefore (4.5) holds and the proof is concluded.

### Important

The assumption that  $I_n$  is **closed** is crucial in Theorem 4.5. Without such assumption the thesis of Theorem 4.5 does not hold in general, as seen in Example 4.6 below.

### Example 4.6

Consider the **open** intervals

$$I_n := \left(0, \frac{1}{n}\right).$$

These are clearly nested

$$I_n \supset I_{n+1}, \quad \forall n \in \mathbb{N}.$$

For this choice of  $I_n$  we have

$$\bigcap_{n=1}^{\infty} I_n = \emptyset. \quad (4.7)$$

Indeed, suppose by contradiction that the intersection is non-empty. Then there exists  $x \in \mathbb{N}$  such that

$$x \in I_n, \quad \forall n \in \mathbb{N}.$$

By definition of  $I_n$  the above reads

$$0 < x < \frac{1}{n}, \quad \forall n \in \mathbb{N}. \quad (4.8)$$

Since  $x > 0$ , by the Archimedean Property in Theorem 4.2 Point 1, there exists  $n_0 \in \mathbb{N}$  such that

$$0 < \frac{1}{n_0} < x.$$

The above contradicts (4.8). Therefore (4.7) holds.

## 4.4 Revisiting Sup and inf

We now investigate some of the properties of supremum and infimum in  $\mathbb{R}$ . The first property is an alternative characterization of the supremum, which we will often use.

### Proposition 4.7: Characterization of Supremum

Let  $A \subset \mathbb{R}$  be a non-empty set. Suppose that  $s \in \mathbb{R}$  is an upper bound for  $A$ . They are equivalent:

1.  $s = \sup A$
2. For every  $\varepsilon > 0$  there exists  $x \in A$  such that

$$s - \varepsilon < x.$$

### Proof

*Step 1.* Assume that

$$s = \sup A.$$

Let  $\varepsilon > 0$  be arbitrary. We clearly have that

$$s - \varepsilon < s. \tag{4.9}$$

Therefore  $(s - \varepsilon)$  cannot be an upper bound of  $A$ . Indeed, if by contradiction  $(s - \varepsilon)$  was an upper bound, then we would have

$$s \leq (s - \varepsilon),$$

since  $s$  is the smallest upper bound. The above contradicts (4.9), and therefore  $(s - \varepsilon)$  is not an upper bound for  $A$ . Hence there exists some  $x \in A$  such that

$$s - \varepsilon < x,$$

concluding.

*Step 2.* Assume that Point 2 in the statement of Proposition 4.7 holds. By assumption we have that  $s$  is an upper bound for  $A$ . Suppose by contradiction that

$$s \neq \sup A.$$

This is equivalent to the statement

$$s \text{ is not the smallest upper bound of } A. \tag{4.10}$$

Hence there exists an upper bound  $b$  of  $A$  such that

$$b < s.$$

Let

$$\varepsilon := s - b.$$

By assumption there exists  $x \in A$  such that

$$s - \varepsilon < x.$$

Substituting the definition of  $\varepsilon$  we get

$$s - s + b < x \implies b < x.$$

Since  $b$  is an upper bound for  $A$  and  $x \in A$ , the above is a contradiction. Therefore (4.10) is false, and  $s$  is the smallest upper bound of  $A$ . Thus  $s = \sup A$ .

The analogue of Proposition 4.7 is as follows. The proof is left as an exercise.

**Proposition 4.8:** Characterization of Infimum

Let  $A \subset \mathbb{R}$  be a non-empty set. Suppose that  $i \in \mathbb{R}$  is a lower bound for  $A$ . They are equivalent:

1.  $i = \inf A$
2. For every  $\varepsilon \in \mathbb{R}$ , with  $\varepsilon > 0$ , there exists  $x \in A$  such that

$$x < i + \varepsilon.$$

With the above characterizations of supremum and infimum, it is now easier to prove that some candidate number is the supremum or infimum of some set. As an example, let us characterize supremum and infimum of an open interval of  $\mathbb{R}$ .

**Proposition 4.9**

Let  $a, b \in \mathbb{R}$  with  $a < b$ . Let

$$A := (a, b) = \{x \in \mathbb{R} : a < x < b\}.$$

Then

$$\inf A = a, \quad \sup A = b.$$

**Proof**

We will only prove that

$$\inf A = a,$$

since the proof of  $\sup A = b$  is similar.

By definition of  $A$ , we have that  $a$  is a lower bound for  $A$ , since

$$a < x, \quad \forall x \in A.$$



We need to prove that  $a$  is the largest lower bound of  $A$ . To this end, let  $L$  be a lower bound for  $A$ , that is,

$$L \leq x, \quad \forall x \in A.$$

We have to prove that

$$L \leq a. \tag{4.11}$$

Indeed suppose by contradiction that (4.11) does not hold, namely that

$$a < L.$$

Since  $a < b$  we have

$$a = \frac{2a}{2} < \frac{a+b}{2} < \frac{2b}{2} = b,$$

showing that the midpoint between  $a$  and  $b$  satisfies

$$\frac{a+b}{2} \in A.$$

Since  $L$  is a lower bound for  $A$  we have

$$L \leq \frac{a+b}{2} < b. \tag{4.12}$$

Consider the midpoint

$$M := \frac{a+L}{2}.$$

We have that

$$M \in A.$$

Indeed, recalling that  $a < L$ , we have

$$a = \frac{2a}{2} < \frac{a+L}{2} = M.$$

Moreover by (4.12) we have  $L < b$ . Thus

$$M = \frac{a+L}{2} \leq \frac{a+b}{2} < b.$$

This shows  $M \in A$ .

Moreover

$$M < L.$$

This is because  $a < L$ , and therefore

$$M = \frac{a+L}{2} < \frac{2L}{2} = L.$$

This is a contradiction, since by assumption  $L$  is a lower bound for  $A$ , and thus we should have

$$L \leq M.$$

Therefore (4.11) holds, showing that  $a$  is the largest lower bound of  $A$ . Thus  $a = \inf A$ .

As a corollary of the above we have that the maximum and minimum of an open interval do not exist.

### Corollary 4.10

Let  $a, b \in \mathbb{R}$  with  $a < b$ . Let

$$A := (a, b) = \{x \in \mathbb{R} : a < x < b\}.$$

Then  $\min A$  and  $\max A$  do not exist.

### Proof

Suppose by contradiction that  $\min A$  exists. We have shown that if the minimum of a set exists, then it must be

$$\min A = \inf A.$$

Since

$$\inf A = a,$$

by Proposition 4.9, we would obtain that

$$\min A = a.$$

By definition  $\min A \in A$ , so that  $a \in A$ . This is contradiction. Then  $\min A$  does not exist. The proof that  $\max A$  does not exist is similar, and is left as an exercise.

We can also consider intervals for which one or both of the sides are closed.

### Corollary 4.11

Let  $a, b \in \mathbb{R}$  with  $a < b$ . Let

$$A := [a, b) = \{x \in \mathbb{R} : a \leq x < b\}.$$

Then

$$\min A = \inf A = a, \quad \sup A = b,$$

$\max A$  does not exist.

The proof is very similar to the ones above, and is left to the reader for exercise. Let us now compute supremum and infimum of a set which is not an interval.

**Proposition 4.12**

Define the set

$$A := \left\{ \frac{1}{n} : n \in \mathbb{N} \right\}.$$

Then

$$\inf A = 0, \quad \sup A = \max A = 1.$$

**Proof**

*Part 1.* We have

$$\frac{1}{n} \leq 1, \quad \forall n \in \mathbb{N}.$$

Since  $1 \in A$ , we conclude that

$$\max A = \sup A = 1.$$

*Part 2.* We have

$$\frac{1}{n} > 0, \quad \forall n \in \mathbb{N},$$

showing that 0 is a lower bound for  $A$ . Suppose by contradiction that 0 is not the infimum. Therefore 0 is not the largest lower bound. Then there exists  $\varepsilon \in \mathbb{R}$  such that:

- $\varepsilon$  is a lower bound for  $A$ , that is,

$$\varepsilon \leq \frac{1}{n}, \quad \forall n \in \mathbb{N}, \tag{4.13}$$

- $\varepsilon$  is larger than 0:

$$0 < \varepsilon.$$

As  $\varepsilon > 0$ , by the Archimedean Property there exists  $n_0 \in \mathbb{N}$  such that

$$0 < \frac{1}{n_0} < \varepsilon.$$

This contradicts (4.13). Thus 0 is the largest lower bound of  $A$ , that is,  $0 = \inf A$ .

*Part 3.* We have that  $\min A$  does not exist. Indeed suppose by contradiction that  $\min A$  exists. Then

$$\min A = \inf A.$$

As  $\inf A = 0$  by Part 2, we conclude  $\min A = 0$ . As  $\min A \in A$ , we obtain  $0 \in A$ , which is a contradiction.

## 4.5 Density of $\mathbb{Q}$ in $\mathbb{R}$

In this section we show that  $\mathbb{Q}$  is dense in  $\mathbb{R}$ . Density can be explained in two equivalent ways:

- For every given number  $x \in \mathbb{R}$ , there exists a rational number  $r \in \mathbb{Q}$  which is arbitrarily close to  $x$ .
- For every pair of  $a, b \in \mathbb{R}$  with  $a < b$ , there exists  $r \in \mathbb{Q}$  such that

$$a < r < b.$$

In other words, the above statements are saying that  $\mathbb{Q}$  and  $\mathbb{R}$  are tightly knitted together. Let us give a proof of this statement.

**Theorem 4.13:** Density of  $\mathbb{Q}$  in  $\mathbb{R}$ 

Let  $a, b \in \mathbb{R}$ , with  $a < b$ . There exists  $r \in \mathbb{Q}$  such that

$$a < r < b.$$

The density result of Theorem 4.13 has a surprising corollary. Indeed, we have constructed the real numbers  $\mathbb{R}$  so that they would fill the **gaps** of  $\mathbb{Q}$ .

**Question 4.14**

How many gaps does  $\mathbb{Q}$  have? In other words, how many irrational numbers are out there?

The answer is that there are as many irrational numbers as there are real numbers. Moreover for every pair of real numbers, we can always find an irrational number in between.

**Corollary 4.15**

Let  $a, b \in \mathbb{R}$ , with  $a < b$ . There exists  $t \in (\mathbb{R} \setminus \mathbb{Q})$  such that

$$a < t < b.$$

**Proof**

Abbott

## 4.6 Existence of Square Roots

## 4.7 Cardinality

Brief. Use Dommmers

# 5 Complex Numbers

Coming soon

# 6 Sequences

Coming soon

$\mathbb{R}$

## 6.1 Example: Heron's Method

The first explicit algorithm for approximating

$$\sqrt{x}$$

for  $x > 0$  is known as **Heron's method**, after the first-century Greek mathematician [Heron of Alexandria](#) who described the method in his AD 60 work *Metrica*, see reference to [Wikipedia page](#).

Let us see what is the idea of the algorithm:

- Suppose that  $a_1$  is an approximation of  $\sqrt{x}$  from above, that is,

$$\sqrt{x} < a_1. \tag{6.1}$$

- Multiplying (6.1) by  $\sqrt{x}/a_1$  we obtain

$$\frac{x}{a_1} < \sqrt{x}, \tag{6.2}$$

obtaining an approximation of  $\sqrt{x}$  from below.


- Therefore, putting together the above inequalities,

$$\frac{x}{a_1} < \sqrt{x} < a_1 \tag{6.3}$$

- If we take the average of the points  $x/a_1$  and  $a_1$ , it is reasonable to think that we find a better approximation of  $\sqrt{x}$ . Thus our next approximation is

$$a_2 := \frac{1}{2} \left( a_1 + \frac{x}{a_1} \right),$$

see figure below.

Figure 6.1: Heron's Algorithm for approximating  $\sqrt{x}$ 

Iterating, we define by recurrence the sequence

$$a_{n+1} := \frac{1}{2} \left( a_n + \frac{x}{a_n} \right)$$

for all  $n \in \mathbb{N}$ , where the initial guess  $a_1$  has to satisfy (6.1). The aim of the section is to show that

$$\lim_{n \rightarrow \infty} a_n = \sqrt{x}. \quad (6.4)$$

We start by showing that (6.3) holds for all  $n \in \mathbb{N}$ .

### Proposition 6.1

We have

$$\frac{x}{a_n} < \sqrt{x} < a_n \quad (6.5)$$

for all  $n \in \mathbb{N}$ .

### Proof

We prove it by induction:

1. By (6.1) and (6.2) we know that (6.5) holds for  $n = 1$ .
2. Suppose now that (6.5) holds for  $n$ . Then

$$a_{n+1} - \sqrt{x} = \frac{1}{2} \left( a_n + \frac{x}{a_n} \right) - \sqrt{x} \quad (6.6)$$

$$= \frac{1}{2a_n} (a_n^2 + x - 2a_n\sqrt{x}) \quad (6.7)$$

$$= \frac{1}{2a_n} (a_n - \sqrt{x})^2 > 0, \quad (6.8)$$

since we are assuming that  $a_n > \sqrt{x}$ . Therefore

$$\sqrt{x} < a_{n+1}. \quad (6.9)$$

Multiplying the above by  $\sqrt{x}/a_{n+1}$  we get

$$\frac{x}{a_{n+1}} < \sqrt{x}. \quad (6.10)$$



Inequalities (6.9) and (6.10) show that (6.5) holds for  $n + 1$ .

Therefore we conclude (6.5) by the Principle of Induction.

We are now ready to prove error estimates, that is, estimating how far away  $a_n$  is from  $\sqrt{x}$ .

**Proposition 6.2:** Error estimate

For all  $n \in \mathbb{N}$  we have

$$a_{n+1} - \sqrt{x} < \frac{1}{2}(a_n - \sqrt{x}). \quad (6.11)$$

**Proof**

By Proposition 6.1 we know that

$$\frac{x}{a_n} < \sqrt{x}$$

for all  $n \in \mathbb{N}$ . Therefore

$$a_{n+1} = \frac{1}{2} \left( a_n + \frac{x}{a_n} \right) \quad (6.12)$$

$$< \frac{1}{2} (a_n + \sqrt{x}). \quad (6.13)$$

Subtracting  $\sqrt{x}$  from both members in the above inequality we get the thesis.

Inequality (6.11) is saying that the error halves at each step. Therefore we can prove that after  $n$  steps the error is exponentially lower, as detailed in the following proposition.

**Proposition 6.3**

For all  $n \in \mathbb{N}$  we have

$$a_{n+1} - \sqrt{x} < \frac{1}{2^n}(a_1 - \sqrt{x}) \quad (6.14)$$

**Proof**

We prove (6.14) by induction:

1. For  $n = 1$  we have that (6.14) is satisfied, since it coincides with (6.11) for  $n = 1$ .

2. Suppose that (6.14) holds for  $n$ . By (6.11) with  $n$  replaced by  $n + 1$  we have

$$a_{n+2} - \sqrt{x} < \frac{1}{2}(a_{n+1} - \sqrt{x}) \quad (6.15)$$

$$< \frac{1}{2} \cdot \frac{1}{2^n}(a_1 - \sqrt{x}) \quad (6.16)$$

$$= \frac{1}{2^{n+1}}(a_1 - \sqrt{x}) \quad (6.17)$$

where in the second inequality we used the induction hypothesis (6.14). Hence (6.14) holds for  $n + 1$ .

By invoking the Induction Principle we conclude the proof.

Let us comment estimate (6.14). Denote the error at step  $n$  by

$$e_n := a_n - \sqrt{x}.$$

The initial error  $e_1$  depends on how far the initial guess is from  $\sqrt{x}$ . The estimate in (6.14) is telling us that  $e_n$  is a fraction of  $e_1$ , and actually

$$\lim_{n \rightarrow \infty} e_n = 0$$

exponentially fast. From this fact we are finally able to prove (6.4).

#### **Theorem 6.4:** Convergence of Heron's Algorithm

We have that

$$\lim_{n \rightarrow \infty} a_n = \sqrt{x}.$$

#### **Proof**

By Proposition 6.3 we have that

$$a_{n+1} - \sqrt{x} < \frac{1}{2^n}(a_1 - \sqrt{x})$$

Moreover Proposition 6.1 tells us that

$$\sqrt{x} < a_{n+1}.$$

Putting together the two inequalities above we infer

$$\sqrt{x} < a_{n+1} < \sqrt{x} + \frac{1}{2^n}(a_1 - \sqrt{x}). \quad (6.18)$$

Now note that

$$\lim_{n \rightarrow \infty} \frac{1}{2^n} = \lim_{n \rightarrow \infty} \left(\frac{1}{2}\right)^n = 0.$$

Therefore the RHS of (6.18) converges to  $\sqrt{x}$  as  $n \rightarrow \infty$ . Applying the Squeeze Theorem to (6.18) we conclude that  $a_n \rightarrow \sqrt{x}$  as  $n \rightarrow \infty$ .

### 6.1.1 Coding the Algorithm

Heron's Algorithm can be easily coded in Python. For example, see the function below:

```
# x is the number for which to compute sqrt(x)
# guess is the point a_1
# a_1 must be strictly larger than sqrt(x)
# n is the number of iterations
# the function returns a_{n+1}

def herons_algorithm(x, guess, n):
    for i in range(n):
        guess = (guess + x / guess) / 2.0
    return guess
```

For example let us use the Algorithm to compute  $\sqrt{2}$  after 3 iterations. For initial guess we take  $a_1 = 2$ .

```
# Calculate sqrt(2) with 3 iterations and guess 2
sqrt_2 = herons_algorithm(2, 2, 3)

print(f"The sqrt(2) is approximately {sqrt_2}")
```

The `sqrt(2)` is approximately 1.4142156862745097

That is a pretty good approximation in just 3 iterations!

## 6.2 Fibonacci Sequence

## 7 Series

Coming soon

# License

## Reuse

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/)



## Citation

For attribution, please cite this work as:

Fanzon, Silvio. (2023). *Lecture Notes on Numbers, Sequences and Series*.  
<https://www.silviofanzon.com/2023-NSS-Notes/>

BibTex citation:

```
@electronic{Fanzon-NSS-2023,  
  author = {Fanzon, Silvio},  
  title = {Lecture Notes on Numbers, Sequences and Series},  
  url = {https://www.silviofanzon.com/2023-NSS-Notes/},  
  year = {2023}}
```

# References

- [1] S. Abbott. *Understanding Analysis*. Second Edition. Springer, 2015.
- [2] R. G. Bartle and D. R. Sherbert. *Introduction to Real Analysis*. Fourth Edition. Wiley, 2011.
- [3] W. Rudin. *Principles of Mathematical Analysis*. Third Edition. McGraw Hill, 1976.