

\mathbb{E}_2 -ALGEBRA STRUCTURES ON THE DERIVED CENTER OF AN ALGEBRAIC SCHEME

SONJA M. FARR

ABSTRACT. This paper provides an explicit interface between J. Lurie's work on higher centers, and the Hochschild cohomology of an algebraic \mathbb{k} -scheme within the framework of deformation quantization. We first recover a canonical solution to Deligne's conjecture on Hochschild cochains in the affine and global cases, even for singular schemes, by exhibiting the Hochschild complex as an ∞ -operadic center. We then prove that this universal \mathbb{E}_2 -algebra structure precisely agrees with the classical Gerstenhaber bracket and cup product on cohomology in the affine and smooth cases. This last statement follows from our main technical result which allows us to extract the Gerstenhaber bracket of any \mathbb{E}_2 -algebra obtained from a 2-algebra via Lurie's Dunn Additivity Theorem.

CONTENTS

1. Introduction	2
2. Preliminaries	5
2.1. Morphism objects and operadic centers	5
2.2. Tensor product of ∞ -operads	8
2.3. The little cubes operads	10
2.4. Deligne's Conjecture on Hochschild cochains	13
3. The bracket operation on 2-algebras	14
3.1. The bracket operation of an \mathbb{E}_2 -algebra	14
3.2. The bracket operation of a 2-algebra	15
4. Recovering the Hochschild complex as \mathbb{E}_1 -center	20
4.1. The dg nerve	21
4.2. Rectification of algebras over an ∞ -operad	22
4.3. The Hochschild complex	24
4.4. The center as endomorphism object of bimodules	24
4.5. The category of \mathbb{E}_1 -modules in chain complexes	28
4.6. The trouble with universal enveloping algebras	29
4.7. Morphism objects in dg model categories	31
5. The Hochschild complex of a scheme	32
5.1. The ∞ -category of sheaves of \mathbb{k} -modules	32
5.2. The Hochschild complex of a scheme is local	35
5.3. Recovering polydifferential operators as the center of \mathcal{O}_X	40
5.4. Comparison to the classical homotopy Gerstenhaber algebra structure on polydifferential operators	43
Appendix A. The endomorphism ∞ -category	45
References	47

1. INTRODUCTION

M. Kontsevich's Formality Theorem in deformation quantization states that the Hochschild-Kostant-Rosenberg map lifts to a morphism of homotopy Gerstenhaber algebras between polyvector fields and polydifferential operators of a smooth manifold. In [Tam03], D. Tamarkin found an algebraic proof of this theorem, extending it from manifolds to affine space $\mathbb{A}_{\mathbb{k}}^n$ where \mathbb{k} is any field of characteristic zero by showing that the operad of little 2-disks is formal and using Deligne's conjecture on Hochschild cochains. This proof also highlights that these formality morphisms for the smooth Hochschild cochains are non-canonical, since Tamarkin's little disks formality depends on the choice of a Drinfeld Associator.

In fact, using results from D. Bar-Natan [BN98], Tamarkin later showed in [Tam02] that his proof essentially identifies Drinfeld's Associators with operad isomorphisms between the operad of parenthesized braids and the operad of parenthesized chord diagrams which are the identity on objects.

Based on Tamarkin's results, Kontsevich conjectured in [Kon03] that the Grothendieck-Teichmüller group (GT) should act on formality isomorphisms between $T_{\text{poly}}(X)$ and $D_{\text{poly}}(X)$ of a smooth complex variety, and that this action should be of motivic nature, arising as a consequence of the fact that the equations in the Knizhnik-Zamolodchikov Associator are periods. This conjecture was proved by V. Dolgushev, C. Rogers and T. Willwacher in [DRW15]. They were able to show that the Deligne-Drinfeld elements of the Grothendieck-Teichmüller group act by contraction with the odd components of the Chern character of the variety on the cohomology of the sheaf of polyvector fields. In particular, they were able to give examples for which this action is non-trivial.

Dolgushev-Rogers-Willwacher use a result by D. Calaque and M. Van den Bergh in [CVdB10] which shows that the Kontsevich Formality Theorem can be extended to non-affine cases by adding a correction term to the HKR map which depends on the Atiyah class of the variety. Astonishingly, this correction term has the form $J^{1/2}$ with $J = \det(q(\text{At}(X)))$ the Todd class of the variety and

$$q(x) = \frac{x}{1 - e^{-x}}.$$

This is clearly reminiscent of (Kontsevich's generalization of) the classical Duflo Isomorphism Theorem, which states that for any finite dimensional Lie algebra \mathfrak{g} , we get an algebra isomorphism

$$\text{PBW} \circ \det(q^{-1}(\text{ad}))^{1/2} : S(\mathfrak{g})^{\mathfrak{g}} \xrightarrow{\cong} U(\mathfrak{g})^{\mathfrak{g}}.$$

It is a result by A. Alekseev and C. Torossian [AT12] that the Grothendieck-Teichmüller group also acts on "classical" Duflo isomorphisms as above by changing the correction term. Unfortunately, this action was shown in [ABA00] to be trivial.

To this end, note that the codomain $U(\mathfrak{g})^{\mathfrak{g}} = H^0(\mathfrak{g}, U(\mathfrak{g}))$ is the center $Z(\mathfrak{g})$ of the Lie algebra, and can be computed as the cohomology of the universal enveloping algebra. A. Ramadoss [Ram08], J. Roberts and S. Willerton [RW10] and N. Markarian [Mar09] showed that various versions of the Hochschild cochain complex of a smooth scheme X satisfy the universal property of a universal enveloping algebra of the tangent Lie algebra $\mathcal{T}_X[-1]$ in the derived 1-category of \mathcal{O}_X -modules. Therefore, taking hypercohomology, the Hochschild cohomology of X computes a "derived center" of \mathcal{O}_X . The definition of the universal enveloping algebra object in the derived category comes from [HV02] by V. Hinich and A. Vaintrob, where they also defined the center of an associative algebra object. However, this center does not satisfy any universal property, and it also is "external" in the sense that it is just a commutative algebra in the category of sets.

In contrast to this, J. Lurie in [Lur17, Chapter 5] defines a higher center of an algebra A over an ∞ -operad \mathcal{O} in some symmetric monoidal ∞ -category as a universal object in a suitable ∞ -category of algebra actions on A . In particular, the center of A is an associative algebra object in the category of \mathcal{O} -algebras, and in case of the little k -disks ∞ -operads \mathbb{E}_k^\otimes , Lurie uses an ∞ -categorical version of the Dunn Additivity Theorem to show that the higher center of an \mathbb{E}_k -algebra is indeed an \mathbb{E}_{k+1} -algebra. This may be viewed as a generalized Deligne Conjecture.

In this paper we will lay the groundwork for explaining the GT action on Duflo isomorphisms within the theory of centers by connecting Lurie's work on higher centers with the classical results on the Hochschild cohomology of schemes. In particular, we will argue that the Hochschild complex of any quasi-compact separated scheme should be defined as its \mathbb{E}_1 -operadic center, thereby equipping it with a canonical \mathbb{E}_2 -algebra structure.

Given a symmetric monoidal ∞ -category \mathcal{C} and an ∞ -operad \mathcal{O} , we let $\text{Alg}_{\mathcal{O}}(\mathcal{C})$ denote the ∞ -category of \mathcal{O} -algebras in \mathcal{C} . We let $\mathcal{D}(\mathbb{k})$ be the derived ∞ -category of \mathbb{k} -modules. We denote the higher center of an \mathcal{O} -algebra object A by $\mathfrak{Z}_{\mathcal{O}}(A)$.

We will first consider the affine case, and we will prove that we recover a solution to the classical Deligne Conjecture. This is done in Section 4.

Theorem A (Theorem 4.30, Corollary 4.31). *Let A be an associative \mathbb{k} -algebra. The Hochschild complex*

$$C^*(A, A) = \text{Hom}_{\mathbb{k}}(A^{\otimes*}, A)$$

together with the evaluation map is a center for $A \in \text{Alg}_{\mathbb{E}_1}(\mathcal{D}_{\infty}(\mathbb{k}))$. In particular, it is an element of $\text{Alg}_{\mathbb{E}_1}(\text{Alg}_{\mathbb{E}_1}(\mathcal{D}_{\infty}(\mathbb{k}))) \simeq \text{Alg}_{\mathbb{E}_2}(\mathcal{D}_{\infty}(\mathbb{k}))$. Its underlying Gerstenhaber bracket and cup product in cohomology agree with the classical Gerstenhaber algebra structure obtained from the Braces-algebra structure.

In Section 5 we recall a construction of the ∞ -category $\text{Sh}_{\infty}(X)$ of dg-sheaves over a quasi-compact separated scheme X , and examine some basic properties of the center of the structure sheaf as an \mathbb{E}_1 -algebra in this ∞ -category. We define the Hochschild cochain complex of X as the center of \mathcal{O}_X in the \mathbb{k} -linear derived ∞ -category of sheaves on X . We show that it has the desired local properties, even in the singular case. In particular, we show the following.

Theorem B (Theorem 5.51). *Let $U = \text{Spec}(A) \subseteq X$ be an affine open. The map $\mathbb{R}\Gamma_U : \text{Sh}_{\infty}(X) \rightarrow \mathcal{D}_{\infty}(\mathbb{k})$ is lax symmetric monoidal and hence induces a map $\mathbb{R}\Gamma_U : \text{Alg}_{\mathbb{E}_2}(\text{Sh}_{\infty}(X)) \rightarrow \text{Alg}_{\mathbb{E}_2}(\mathcal{D}_{\infty}(\mathbb{k}))$. We have*

$$\mathbb{R}\Gamma_U(\mathfrak{Z}_{\mathbb{E}_1}(\mathcal{O}_X)) \simeq \mathfrak{Z}_{\mathbb{E}_1}(A).$$

In the smooth case, Kontsevich defined the Hochschild cochains to be the sheaf $\mathcal{D}_{\text{poly}}(X)$ of polydifferential operators. This sheaf comes with the structure of a Braces-algebra inherited from the local Hochschild complex, and therefore has the structure of a homotopy Gerstenhaber algebra by Tamarkin's results. In order to compare to the existing GT actions, we need to compare our newly obtained \mathbb{E}_2 -algebra structure to this homotopy Gerstenhaber algebra structure.

Theorem C (Theorem 5.59, Proposition 5.62). *For a smooth quasi-compact separated scheme X of finite type over \mathbb{k} , the sheaf of polydifferential operators $\mathcal{D}_{\text{poly}}(X)$ is a center of \mathcal{O}_X :*

$$\mathcal{D}_{\text{poly}}(X) \simeq \mathfrak{Z}_{\mathbb{E}_1}(\mathcal{O}_X).$$

This equips $\mathcal{D}_{\text{poly}}(X)$ with the structure of an \mathbb{E}_2 -algebra. The corresponding Gerstenhaber algebra in the \mathbb{k} -linear derived 1-category agrees with the classical one coming from the Braces-algebra structure.

In the course of proving these results, we also obtain a couple of technical results about higher centers. In particular, we show how to explicitly obtain the Gerstenhaber bracket of a 2-algebra up to homotopy using Lurie's version of the Dunn Additivity Theorem (see 3.21). We also show that the 2-algebra structure on a center which is obtained as an endomorphism object indeed corresponds to the composition product (Yoneda product) and the convolution product (see 4.35). In a similar manner, we further obtain a stability result stating that the space of Gerstenhaber algebra structures on a center is contractible, which we expect will be helpful in examining action of the Grothendieck-Teichmüller group later on.

Related work. The connection between the Hochschild cochain complex of an associative algebra and its higher center in the form of the universal 2-algebra acting on it was already described by Kontsevich in [Kon03]. In fact, in Deligne's original letter outlining the conjecture, he already stated that he expected the little 2-disk algebra structure to come from the composition and convolution product via some type of Eckmann-Hilton argument. P. Hu, I. Kriz and A. Voronov in [HKV06] proved a simplicial version of Deligne's Conjecture using this idea and explicit models of the little disks operads. In particular, for their result they proved a version of the Dunn Additivity Theorem for these specific models of the little k -disks operads.

In 2013, J. Francis [Fra13] used Lurie's Dunn Additivity Theorem to examine centers of stable ∞ -categories and to relate the center to the module of derivations. He also defines the Hochschild cohomology of an algebra over an ∞ -operad \mathcal{O} as the hom-set of \mathcal{O} -module maps over A from A to itself, which is closely related to Lurie's definition of the center.

In 2020, I. Iwanari [Iwa20] used this definition of the Hochschild cochain complex to show that pair of Hochschild cohomology and homology of a linear category over some commutative ring spectrum gives an algebra over the KS operad, which is a generalization of the \mathbb{E}_2 -operad.

In 2023, C. Brav and N. Rozenblyum [BR23] proved a cyclic version of Deligne's Conjecture (i.e. replacing the little 2-disk operad by the framed version) also using the above described techniques. In particular, their method also relies on Lurie's version of Dunn Additivity. However, none of the above results provide a direct comparison to the classical solutions of Deligne's Conjecture, or make any claim about the underlying Gerstenhaber algebra structure. Similarly, the author is not aware of any comparison of the center of a scheme to the classical sheaf of polydifferential operators.

Conventions. Throughout this paper, \mathbb{k} is a field of characteristic zero. The term "operad" is reserved for non-reduced unital symmetric operads. Complexes are generally chain graded unless states otherwise, and we view non-negatively graded cochain complexes as non-positive chain complexes. We denote the presheaf tensor product simply by " \otimes ", and we decorate symbols with " $(-)^a$ " to indicate sheafification. We try to use the term " ∞ -category" for ∞ -categories, but if nothing else is stated "category" refers to ∞ -category and "1-category" refers to ordinary categories.

Acknowledgments. I thank my thesis advisor, Chris Rogers, for proposing the problem and providing essential guidance. I would also like to thank Nick Rozenblyum for invaluable comments on J. Lurie's proof of the Dunn Additivity Theorem and Dennis Borisov for helpful discussions on dg sheaves. This work was supported by NSF Grant DMS-2305407.

2. PRELIMINARIES

We follow Lurie's formalism for ∞ -operads as developed in [Lur17]. In particular, an ∞ -operad is a morphism $p : \mathcal{O}^\otimes \rightarrow \mathcal{F}\text{in}_*$ of ∞ -categories satisfying a list of conditions making \mathcal{O}^\otimes into an ∞ -category of operators. We will freely use notation from [Lur17] regarding ∞ -operads, algebras and modules over these.

2.1. Morphism objects and operadic centers. We review Lurie's theory of morphism objects and operadic centers. The definitions we use can be found in [Lur17, Section 4.2, 4.7 and 5.3].

Definition 2.1. Let \mathfrak{a} and \mathfrak{m} be the two colors of the ∞ -operad \mathcal{LM}^\otimes . Let $q : \mathcal{C}^\otimes \rightarrow \mathcal{LM}^\otimes$ be a coCartesian fibration of ∞ -operads. We then say that q exhibits the ∞ -category $\mathcal{M} := \mathcal{C}_\mathfrak{m}$ as left tensored over the monoidal ∞ -category $\mathcal{C}_\mathfrak{a}^\otimes := \mathcal{C}^\otimes \times_{\mathcal{LM}^\otimes} \text{Assoc}^\otimes$. In particular, q determines a tensoring

$$\otimes : \mathcal{C}_\mathfrak{a} \times \mathcal{M} \rightarrow \mathcal{M}$$

well-defined up to homotopy that is compatible with the monoidal structure on $\mathcal{C}_\mathfrak{a}$ up to homotopy.

Definition 2.2. Let $q : \mathcal{C}^\otimes \rightarrow \mathcal{LM}^\otimes$ exhibit \mathcal{M} as left tensored over $\mathcal{C}_\mathfrak{a}^\otimes$. Then we denote by

$$\text{LMod}(\mathcal{M}) = \text{Alg}_{/\mathcal{LM}^\otimes}(\mathcal{C})$$

the ∞ -category of pairs of associative algebras in $\mathcal{C}_\mathfrak{a}$ and left modules over them.

In particular, any monoidal ∞ -category is left tensored over itself.

Recall that for ordinary categories, internal homs and more generally enrichments are right adjoint to a tensoring. Similarly, one makes the following definition.

Definition 2.3. Let $\mathcal{C}^\otimes \rightarrow \mathcal{LM}^\otimes$ be a coCartesian fibration of ∞ -operads exhibiting \mathcal{M} as left tensored over $\mathcal{C}_\mathfrak{a}^\otimes$. If $M, N \in \mathcal{M}$, a morphism object for M and N is an object $\text{Mor}(M, N) \in \mathcal{C}_\mathfrak{a}$ together with a map $\alpha \in \text{Map}_\mathcal{M}(\mathcal{C} \otimes M, N)$ such that for each $C \in \mathcal{C}_\mathfrak{a}$, post-composition with α induces a homotopy equivalence

$$(1) \quad \text{Map}_{\mathcal{C}_\mathfrak{a}}(C, \text{Mor}(M, N)) \xrightarrow{\simeq} \text{Map}_\mathcal{M}(C \otimes M, N).$$

We call \mathcal{M} enriched over $\mathcal{C}_\mathfrak{a}^\otimes$ if all the morphisms objects exist.

The following result shows that we can think of morphism objects as the classifying object of action maps $A \otimes M \rightarrow N$ with $A \in \mathcal{C}_\mathfrak{a}$.

Proposition 2.4. *Let $q : \mathcal{C}^\otimes \rightarrow \mathcal{LM}^\otimes$ be a coCartesian fibration of ∞ -operads. Let $M, N \in \mathcal{M}$. Then an object $\text{Mor}(M, N) \in \mathcal{C}_\mathfrak{a}$ together with a map $\alpha : \text{Mor}(M, N) \otimes M \rightarrow N$ is a morphism object of M and N if and only if $(\text{Mor}(M, N), \alpha) \in \mathcal{C}_\mathfrak{a} \times_\mathcal{M} \mathcal{M}_{/N}$ (with map given by $- \otimes M : \mathcal{C}_\mathfrak{a} \rightarrow \mathcal{M}$) is final.*

Proof. Note that $\mathcal{M}_{/N} \rightarrow \mathcal{M}$ is a right fibration, and since these are stable under base change, so is $f : \mathcal{C}_\mathfrak{a} \times_\mathcal{M} \mathcal{M}_{/N} \rightarrow \mathcal{C}_\mathfrak{a}$. Consider the functor $F : \mathcal{C}_\mathfrak{a}^{\text{op}} \rightarrow \mathbf{An}$ classifying f . Then by [Lur09, Lemma 2.2.2.4], its underlying functor $hF : h\mathcal{C}_\mathfrak{a}^{\text{op}} \rightarrow \mathcal{H}$ can be recovered as follows. On objects, $X \in \mathcal{C}_\mathfrak{a}$ is sent to its fiber

$$(\mathcal{C}_\mathfrak{a} \times_\mathcal{M} \mathcal{M}_{/N}) \times_{\mathcal{C}_\mathfrak{a}} \{X\} \simeq \{X \otimes M\} \times_\mathcal{M} \mathcal{M}_{/N} \simeq \text{Map}_\mathcal{M}(X \otimes M, N).$$

Given a morphism $e : Y \rightarrow X \in \text{Mor}_{\mathcal{C}_\mathfrak{a}}(Y, X)$ in $h\mathcal{C}_\mathfrak{a}$, the induced map between the fibers comes from solving the lifting problem

$$\begin{array}{ccc}
 \{1\} \times \mathrm{Map}_{\mathcal{M}}(X \otimes M, N) & \hookrightarrow & \mathcal{C}_{\mathfrak{a}} \times_{\mathcal{M}} \mathcal{M}_{/N} \\
 \downarrow & \nearrow \text{dashed} & \downarrow f \\
 \Delta^1 \times \mathrm{Map}_{\mathcal{M}}(X \otimes M, N) & & \mathcal{C}_{\mathfrak{a}} \\
 & \searrow e & \\
 & \Delta^1 &
 \end{array} ,$$

and restricting the lift to $\{0\} \times \mathrm{Map}_{\mathcal{M}}(X \otimes M, N)$. Since f is a pullback of the right fibration $\mathcal{M}_{/N} \rightarrow \mathcal{M}$, the lift above is induced by the solution to

$$\begin{array}{ccc}
 \{1\} \times \mathrm{Map}_{\mathcal{M}}(X \otimes M, N) & \longrightarrow & \mathcal{M}_{/N} \\
 \downarrow & \nearrow \text{dashed} & \downarrow \\
 \Delta^1 \times \mathrm{Map}_{\mathcal{M}}(X \otimes M, N) & \xrightarrow{e \otimes \mathrm{id}_M} & \mathcal{M}
 \end{array} .$$

But the restriction of this lift to $\{0\} \times \mathrm{Map}_{\mathcal{M}}(X \otimes M, N)$ is given by pre composition with $e \otimes \mathrm{id}_M$. Therefore, we see that hF is given by the composition of $- \otimes M$ and $\mathrm{Map}_{\mathcal{M}}(-, N)$. Recall that by [Lur09, Proposition 4.4.4.5], an object $(X, X \otimes M \xrightarrow{\eta} N)$ is final in $\mathcal{C}_{\mathfrak{a}} \times_{\mathcal{M}} \mathcal{M}_{/N}$ if and only if the pair $(X, \eta \in hF(X))$ represents hF . Then we are done after noting that by definition, $(\mathrm{Mor}(M, N), \alpha)$ is a morphism object exactly if it represents the functor $X \mapsto \mathrm{Map}_{\mathcal{M}}(X \otimes M, N)$. \square

Now put $N = M$. Then by the above proposition, $\mathrm{End}(M) := \mathrm{Mor}(M, M) \in \mathcal{C}_{\mathfrak{a}}$ classifies actions of elements of $\mathcal{C}_{\mathfrak{a}}$ on M . Note however that these are plain actions $A \otimes M \rightarrow M$ that do not require A to be an algebra object and do not require M to satisfy the axioms of a module over A . However, we do expect $\mathrm{End}(M)$ to carry the structure of an associative algebra coming from composition, and M to be a module over it.

Definition 2.5. Let $q : \mathcal{C}^{\otimes} \rightarrow \mathcal{LM}^{\otimes}$ be a coCartesian fibration of ∞ -operads. Define the endomorphism ∞ -category of $M \in \mathcal{M}$ as

$$\mathcal{C}_{\mathfrak{a}}[M] = \mathcal{C}_{\mathfrak{a}} \times_{\mathcal{M}} \mathcal{M}_{/M}.$$

We show in Appendix A that $\mathcal{C}_{\mathfrak{a}}[M]$ agrees with Lurie's endomorphism ∞ -category as in [Lur17, Definition 4.7.1.1]. In particular, it admits the structure of a monoidal ∞ -category with the tensor product given up to homotopy by

$$(A, A \otimes M \xrightarrow{\eta} M) \otimes (B, B \otimes M \xrightarrow{\nu} M) = (A \otimes B, A \otimes B \otimes M \xrightarrow{\mathrm{id} \otimes \nu} A \otimes M \xrightarrow{\eta} M).$$

Then by [Lur17, Corollary 3.2.2.5] with $K = \emptyset$ and $\mathcal{O}^{\otimes} = \mathrm{Assoc}^{\otimes}$ we automatically get the following.

Corollary 2.6. *Assume the ∞ -category $\mathcal{C}_{\mathfrak{a}}[M]$ has a final object $(\mathrm{End}(M), \alpha)$. Then $(\mathrm{End}(M), \alpha)$ can be promoted to an object of $\mathrm{Alg}_{\mathbb{E}_1}(\mathcal{C}_{\mathfrak{a}}[M])$ in an essentially unique way. We denote this object again by $\mathrm{End}(M)$. Note that $\mathrm{End}(M)$ is automatically final in $\mathrm{Alg}_{\mathbb{E}_1}(\mathcal{C}_{\mathfrak{a}}[M])$.*

Up to homotopy, the tensor product of $\mathrm{End}(M)$ with itself is given by the action

$$(\mathrm{End}(M) \otimes \mathrm{End}(M)) \otimes M \simeq \mathrm{End}(M) \otimes (\mathrm{End}(M) \otimes M) \xrightarrow{\mathrm{id}_{\mathrm{End}(M)} \otimes \alpha} \mathrm{End}(M) \otimes M \xrightarrow{\alpha} M,$$

and hence the algebra structure on $\mathrm{End}(M)$ has a multiplication \circ making the following diagram commute

$$\begin{array}{ccc}
 & \text{End}(M) \otimes M & \\
 \nearrow \circ \otimes \text{id} & & \searrow \alpha \\
 (\text{End}(M) \otimes \text{End}(M)) \otimes M & \xrightarrow{\alpha \circ (\text{id} \otimes \alpha)} & M
 \end{array}$$

Further, M is automatically a module over the algebra $\text{End}(M)$, and this action $\text{End}(M) \otimes M \rightarrow M$ is universal among algebra actions on M as a module.

Corollary 2.7. *In the above situation, we have an equivalence of ∞ -categories*

$$\text{Alg}(\mathcal{C}_a[M]) \rightarrow \text{LMod}(\mathcal{M}) \times_{\mathcal{M}} \{M\}.$$

Proof. This follows directly from [Lur17, Theorem 4.7.1.34] and the results of Chapter 4.1.3 in [Lur17]. \square

There are a variety of interesting situations in which such an (endo)morphism object fails to exist, in particular if we consider ∞ -categories arising as categories of algebra objects. The archetypal example is the following.

Example 2.8. Let \mathbb{k} be a field and consider the (symmetric) monoidal category $\mathcal{C} = \text{Alg}_{\mathbb{k}}$ as left tensored over itself. Let $M \in \mathcal{C}$ be some \mathbb{k} -algebra. Then for any endomorphism $\varphi \in \text{Hom}_{\mathcal{C}}(M, M)$, the pair $(k, k \otimes M \cong M \xrightarrow{\varphi} M)$ is an object of $\mathcal{C}[M]$. Therefore, if $(A, \eta) \in \mathcal{C}[M]$ were a final object, then $\eta \circ (u_A \otimes \text{id}_M) = \varphi$ for any endomorphism $\varphi : M \rightarrow M$. But this not possible unless $M = \mathbb{k}$ is the trivial \mathbb{k} -algebra. This was to be expected, since we know that the monoidal category of \mathbb{k} -algebras is not closed.

The solution to this problem is to relax our expectations on the morphism object. In the above discussion, we start out requiring that $\text{Mor}(M, N)$ classify all (plain) actions $C \otimes M \rightarrow N$, and then in the case $N = M$ get for free that $\text{End}(M)$ also classifies algebra actions of algebras on M . Instead, we now consider objects that only classify the algebra actions.

Definition 2.9. Let $\mathcal{C}^{\otimes} \rightarrow \mathcal{LM}^{\otimes}$ be a coCartesian fibration of ∞ -operads, and let $M \in \mathcal{M}$. A center $\mathfrak{Z}(M)$ of M is a final object of $\text{LMod}(\mathcal{M}) \times_{\mathcal{M}} \{M\}$. We generally identify $\mathfrak{Z}(M)$ with its image in $\text{Alg}_{\mathbb{E}_1}(\mathcal{C}_a)$.

Clearly if M admits an endomorphism object, then this endomorphism object is also a center of M . The converse does not hold: The category $\text{Alg}(\mathcal{C}_a[M])$ might have final objects although $\mathcal{C}_a[M]$ does not.

Example 2.10 (Example 2.8 continued). The ordinary center $Z(A)$ of an associative \mathbb{k} -algebra A is indeed the universal algebra object acting on A . To see this, note first that the center is a commutative algebra, and therefore an algebra object in the category of associative algebras. It comes with a natural action on A given by multiplication in A . Now suppose that B is a commutative algebra with action $\eta : B \otimes A \rightarrow A$ making A into a B -module. Then the restriction of η to A yields the identity on A and η must be an algebra morphism. Hence

$$\begin{aligned}
 \eta(b \otimes a) &= \eta(b \otimes 1) \cdot \eta(1 \otimes a) = \eta(b \otimes 1) \cdot a \quad \text{and} \\
 \eta(b \otimes a) &= \eta(1 \otimes a) \cdot \eta(b \otimes 1) = a \cdot \eta(b \otimes 1),
 \end{aligned}$$

showing that η sends B to $Z(A)$.

There also is a relative version of the center.

Definition 2.11. Let $\mathcal{C}^\otimes \rightarrow \mathcal{LM}^\otimes$ be a coCartesian fibration of ∞ -operads, let $\mathbb{1}$ denote the monoidal unit of \mathcal{C}_a^\otimes , and let $f : M \rightarrow N$ be a morphism in \mathcal{M} . A centralizer $\mathfrak{Z}(f)$ of f is a final object in

$$\mathrm{Act}(f) := (\mathcal{C}_a)_{\mathbb{1}/} \times_{\mathcal{M}_{M/}} \mathcal{M}_{M//N}.$$

We generally identify $\mathfrak{Z}(f)$ with its image in \mathcal{C}_a .

The objects of this ∞ -category are given by commuting triangles in \mathcal{M}

$$\begin{array}{ccc} & C \otimes M & \\ \nearrow & & \searrow \\ \mathbb{1} \otimes M & \xrightarrow{f} & N \end{array}.$$

In particular, the centralizer is equipped with an action $\mathfrak{Z}(f) \otimes M \rightarrow N$ making the above diagram commute.

Lemma 2.12. *Let $f : M \rightarrow N$ be a morphism in \mathcal{M} . Let $\overline{M} \in \mathrm{LMod}_{\mathbb{1}}(\mathcal{M})$ be a lift of M as module over the trivial algebra. Let $\mathcal{C}_{\overline{M} \otimes \mathcal{M}}^\otimes \rightarrow \mathcal{LM}^\otimes$ be defined as in [Lur17, Definition 4.2.1.28]. Then centralizers of f can be identified with morphism objects*

$$\mathrm{Mor}_{\mathcal{M}_{M/}}(\mathrm{id}_M, f) \in (\mathcal{C}_a)_{\mathbb{1}/}.$$

Proof. Let $\mathcal{C}'^\otimes := \mathcal{C}_{\overline{M} \otimes \mathcal{M}}^\otimes$. By Proposition 2.4, it suffices to show that we have an equivalence of ∞ -categories $\mathrm{Act}(f) \simeq (\mathcal{C}')_a \times_{\mathcal{M}'} \mathcal{M}'_{/f}$. But we have $(\mathcal{C}')_a \times_{\mathcal{M}'} \mathcal{M}'_{/f} \simeq (\mathcal{C}_a)_{\mathbb{1}/} \times_{\mathcal{M}_{M/}} (\mathcal{M}_{M/})_{/N}$, so this is clear. \square

We would like to see that these notions are compatible, in the sense that the centralizer of an identity morphism recovers the center.

Proposition 2.13 (Proposition 5.3.1.8 [Lur17]). *Let $M \in \mathcal{M}$, and suppose there exists a centralizer $\mathfrak{Z}(\mathrm{id}_M) \in \mathcal{C}_a$. Then there exists a center $\mathfrak{Z}(M) \in \mathrm{Alg}(\mathcal{C}_a)$. Further, a lift of M to a module over an algebra $A \in \mathrm{Alg}_{\mathbb{E}_1}(\mathcal{C}_a)$ exhibits A as a center of M if and only if the action map $A \otimes M \rightarrow M$ exhibits A as a centralizer of id_M .*

Proof. By Lemma 2.12, the centralizer of the identity is a morphism object $\mathrm{Mor}_{\mathcal{M}_{M/}}(\mathrm{id}_M, \mathrm{id}_M)$. By Corollary 2.6, this morphism object admits an essentially unique structure of an algebra object in $(\mathcal{C}_a)_{\mathbb{1}/}$, and id_M lifts to a module over this algebra structure. In particular, $\mathfrak{Z}(\mathrm{id}_M)$ admits a canonical algebra structure making it into the center of id_M in $\mathcal{M}_{M/}$. Now use [Lur17, Lemma 5.3.1.10] to see that the forgetful functor $\mathrm{LMod}(\mathcal{M}_{M/}) \times_{\mathcal{M}_{M/}} \{\mathrm{id}_M\} \rightarrow \mathrm{LMod}(\mathcal{M}) \times_{\mathcal{M}} \{M\}$ preserves final objects. \square

2.2. Tensor product of ∞ -operads. The Boardman-Vogt tensor product on ordinary operads is designed such that algebras over the tensor product $\mathcal{P} \boxtimes_{\mathrm{BV}} \mathcal{O}$ are given by \mathcal{P} -algebras in the category of \mathcal{O} -algebras. However, it is well-known that this tensor product does not make the category of (reduced) operads into a monoidal model category. We briefly review the corresponding construction for ∞ -operads following [Lur17, Section 2.5.5].

We want to capture bilinearity of a map between ∞ -operads. To this end, define a functor $\wedge : \mathrm{Fin}_* \times \mathrm{Fin}_* \rightarrow \mathrm{Fin}_*$ by sending $(\langle m \rangle, \langle n \rangle)$ to the pointed set $(\langle m \rangle^\circ \times \langle n \rangle^\circ)_+ \cong \langle mn \rangle$, where the

isomorphism is given by the lexicographic ordering, and by sending $(f : \langle m \rangle \rightarrow \langle n \rangle, g : \langle m' \rangle \rightarrow \langle n' \rangle)$ to

$$\langle mm' \rangle \xrightarrow{\cong} (\langle m \rangle^\circ \times \langle m' \rangle^\circ)_+ \xrightarrow{f \times g} (\langle n \rangle^\circ \times \langle n' \rangle^\circ)_+ \xrightarrow{\cong} \langle nn' \rangle.$$

Definition 2.14. We call a map of simplicial sets $F : \mathcal{O}^\otimes \times \mathcal{O}'^\otimes \rightarrow \mathcal{O}''^\otimes$ a bifunctor of ∞ -operads if the diagram below commutes, and if F sends pairs of inert maps to an inert map in \mathcal{O}''^\otimes .

$$\begin{array}{ccc} \mathcal{O}^\otimes \times \mathcal{O}'^\otimes & \xrightarrow{F} & \mathcal{O}''^\otimes \\ \downarrow & & \downarrow \\ \mathrm{Fin}_* \times \mathrm{Fin}_* & \xrightarrow{\wedge} & \mathrm{Fin}_* \end{array}$$

In particular, define $\mathrm{Bil}(\mathcal{O}^\otimes, \mathcal{O}'^\otimes; \mathcal{O}''^\otimes)$ to be the full subcategory of $\mathrm{Fun}_{\mathrm{Fin}_*}(\mathcal{O}^\otimes \times \mathcal{O}'^\otimes, \mathcal{O}''^\otimes)$ spanned by the bifunctors.

We claim that for a symmetric monoidal ∞ -category \mathcal{C}^\otimes , the ∞ -category of bifunctors from $\mathcal{O}^\otimes \times \mathcal{O}'^\otimes$ to \mathcal{C}^\otimes is equivalent to the appropriate ∞ -category of \mathcal{O} -algebras in the symmetric monoidal ∞ -category $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes$. To this end, recall the symmetric monoidal structure on the ∞ -category $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})$. This is supposed to capture the fact that the tensor product in \mathcal{C} descends to a tensor product of \mathcal{O}' -algebras in \mathcal{C} . We define a map of simplicial sets $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes \rightarrow \mathrm{Fin}_*$ by the following universal property. If $K \rightarrow \mathrm{Fin}_*$ is a map of simplicial sets, then there is a natural bijection between $\mathrm{Hom}_{\mathrm{Fin}_*}(K, \mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes)$ and the set of diagrams

$$\begin{array}{ccc} K \times \mathcal{O}'^\otimes & \xrightarrow{F} & \mathcal{C}^\otimes \\ \downarrow & & \downarrow \\ \mathrm{Fin}_* \times \mathrm{Fin}_* & \xrightarrow{\wedge} & \mathrm{Fin}_* \end{array}$$

such that for $v \in K$ a vertex and f an inert morphisms in \mathcal{O}'^\otimes , the map $F(s_0(v), f)$ is inert in \mathcal{C}^\otimes . In particular, the fiber $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})_{\langle 1 \rangle}^\otimes$ over $\langle 1 \rangle \in \mathrm{Fin}_*$ is given by the full subcategory of $\mathrm{Fun}_{\mathrm{Fin}_*}(\mathcal{O}'^\otimes, \mathcal{C}^\otimes)$ of maps that preserve inert morphisms, and hence can be identified with the ∞ -category $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})$. Fixing $\langle n \rangle \in \mathrm{Fin}_*$, we get a new map $\mathcal{O}'^\otimes \rightarrow \mathrm{Fin}_*$ given by the following diagram

$$\begin{array}{ccc} \mathcal{O}'^\otimes & \xrightarrow{\quad} & \mathrm{Fin}_* \\ \cong \downarrow & & \uparrow \wedge \\ \Delta^0 \times \mathcal{O}'^\otimes & \xrightarrow{\quad} & \mathrm{Fin}_* \times \mathrm{Fin}_* \end{array}$$

where the lower horizontal map picks out $\langle n \rangle \in \mathrm{Fin}_*$. We can informally describe this map as $\langle n \rangle \wedge p$, if $p : \mathcal{O}'^\otimes \rightarrow \mathrm{Fin}_*$ is the map making \mathcal{O}'^\otimes an ∞ -operad. In particular, it sends $X \in \mathcal{O}'_{\langle m \rangle}$ to $\langle nm \rangle = \langle n \rangle \wedge p(X)$ and $f : X \rightarrow Y$ to $\mathrm{id}_{\langle n \rangle} \wedge p(f)$. The fiber $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})_{\langle n \rangle}^\otimes$ is given by the full subcategory of $\mathrm{Fun}_{\mathrm{Fin}_*}(\mathcal{O}'^\otimes, \mathcal{C}^\otimes)$ sending inert maps to inert maps, where now the map $\mathcal{O}'^\otimes \rightarrow \mathrm{Fin}_*$ is given by this new map $\langle n \rangle \wedge p$. If $F : \mathcal{O}'^\otimes \rightarrow \mathcal{C}^\otimes \in \mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})_{\langle n \rangle}^\otimes$, and $X \in \mathcal{O}'$, then $F(X) \in \mathcal{C}_{\langle n \rangle}^\otimes$ can be described by a tuple $(F(X)_1, \dots, F(X)_n) \in \mathcal{C}^n$. This argument shows that F can indeed be identified with a tuple $(F_1, \dots, F_n) \in \mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})^n$. By [Lur17, Proposition 3.2.4.3] the induced map $q' : \mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes \rightarrow \mathrm{Fin}_*$ is a coCartesian fibration and a morphism f in $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes$ is q' -coCartesian if and only if for every $X \in \mathcal{O}'$, the image $f(X)$ is q -coCartesian in \mathcal{C}^\otimes . In particular, $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes$ is again a symmetric monoidal ∞ -category. For $X \in \mathcal{O}'$, the evaluation map $\mathrm{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes \rightarrow \mathcal{C}^\otimes$ is a morphism of ∞ -operads, hence a lax symmetric monoidal functor, and we see that the symmetric

monoidal structure on $\text{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes$ is given by the pointwise tensor product on \mathcal{C}^\otimes .

An \mathcal{O} -algebra in $\text{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes$ is given by a morphism of simplicial sets $\mathcal{O}^\otimes \rightarrow \text{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes$ over Fin_* sending inert morphisms to inert morphisms. In particular, by the construction of $\text{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes$, such an \mathcal{O} -algebra is given by a diagram

$$\begin{array}{ccc} \mathcal{O}^\otimes \times \mathcal{O}'^\otimes & \longrightarrow & \mathcal{C}^\otimes \\ \downarrow & & \downarrow \\ \text{Fin}_* \times \text{Fin}_* & \xrightarrow{\wedge} & \text{Fin}_* \end{array}$$

such that for every $X \in \mathcal{O}^\otimes$ and every inert map f in \mathcal{O}'^\otimes , the tuple (id_X, f) is sent to an inert map in \mathcal{C}^\otimes . The condition that inert morphisms in \mathcal{O}^\otimes are sent to inert maps in $\text{Alg}_{\mathcal{O}'}(\mathcal{C})^\otimes$ translates to the fact that for inert maps f in \mathcal{O}^\otimes and $X \in \mathcal{O}'$, the tuple (f, id_X) is sent to an inert map in \mathcal{C}^\otimes ; and together those two conditions say exactly that tuples of inert maps are sent to an inert map. But this is clearly the same as a bifunctor $\mathcal{O}^\otimes \times \mathcal{O}'^\otimes \rightarrow \mathcal{C}^\otimes$.

Definition 2.15. We say that a bifunctor $F : \mathcal{O}^\otimes \times \mathcal{O}'^\otimes \rightarrow \mathcal{O}''^\otimes$ exhibits \mathcal{O}''^\otimes as a tensor product of \mathcal{O}^\otimes and \mathcal{O}'^\otimes if for every ∞ -operad \mathcal{C}^\otimes , precomposition with F determines an equivalence of ∞ -categories

$$\text{Alg}_{\mathcal{O}''}(\mathcal{C}) \rightarrow \text{Bil}(\mathcal{O}^\otimes, \mathcal{O}'^\otimes; \mathcal{C}^\otimes).$$

In particular, in this case, if \mathcal{C}^\otimes is a symmetric monoidal ∞ -category, the above discussion shows that we have an equivalence of ∞ -categories

$$\text{Alg}_{\mathcal{O}''}(\mathcal{C}) \rightarrow \text{Alg}_{\mathcal{O}}(\text{Alg}_{\mathcal{O}'}(\mathcal{C})).$$

In this sense, the tensor product of ∞ -operads is a derived version of the Bordmann-Vogt tensor products of operads.

2.3. The little cubes operads. Consider the topological (one-colored) operad \mathbb{E}_k^T with $\mathbb{E}_k^T(n) = \text{Rect}(\square^k \times \{1, \dots, n\}, \square^k)$ the space of rectilinear embeddings. Let $\mathbb{E}_k^{T, \otimes}$ denote its topological category of operators, and consider the corresponding simplicial category $\text{Sing}_\bullet(\mathbb{E}_k^{T, \otimes})$. Taking the homotopy coherent nerve, we obtain an ∞ -category \mathbb{E}_k^\otimes , which is an ∞ -operad since the underlying simplicial operad is fibrant. In particular, objects in \mathbb{E}_k^\otimes are given by $\langle n \rangle$ for $n \in \mathbb{N}$, morphisms are given by points in

$$\text{Map}_{\mathbb{E}_k^\otimes}(\langle n \rangle, \langle m \rangle) = \coprod_{f: \langle n \rangle \rightarrow \langle m \rangle} \coprod_{j \in \langle m \rangle^\circ} \text{Rect}(\square^k \times \{1, \dots, n\}, \square^k),$$

and to give a 2-simplex with boundary as shown below is equivalent to giving a path from $F \circ E$ to G in $\text{Map}_{\mathbb{E}_k^\otimes}(\langle m \rangle, \langle k \rangle)$.

$$\begin{array}{ccc} & \langle m \rangle & \\ E \nearrow & & \searrow F \\ \langle n \rangle & \xrightarrow{G} & \langle k \rangle \end{array}$$

For $k = 1$ we obtain the \mathbb{E}_1^\otimes -operad, which is equivalent to the ∞ -operad Assoc^\otimes and governs homotopy associative algebras. For $k = 2$, we instead recover the ∞ -operadic version of the little 2-disks operad D_2 . In particular, an element in $\text{Mul}_{\mathbb{E}_1}(\langle n \rangle, \langle 1 \rangle)$ is given by a rectangular embedding of n copies of the interval $[0, 1]$ into the interval $[0, 1]$. An element in $\text{Mul}_{\mathbb{E}_2}(\langle n \rangle, \langle 1 \rangle)$ is given by a

rectangular embedding of n copies of the square $[0, 1] \times [0, 1]$ into the square $[0, 1] \times [0, 1]$.

Note that for $k \geq 0$, we have a homotopy equivalence $\text{Mul}_{\mathbb{E}_k}(\langle 2 \rangle, \langle 1 \rangle) \simeq S^{k-1}$ given by drawing a line between the middle points of the two copies of $[0, 1]^k$ inside $[0, 1]^k$ with direction given by going from the label 2 to the label 1, and finding the intersection of this line with the boundary of $[0, 1]^k$ in the positive direction. Then one can deform the boundary of $[0, 1]^k$ into an S^{k-1} and get the corresponding point there. We frequently use this homotopy equivalence as a convenient method to label morphisms in the little cubes operads. In particular, fix a homotopy inverse $S^0 \rightarrow \text{Mul}_{\mathbb{E}_1}(\langle 2 \rangle, \langle 1 \rangle)$ and a homotopy inverse $S^1 \rightarrow \text{Mul}_{\mathbb{E}_2}(\langle 2 \rangle, \langle 1 \rangle)$. Then we get two elements in $\text{Mul}_{\mathbb{E}_1}(\langle 2 \rangle, \langle 1 \rangle)$ named μ_0 and μ_π , and for every $t \in [0, 2\pi)$ we get an element $\mu_t \in \text{Mul}_{\mathbb{E}_2}(\langle 2 \rangle, \langle 1 \rangle)$.






FIGURE 1. The element $\mu_0 \in \text{Mul}_{\mathbb{E}_2}(\langle 2 \rangle, \langle 1 \rangle)$.

Recall that 2-morphisms in \mathbb{E}_k^\otimes are given by paths in the relevant hom-spaces. There are two such 2-morphisms in \mathbb{E}_2^\otimes that will play a special role in the subsequent discussion. On the one hand, for each $t \in [0, 2\pi)$, there is a 2-morphisms $\sigma_t \in \text{Map}_{\mathbb{E}_2^\otimes}(\langle 2 \rangle, \langle 1 \rangle)_1$ with boundary given by t and $t + \pi(\text{mod } 2\pi)$ that is represented by the braid



On the other hand, for each $t \in [0, 2\pi)$ there is a non-trivial 2-morphism $\gamma_t \in \text{Map}_{\mathbb{E}_2^\otimes}(\langle 2 \rangle, \langle 1 \rangle)_1$ between t and itself represented by the composition of braids



The classical Eckmann-Hilton argument shows that, in the 1-categorical case, algebra objects in the category of algebra objects yield commutative algebra objects. In particular, if (A, \cdot) is an associative algebra in a symmetric monoidal category \mathcal{C} and $\ast : (A, \cdot) \otimes (A, \cdot) \rightarrow (A, \cdot)$ endows (A, \cdot) with the structure of an associative algebra object in the $\text{Alg}_{\text{Assoc}}(\mathcal{C})$, then both operations \cdot and \ast agree and they are commutative. In this sense, an \mathbb{E}_1 -algebra inside the symmetric monoidal category of \mathbb{E}_1 -algebras in \mathcal{C} is the same as a commutative algebra inside \mathcal{C} , which in the 1-categorical case is the same as an \mathbb{E}_2 -algebra. In fact, this pattern continues for all the little k -cubes operads, as was shown by Dunn for topological operads and later by Lurie for ∞ -operads. To explain this, for $k, k' \geq 0$, define a topological functor

$$\rho : \mathbb{E}_k^{T, \otimes} \times \mathbb{E}_{k'}^{T, \otimes} \rightarrow \mathbb{E}_{k+k'}^{T, \otimes},$$

given on objects by $\rho(\langle m \rangle, \langle n \rangle) = \langle m \rangle \wedge \langle n \rangle$, and sending a pair of morphisms $(\alpha, \{f_j : \square^k \times \alpha^{-1}(\{j\}) \rightarrow \square^k\}_{j \in \langle n \rangle^\circ})$ and $(\beta, \{g_i : \square^{k'} \times \beta^{-1}(\{i\}) \rightarrow \square^{k'}\}_{i \in \langle n' \rangle^\circ})$ to

$$(\alpha \wedge \beta, \{f_j \times g_i : \square^{k+k'} \times \alpha^{-1}(\{j\}) \times \beta^{-1}(\{i\}) \rightarrow \square^{k+k'}\}_{j \in \langle n \rangle^\circ, i \in \langle n' \rangle^\circ}).$$

In order for this to make sense, we note that viewing a tuple $(j, i) \in \langle n \rangle^\circ \times \langle n' \rangle^\circ$ as an element of $\langle nn' \rangle^\circ$, we have $(\alpha \wedge \beta)^{-1}((j, i)) = \alpha^{-1}(\{j\}) \times \beta^{-1}(\{i\})$. This descends to a simplicial functor, and then taking the homotopy coherent nerve to a map of ∞ -categories $\rho : \mathbb{E}_k^\otimes \times \mathbb{E}_{k'}^\otimes \rightarrow \mathbb{E}_{k+k'}^\otimes$. By construction, the diagram

$$\begin{array}{ccc} \mathbb{E}_k^\otimes \times \mathbb{E}_{k'}^\otimes & \xrightarrow{\rho} & \mathbb{E}_{k+k'}^\otimes \\ \downarrow & & \downarrow \\ \text{Fin}_* \times \text{Fin}_* & \xrightarrow{\wedge} & \text{Fin}_* \end{array}$$

commutes, and clearly ρ sends pairs of inert morphisms to inert morphisms. Thus, ρ is a bifunctor of ∞ -operads.

Theorem 2.16 (Dunn Additivity Theorem, Theorem 5.1.2.2 [Lur17]). *The bifunctor $\rho : \mathbb{E}_k^\otimes \times \mathbb{E}_{k'}^\otimes \rightarrow \mathbb{E}_{k+k'}^\otimes$ exhibits the ∞ -operad $\mathbb{E}_{k+k'}^\otimes$ as a tensor product of \mathbb{E}_k^\otimes and $\mathbb{E}_{k'}^\otimes$.*

This means that for every symmetric monoidal ∞ -category \mathcal{C}^\otimes , precomposition with ρ determines an equivalence of ∞ -categories

$$\rho^* : \text{Alg}_{\mathbb{E}_{k+k'}}(\mathcal{C}) \rightarrow \text{Alg}_{\mathbb{E}_k}(\text{Alg}_{\mathbb{E}_{k'}}(\mathcal{C})).$$

In particular, for every \mathbb{E}_k -algebra A in $\mathbb{E}_{k'}$ -algebras in \mathcal{C}^\otimes , there exists a $\mathbb{E}_{k+k'}$ -algebra \tilde{A} in \mathcal{C}^\otimes such that $\tilde{A} \circ \rho$ is equivalent to A in the ∞ -category $\text{Alg}_{\mathbb{E}_k}(\text{Alg}_{\mathbb{E}_{k'}}(\mathcal{C}))$.

2.4. Deligne's Conjecture on Hochschild cochains. Let A be an associative k -algebra. The 1-category of left modules over the algebra $A \otimes A^{\text{op}}$ is isomorphic to the 1-category of A -bimodules. This is a monoidal category with tensor unit given by A viewed as a bimodule over itself.

Definition 2.17. The Hochschild complex of A with coefficients in an A -bimodule M is given by

$$C^*(A, M) = \mathbb{R}\text{Hom}_{A \otimes A^{\text{op}}}(A, M).$$

This is well-defined up to quasi-isomorphism. The Hochschild cohomology $\text{HH}^*(A, M)$ of A with coefficients in M is given by the cohomology of this complex. We will mainly be interested in the case $M = A$.

In particular, Hochschild cohomology groups can be viewed as the Ext-groups in the category of bimodules, and can be computed as the cohomology of

$$\text{Hom}_{A \otimes A^{\text{op}}}(P, A)$$

for any projective resolution $P \xrightarrow{\simeq} A$. In practice, one usually takes the Bar complex $B(A)$ of A to get the Hochschild cochain complex

$$\text{Hom}_{A \otimes A^{\text{op}}}(B(A), A) \cong \text{Hom}_k(A^{\otimes *}, A)$$

The Hochschild cohomology of A encodes the deformation theory of A as a bimodule over itself. In particular

$$\text{HH}^0(A, A) = \text{Hom}_{A \otimes A^{\text{op}}}(A, A) = Z(A).$$

In 1962, M. Gerstenhaber [Ger63] noticed that Hochschild cohomology is equipped with a shifted Lie bracket $[-, -]_G$ and a commutative product \smile such that $[f, -]_G$ is a derivation for the product. Such a structure is now called a Gerstenhaber algebra. We denote the dg operad governing Gerstenhaber algebras by Ger . By a result of F. Cohen, the Gerstenhaber operad is the singular homology of the topological operad of little 2-disks D_2 . In particular, if A is an algebra over $C_*(D_2)$, then $H_*(A)$ is an algebra over Ger . This inspired P. Deligne in a 1993 letter to make the following conjecture.

Conjecture 2.18 (Deligne's Conjecture). *The Hochschild cochain complex of an associative algebra A is an algebra over the chains on little 2-disks operad in such a way that the induced Gerstenhaber algebra structure on cohomology recovers Gerstenhaber's original one.*

In his letter, Deligne explained that the $C_*(D_2)$ -algebra structure on the cochain level should come from the two multiplications on $\mathbb{R}\text{Hom}_{A \otimes A^{\text{op}}}(A, A)$ induced by A being a monoidal unit. In particular, we have an "inner multiplication" coming from the bialgebra structure of A in the bimodule category. Let $\text{End}(A) := \text{Hom}_{A \otimes A^{\text{op}}}(A, A)$. Then we get

$$\begin{aligned} \text{End}(A) \otimes_A \text{End}(A) \otimes_A A &\xrightarrow{\simeq} \text{End}(A) \otimes_A \text{End}(A) \otimes_A A \otimes_A A \\ &\cong \text{End}(A) \otimes_A A \otimes_A \text{End}(A) \otimes_A A \rightarrow A \otimes_A A \xrightarrow{\simeq} A \end{aligned}$$

inducing a multiplication $\text{End}(A) \otimes_A \text{End}(A) \rightarrow \text{End}(A)$. We also have an "outer multiplication" given by composition.

Multiple different proofs have been given that this is in fact true, see for example [Tam98], [Vor00], [MS02]. In particular, D. Tamarkin in [Tam98] constructed a map $\Psi_T : \mathcal{G}er_\infty \rightarrow \mathcal{B}races$ from the operad of homotopy Gerstenhaber algebras to the Braces-operad, and also proved a formality result for the Gerstenhaber operad. Since $C^*(A, A)$ is canonically a Braces-algebra, this solves Deligne's conjecture. Notably, Tamarkin's map Ψ_T depends on the choice of a Drinfeld associator.

There is a global version of Deligne's Conjecture. Let X be a separated finite type scheme over \mathbb{k} . Then the Hochschild cohomology of X should be a global version of the above definitions, such that if $X = \text{Spec}(A)$ we recover definition 2.17. Historically, there have been multiple proposed definitions, for example by Grothendieck-Loday, Gerstenhaber-Schack and Swan. The perhaps most straight-forward generalization from the affine case was given by Swan, who defined the Hochschild cohomology of X to be

$$\text{HH}_S^*(X) = \text{Ext}_{\mathcal{O}_X \times_{\mathbb{k}} X}^*(\Delta_* \mathcal{O}_X, \Delta_* \mathcal{O}_X).$$

Unfortunately, this does not carry a Gerstenhaber bracket. For the case that X is smooth, Kontsevich gave an alternative definition, which does carry the structure of a Gerstenhaber algebra. He defined the complex of polydifferential operators on a regular algebra A to be the subcomplex

$$D_{\text{poly}}(A) \subseteq C^*(A, A)$$

of maps $f : A^{\otimes n} \rightarrow A$ that are differential operators in each variable separately. These glue together to yield the sheaf of polydifferential operators on X

$$\mathcal{D}_{\text{poly}}(X)(\text{Spec}(A)) = D_{\text{poly}}(A).$$

Then the Hochschild cohomology of X is the hypercohomology of the sheaf of polydifferential operators,

$$\text{HH}_K^*(X) = \mathbb{H}^*(X, \mathcal{D}_{\text{poly}}(X)).$$

Since $\mathcal{D}_{\text{poly}}(X)$ is a sheaf of homotopy Gerstenhaber algebras by Tamarkin's solution of the Deligne conjecture, the hypercohomology inherits a Gerstenhaber algebra structure.

3. THE BRACKET OPERATION ON 2-ALGEBRAS

3.1. The bracket operation of an \mathbb{E}_2 -algebra. Consider the topological operad \mathbb{E}_2 of little 2-disks and its corresponding dg-operad $C_*(\mathbb{E}_2)$. If $C_*(\mathbb{E}_2) \rightarrow \text{End}(A)$ is an operad algebra in the category of chain complexes over \mathbb{k} , we have an action of the 2-ary operation space

$$C_*(\mathbb{E}_2(2)) \otimes A^{\otimes 2} \rightarrow A,$$

and recalling that $\mathbb{E}_2(2) \simeq S^1$, taking homology yields a map

$$H_*(S^1) \otimes H_*(A)^{\otimes 2} \rightarrow H_*(A).$$

Since $H_*(S^1) \cong \mathbb{Z}[p] \oplus \mathbb{Z}[\gamma]$ for some choice of basepoint $p \in S^1$ and generating loop $\gamma : [0, 1] \rightarrow S^1$, this yields two 2-ary operations on $H_*(A)$; one of degree 0 induced by $[p]$

$$\smile : H_*(A) \otimes H_*(A) \rightarrow H_*(A)$$

and one of degree 1 induced by $[\gamma]$

$$[\cdot, \cdot] : H_*(A) \otimes H_*(A) \rightarrow H_*(A)[-1].$$

These two operations make $H_*(A)$ into a Gerstenhaber algebra. This discussion in particular shows that the bracket operation is induced by the chain level operation $A^{\otimes 2} \rightarrow A$ corresponding to a choice of generating loop γ of the homology of S^1 .

Definition 3.19. Let \mathcal{C}^\otimes be a symmetric monoidal ∞ -category and let $A : \mathbb{E}_2^\otimes \rightarrow \mathcal{C}^\otimes$ be an \mathbb{E}_2 -algebra in \mathcal{C}^\otimes . Then we call the image under A of $\gamma_t \in \text{Map}_{\mathbb{E}_2^\otimes}(\langle 2 \rangle, \langle 1 \rangle)_1$ the bracket operation of A at $m_t \in \text{Map}_{\mathbb{E}_2^\otimes}(\langle 2 \rangle, \langle 1 \rangle)_0$.

Let $A : \mathbb{E}_2^\otimes \rightarrow N_{\text{dg}}(C^\circ)^\otimes$ be an \mathbb{E}_2 -algebra in a symmetric monoidal dg model category C . Then we get an induced map

$$\mathbb{E}_2(2) = \text{Map}_{\mathbb{E}_2^\otimes}^\alpha(\langle 2 \rangle, \langle 1 \rangle) \rightarrow \text{Map}_{N_{\text{dg}}(C^\circ)^\otimes}(A(\langle 2 \rangle), A(\langle 1 \rangle)).$$

Let $A = A(\langle 1 \rangle) \in C^\circ$. Then we have a homotopy equivalence

$$\text{Map}_{N_{\text{dg}}(C^\circ)^\otimes}(A(\langle 2 \rangle), A(\langle 1 \rangle)) \simeq \text{Map}_{N_{\text{dg}}(C^\circ)}(A^{\otimes 2}, A).$$

Hence we get a map (well-defined up to homotopy)

$$\mathbb{E}_2(2) \rightarrow \text{Map}_{N_{\text{dg}}(C^\circ)}(A^{\otimes 2}, A) \simeq \text{DK}\tau_{\geq 0} \text{Map}_C(A^{\otimes 2}, A),$$

and therefore taking homology we get maps

$$(\mathcal{G}er(2))_n \cong H_n(\mathbb{E}_2(2)) \rightarrow \text{Hom}_{\mathcal{D}(C)}(A^{\otimes 2}, A[n]).$$

This procedure yields a Gerstenhaber algebra structure on A in the derived category of C . We see that the bracket of this Gerstenhaber algebra is indeed given by the image of γ_0 .

3.2. The bracket operation of a 2-algebra. If $A : \mathbb{E}_1^\otimes \times \mathbb{E}_1^\otimes \rightarrow \mathcal{C}^\otimes$ is a 2-algebra in \mathcal{C}^\otimes , the Dunn Additivity Theorem 2.16 tells us that there exists an \mathbb{E}_2 -algebra $\tilde{A} : \mathbb{E}_2^\otimes \rightarrow \mathcal{C}^\otimes$ such that the restriction of \tilde{A} along $\rho : \mathbb{E}_1^\otimes \times \mathbb{E}_1^\otimes \rightarrow \mathbb{E}_2^\otimes$ is equivalent to A in the category of bifunctors. Fixing such an \mathbb{E}_2^\otimes -algebra, we can ask whether it is possible to express the bracket operations $\tilde{A}(\gamma_t)$ in terms of the original 2-algebra A .

Denote by $A \in \mathcal{C}$ the image $A(\langle 1 \rangle, \langle 1 \rangle)$, and let $\mu = \mu_0 \in \text{Hom}_{\mathbb{E}_1^\otimes}(\langle 2 \rangle, \langle 1 \rangle)_0$ be the element

$$\begin{array}{c} 2 \quad 1 \\ | \quad | \\ \hline \end{array}.$$

We fix coCartesian lifts of inert maps in \mathbb{E}_1^\otimes by using the correct enumerations of the full intervals. By abuse of notation, we denote those lifts by the inert map in $\mathcal{F}in_*$ they lift. The key observation in expressing the bracket operations in terms of the original 2-algebra is given by the following theorem.

Theorem 3.20. *The images under A of the 2-simplices in $\mathbb{E}_1^\otimes \times \mathbb{E}_1^\otimes$*

$$\begin{array}{ccc} (\langle 2 \rangle, \langle 2 \rangle) & \xrightarrow{(\text{id}_{\langle 2 \rangle}, \mu)} & (\langle 2 \rangle, \langle 1 \rangle) \\ (\mu, \text{id}_{\langle 2 \rangle}) \downarrow & & \downarrow (\mu, \text{id}_{\langle 1 \rangle}) \\ (\langle 1 \rangle, \langle 2 \rangle) & \xrightarrow{(\text{id}_{\langle 1 \rangle}, \mu)} & (\langle 1 \rangle, \langle 1 \rangle) \end{array}$$

$$\begin{array}{ccc}
 (\langle 1 \rangle, \langle 2 \rangle) & & \\
 (\text{id}_{\langle 1 \rangle}, (2,3)) \downarrow & \searrow (\text{id}_{\langle 1 \rangle}, \text{id}_{\langle 2 \rangle}) & \\
 (\langle 1 \rangle, \langle 4 \rangle) & \xrightarrow{(\text{id}_{\langle 1 \rangle}, (\mu, \mu))} & (\langle 1 \rangle, \langle 2 \rangle) \\
 (\langle 2 \rangle, \langle 1 \rangle) & \xrightarrow{((2,3), \text{id}_{\langle 1 \rangle})} & (\langle 4 \rangle, \langle 1 \rangle) \\
 & \searrow (\tau, \text{id}_{\langle 1 \rangle}) & \downarrow ((\mu, \mu), \text{id}_{\langle 1 \rangle}) \\
 & & (\langle 2 \rangle, \langle 1 \rangle)
 \end{array}$$

induce a 2-simplex in \mathcal{C}^\otimes (up to homotopy)

$$\begin{array}{ccccc}
 (A, A) & & \xrightarrow{\text{id}_{(A, A)}} & & (A, A) \\
 & \searrow \iota_{2,3} & & \searrow (m_2, m_2) & \\
 & & (A, A, A, A) & \xrightarrow{(m_2, m_2)} & (A, A) \\
 & \searrow \tau & \downarrow (m_1, m_1) \circ \tau_{2,3} & & \downarrow m_1 \\
 & & (A, A) & \xrightarrow{m_2} & A
 \end{array}$$

whose homotopy class is identified under the isomorphism $\tilde{A} \circ \rho \simeq A$ with the image under \tilde{A} of the half twist between $t = 0$ and $t = \pi$ in $\text{Hom}_{\mathbb{E}_2^\otimes}(\langle 2 \rangle, \langle 1 \rangle)_1$.

Proof. First check that the 2-simplices in $\mathbb{E}_1^\otimes \times \mathbb{E}_1^\otimes$ indeed induce composable 2-simplices in \mathcal{C}^\otimes with the depicted boundaries. We get diagrams

$$\begin{array}{ccccc}
 & & (A, A, A, A) & \xrightarrow{\quad} & (A, A) \\
 & \nearrow \simeq & \downarrow & \nearrow \simeq & \downarrow \\
 A(\langle 2 \rangle, \langle 2 \rangle) & \xrightarrow{\quad} & A(\langle 2 \rangle, \langle 1 \rangle) & & \\
 \downarrow & & \downarrow & & \downarrow \\
 & \nearrow \simeq & (A, A) & \xrightarrow{\quad} & A \\
 & \searrow \simeq & \downarrow & \searrow \simeq & \\
 A(\langle 1 \rangle, \langle 2 \rangle) & \xrightarrow{\quad} & A(\langle 1 \rangle, \langle 1 \rangle) & &
 \end{array}$$

$$\begin{array}{ccccc}
 & & (A, A) & & \\
 & \nearrow \simeq & \downarrow & \searrow & \\
 A(\langle 1 \rangle, \langle 2 \rangle) & & (A, A, A, A) & \xrightarrow{\quad} & (A, A) \\
 \downarrow & \nearrow \simeq & \downarrow & \searrow \simeq & \\
 A(\langle 1 \rangle, \langle 4 \rangle) & \xrightarrow{\quad} & A(\langle 1 \rangle, \langle 2 \rangle) & &
 \end{array}$$

$$\begin{array}{ccccc}
 & & (A, A) & \longrightarrow & (A, A, A, A) \\
 & \nearrow \simeq & & \searrow \simeq & \downarrow \\
 A(\langle 2 \rangle, \langle 1 \rangle) & \longrightarrow & A(\langle 4 \rangle, \langle 1 \rangle) & & (A, A) \\
 & \searrow & \downarrow & \nearrow \simeq & \\
 & & A(\langle 2 \rangle, \langle 1 \rangle) & &
 \end{array}$$

so it suffices to show that these fit together into the depicted 2-simplex. To this end we check that the maps $(A, A, A, A) \rightarrow (A, A)$ in the square agree with the respective maps $(A, A, A, A) \rightarrow (A, A)$ in the triangles, and similarly for the two maps $(A, A) \rightarrow (A, A, A, A)$ in the different triangles. To this end note that it suffices to show that each of those pairs of maps agrees after postcomposition with coCartesian lifts of the ρ^i . Consider first the maps induced by $(\langle 2 \rangle, \langle 2 \rangle) \xrightarrow{\text{id}_{\langle 2 \rangle}, \mu} (\langle 2 \rangle, \langle 1 \rangle)$ and $(\langle 1 \rangle, \langle 4 \rangle) \xrightarrow{\text{id}_{\langle 1 \rangle}, (\mu, \mu)} (\langle 1 \rangle, \langle 2 \rangle)$. We have a factorization

$$\begin{array}{ccccc}
 (\langle 2 \rangle, \langle 2 \rangle) & \xrightarrow{\text{id}, \mu} & (\langle 2 \rangle, \langle 1 \rangle) & \xrightarrow{\rho^i, \text{id}} & (\langle 1 \rangle, \langle 1 \rangle) \\
 & \searrow \rho^i, \text{id} & & \nearrow \text{id}, \mu & \\
 & & (\langle 1 \rangle, \langle 2 \rangle) & &
 \end{array}$$

in $\mathbb{E}_1^\otimes \times \mathbb{E}_1^\otimes$, where the lower left hand side map lies over the inert map $1, 2, *, * : \langle 4 \rangle \rightarrow \langle 2 \rangle$ if $i = 1$ and over the inert map $*, *, 1, 2$ if $i = 2$. Since inert pairs are sent to inert maps by A , this diagram maps to

$$\begin{array}{ccccc}
 (A, A, A, A) & \xrightarrow{A(\text{id}, \mu)} & (A, A) & \xrightarrow{\rho^i} & A \\
 & \searrow & & \nearrow m_2 & \\
 & & (A, A) & &
 \end{array}$$

where $(A, A, A, A) \rightarrow (A, A)$ is an inert lift. Now on the other hand, we also have a factorization


$$\begin{array}{ccccc}
 (\langle 1 \rangle, \langle 4 \rangle) & \xrightarrow{\text{id}, (\mu, \mu)} & (\langle 1 \rangle, \langle 2 \rangle) & \xrightarrow{\text{id}, \rho^i} & (\langle 1 \rangle, \langle 1 \rangle) \\
 & \searrow \text{id}, f_i & & \nearrow \text{id}, \mu & \\
 & & (\langle 1 \rangle, \langle 2 \rangle) & &
 \end{array}$$

where f_i is $1, 2, *, *$ if $i = 1$ and $*, *, 1, 2$ if $i = 2$. This again is sent by A to

$$\begin{array}{ccccc}
 (A, A, A, A) & \xrightarrow{A(\text{id}, (\mu, \mu))} & (A, A) & \xrightarrow{\rho^i} & A \\
 & \searrow & & \nearrow m_2 & \\
 & & (A, A) & &
 \end{array}$$




showing that our two maps $(A, A, A, A) \rightarrow (A, A)$ indeed agree up to homotopy. An analogous analysis can be carried out with the other two maps $(A, A, A, A) \rightarrow (A, A)$. For the two inclusions $(\langle 1 \rangle, \langle 2 \rangle) \rightarrow (\langle 1 \rangle, \langle 4 \rangle)$ and $(\langle 2 \rangle, \langle 1 \rangle) \rightarrow (\langle 4 \rangle, \langle 1 \rangle)$, it suffices to show that the two unit maps coming from $(\langle 1 \rangle, \langle 0 \rangle) \rightarrow (\langle 1 \rangle, \langle 1 \rangle)$ and $(\langle 0 \rangle, \langle 1 \rangle) \rightarrow (\langle 1 \rangle, \langle 1 \rangle)$ agree. To this end, note that both $(\langle 0 \rangle, \langle 0 \rangle) \rightarrow$

$(\langle 1 \rangle, \langle 0 \rangle)$ and $(\langle 0 \rangle, \langle 0 \rangle) \rightarrow (\langle 0 \rangle, \langle 1 \rangle)$ are sent to the identity on the empty tuple by A , up to homotopy, since $\mathcal{C}_{\langle 0 \rangle}^\otimes$ is contractible. Then the image of the diagram



under A shows that the two unit maps are homotopic.


We now show that the depicted 2-simplex in \mathcal{C}^\otimes indeed corresponds to the image of the half twist under \tilde{A} . To this end, examine the images of the three 2-simplices in $\mathbb{E}_1^\otimes \times \mathbb{E}_1^\otimes$ under $\rho : \mathbb{E}_1^\otimes \times \mathbb{E}_1^\otimes \rightarrow \mathbb{E}_2^\otimes$. We get




Note that the paths forming the fillings of the two triangles in the square diagram are both the constant path at

3	1
4	2

while the fillings of the triangle diagrams are given by continuously enlarging the respective rectangles. Composing those in the simplicial category $\mathbb{E}_2^{\Delta, \otimes}$ we hence get the half twist




Applying \tilde{A} to the above simplices induces a diagram of 2-simplices in \mathcal{C}^{\otimes}



and by construction the isomorphism between $\tilde{A} \circ \rho$ and A identifies this diagram with the one in the statement. Since maps between ∞ -categories respect composition, this proves the claim. \square

We can repeat the analysis in this theorem for the other parts of the classical Eckmann-Hilton argument. In particular, we get representations of the image under \tilde{A} of all four different parts of the double twist in terms of compositions of 2-simplices in the image of A . As a corollary, we obtain

Corollary 3.21. *The homotopy class of the bracket on \tilde{A} at $t = 0$ can be produced as a composition of the following 2-simplices in \mathcal{C}^{\otimes}*



$$\begin{array}{c}
 \begin{array}{ccc}
 (A, A) & \xrightarrow{\text{id}} & (A, A) \\
 \downarrow \iota_{1,4} & \searrow (m_2^{\text{op}}, m_2^{\text{op}}) & \downarrow m_1^{\text{op}} \\
 (A, A, A, A) & \xrightarrow{(m_2^{\text{op}}, m_2^{\text{op}})} & (A, A) \\
 \downarrow (m_1^{\text{op}}, m_1^{\text{op}}) \circ \tau_{2,3} & & \downarrow m_1^{\text{op}} \\
 (A, A) & \xrightarrow{m_2^{\text{op}}} & A
 \end{array} \\
 \downarrow \text{id} \\
 \begin{array}{ccc}
 (A, A) & \xrightarrow{\text{id}} & (A, A) \\
 \downarrow \iota_{2,3} & \searrow (m_2^{\text{op}}, m_2^{\text{op}}) & \downarrow m_1^{\text{op}} \\
 (A, A, A, A) & \xrightarrow{(m_2^{\text{op}}, m_2^{\text{op}})} & (A, A) \\
 \downarrow (m_1^{\text{op}}, m_1^{\text{op}}) \circ \tau_{2,3} & & \downarrow m_1^{\text{op}} \\
 (A, A) & \xrightarrow{m_2^{\text{op}}} & A
 \end{array} \\
 \downarrow \tau \\
 \begin{array}{ccc}
 (A, A) & \xrightarrow{\text{id}} & (A, A) \\
 \downarrow \iota_{1,4} & \searrow (m_2, m_2) & \downarrow m_1 \\
 (A, A, A, A) & \xrightarrow{(m_2, m_2)} & (A, A) \\
 \downarrow (m_1, m_1) \circ \tau_{2,3} & & \downarrow m_1 \\
 (A, A) & \xrightarrow{m_2} & A
 \end{array}
 \end{array}$$

Corollary 3.22. *If C^\otimes is the dg nerve of a symmetric monoidal dg model category C , then the bracket at m_0 on \tilde{A} is given by the chain homotopy*

$$\gamma_0 \simeq h\iota_{2,3} + h^{\text{op}}\iota_{1,4} + h^{\text{op}}\iota_{2,3} + h\iota_{1,4}.$$

Here h and h^{op} are the chain homotopies corresponding to the image of the square diagram in $\mathbb{E}_1 \times \mathbb{E}_1$ for the multiplications and their opposite multiplications respectively.

Proof. Note first that by construction of the symmetric monoidal structure on \mathcal{C} , a morphism $\underbrace{(A, \dots, A)}_{m \text{ times}} \rightarrow \underbrace{(A, \dots, A)}_{n \text{ times}}$ in \mathcal{C}^\otimes corresponds to a map $A^{\otimes m} \rightarrow A^{\otimes n}$ in C which is unique up to chain homotopy. Similarly, a 2-simplex between such maps corresponds to a chain homotopy between the corresponding maps in C . Fix such maps in C corresponding to all the involved diagrams. Horizontal composition of maps is strictly defined in dg categories, and hence we have well defined whiskering compositions $h\iota_{2,3}$, $h\iota_{1,4}$, $h^{\text{op}}\iota_{2,3}$ and $h^{\text{op}}\iota_{1,4}$. Finally, note that horizontal composition of chain homotopies is given by addition. \square

4. RECOVERING THE HOCHSCHILD COMPLEX AS \mathbb{E}_1 -CENTER

Often the Hochschild cohomology of a \mathbb{k} -algebra is called its "derived center". In this section we will show that this statement is true in a very precise sense. Namely, the Hochschild complex is the \mathbb{E}_1 -center in the derived ∞ -category of chain complexes.

4.1. The dg nerve. If C is a dg category, denote by C_0 the underlying 1-category. If C is a dg model category, we view C_0 as a model category.

Theorem 4.23. *Let C be a dg category that is tensored over the category of chain complexes of \mathbb{k} -modules, let C' be a full dg subcategory, and let W be a collection of morphisms in C' that are isomorphisms in the homotopy category. Assume that the following conditions are satisfied:*

- *Every isomorphism in C' belongs to W .*
- *The set W satisfies the 2-out-of-3 property.*
- *For all $X \in C'$, we also have $N_*(\Delta^1) \otimes X \in C'$.*
- *For each $X \in C'$, the map $N_*(\Delta^1) \otimes X \rightarrow X$ induced by the map $[1] \rightarrow [0]$ belongs to W .*

Then the canonical map $\theta : N(C'_0) \rightarrow N_{dg}(C')$ induces an equivalence of ∞ -categories $\theta' : N(C'_0)[W^{-1}] \simeq N_{dg}(C')$.

Proof. The above conditions are exactly what is needed to repeat the proof of [Lur17, Proposition 1.3.4.5] replacing $\text{Ch}(\mathcal{A})$ with \mathcal{C} and $\text{Ch}(\mathcal{A})'$ with C' . \square

Corollary 4.24. *Let C be a dg model category, and let C° be the full subcategory on bifibrant objects. Then the map $N(C_0^\circ) \rightarrow N_{dg}(C^\circ)$ exhibits the dg nerve as the underlying ∞ -category of C_0 .*

Proof. Let C_Δ be the simplicial category obtained from \mathcal{C} . Since $N(C_0^\circ)[W^{-1}] \simeq N(C_0^\circ)[W^{-1}]$ it suffices to take $C' = C^\circ$ above and W the set of homotopy equivalences. In a dg model category, left homotopy between bifibrant objects agrees with chain homotopy in the dg sense, i.e. a homotopy is a map $h : N(\Delta^1) \otimes X \rightarrow Y$ with the correct restrictions to $\{0\}$ and $\{1\}$. In particular,

$$\begin{aligned} \text{Hom}_{C_0}(N_*(\Delta^1) \otimes X, Y) &\cong \text{Hom}_{\text{Ch}(\mathbb{k})}(N_*(\Delta^1), \text{Map}_C(X, Y)) \\ &\cong \text{Hom}_{\text{Ch}_{\geq 0}(\mathbb{k})}(N_*(\Delta^1), \tau_{\geq 0} \text{Map}_C(X, Y)) \\ &\cong \text{Hom}_{\text{sSet}}(\Delta^1, \text{DK}_\bullet \tau_{\geq 0} \text{Map}_C(X, Y)) \\ &\cong \text{Map}_{C_\Delta}(X, Y)_1 \end{aligned}$$

so chain homotopies correspond to 1-chains of the mapping complex of C . One checks directly that the diagram making $h : N_*(\Delta^1) \otimes X \rightarrow Y$ into a homotopy between $f, g \in \text{Hom}_{C_0}(X, Y)$ forces the corresponding 1-chain $z \in \text{Map}_C(X, Y)_1$ to satisfy $dz = f - g$. This shows that homotopy equivalences in C_0° become isomorphisms in the homotopy category hC_Δ , which is isomorphic to the homotopy category hC . Clearly every isomorphism is a homotopy equivalence and the set of homotopy equivalences satisfies 2-out-of-3. By assumption, the map $\otimes : \text{Ch}(\mathbb{k}) \times \mathcal{C} \rightarrow \mathcal{C}$ is a left Quillen bifunctor. The complex $N_*(\Delta^1)$ is bounded below and therefore cofibrant, and hence $N_*(\Delta^1) \otimes -$ preserves cofibrant objects. At the same time, C is also powered over $\text{Ch}(\mathbb{k})$ and hence the functor $N_*(\Delta^1) \otimes -$ is right adjoint to the left Quillen bifunctor $N_*(\Delta^1)^\vee \otimes -$. In particular, it preserves fibrant objects. This shows that $N_*(\Delta^1) \otimes X \in C^\circ$. Finally, note that $d_0 : \mathbb{k} \rightarrow N_*(\Delta^1)$ is a trivial cofibration in $\text{Ch}(\mathbb{k})$, and thus if $X \in C^\circ$, the map $X \cong \mathbb{k} \otimes X \rightarrow N_*(\Delta^1) \otimes X$ is again a trivial cofibration. Now the map $N_*(\Delta^1) \rightarrow \mathbb{k}$ is a left inverse to d_0 and in particular

$$X \rightarrow N_*(\Delta^1) \otimes X \rightarrow X$$

is the identity on X and thus a weak equivalence. By 2-out-of-3, this means that $N_*(\Delta^1) \otimes X \rightarrow X$ must be a weak equivalence. \square

Remark 4.25. Note that for any dg category C , we have an equivalence of ∞ -categories $N_{\text{hc}}(C_\Delta) \rightarrow N_{\text{dg}}(C)$. For simplicial model categories, the homotopy coherent nerve of the bifibrant objects is

always the ∞ -category underlying the model category, but C_Δ is not tensored and cotensored over \mathbf{sSet} and therefore does not satisfy the requirements of this theorem. The above corollary then shows that we get this relationship between the homotopy coherent nerve and the model category regardless.

If C is a (symmetric) monoidal model category, then [Lur17, Example 4.1.7.6] shows that $N(C^c)[W^{-1}]$ is a (symmetric) monoidal ∞ -category. If C is also a simplicial model category and the (symmetric) monoidal structure is compatible with the simplicial enrichment, then [Lur17, Corollary 4.1.7.16] shows that the (symmetric) monoidal structure on this ∞ -category is given by $N_{\text{hc}}((C^\circ)^\otimes)$, and in fact one readily checks that the same hold if C is just weakly simplicial in the same sense as above. We would like to use this to argue that the dg nerve of a (symmetric) monoidal dg model category C is a (symmetric) monoidal ∞ -category and also presents the (symmetric) monoidal structure of $N(C^c)[W^{-1}]$, but unfortunately the Dold-Kan functor is not symmetric, and thus does not send operads to operads. Nevertheless, it is homotopy symmetric lax monoidal, and in fact V. Hinich proved the following.

Proposition 4.26 ([Hin15], Theorem 3.2.3). *The dg nerve $N_{\text{dg}} : N(\text{Cat}_{\text{dg}})[W_{\text{dg}}^{-1}] \rightarrow N(\text{Cat}_\Delta)[W_\Delta^{-1}] \simeq \text{Cat}_\infty$ from the symmetric monoidal ∞ -category of dg-categories to the symmetric monoidal ∞ -category of ∞ -categories is lax symmetric monoidal. In particular, it is a morphism of ∞ -operads. It thus induces a map from the ∞ -category of symmetric monoidal dg-categories to the ∞ -category of symmetric monoidal ∞ -categories.*

Corollary 4.27. *The dg nerve induces a map $\text{Alg}_{\mathcal{LM}^\otimes}(N(\text{Cat}_{\text{dg}})[W_{\text{dg}}^{-1}]) \rightarrow \text{Alg}_{\mathcal{LM}^\otimes}(\text{Cat}_\infty)$. By 4.28, this means that a dg category left tensored over a monoidal dg category yields an ∞ -category left tensored over a monoidal ∞ -category.*

4.2. Rectification of algebras over an ∞ -operad. Let \mathcal{O} be a topological operad. Then we get a dg operad $C_*(\mathcal{O})$ by applying the singular chains functor with coefficients in our field k . We want to view algebras in a symmetric monoidal dg model category C over this dg operad as algebras over the ∞ -operad $N^\otimes(\text{Sing}_\bullet \mathcal{O})$ in the symmetric monoidal ∞ -category $N_{\text{dg}}(C^\circ)$. We generalize the rectification results of Hinich [Hin15] and of D. Pavlov and J. Scholbach [PS18a].

Let C be a symmetric monoidal model category that is enriched over the projective model category of chain complexes over k . Suppose further that C is cofibrantly generated and symmetrically flat, and that $C_*(\mathcal{O})$ is admissible and well-pointed in C , and that it admits a lax monoidal fibrant replacement functor. The construction in [Hin15, Section 4.2] generalizes directly to give a functor

$$\phi : N(\text{Alg}_{C_*(\mathcal{O})}(C)^c) \rightarrow \text{Alg}_{N^\otimes(\mathcal{O})}(N_{\text{dg}}(C^\circ))$$

that carries weak equivalences to equivalences, and therefore yields a comparison map

$$\Phi : N(\text{Alg}_{C_*(\mathcal{O})}(C)^c)[W_{\text{Alg}_{C_*(\mathcal{O})}(C)}^{-1}] \rightarrow \text{Alg}_{N^\otimes(\mathcal{O})}(N_{\text{dg}}(C^\circ)).$$

We want to show that this functor is an equivalence of ∞ -categories.

Theorem 4.28. *Let \mathcal{O} and C as above. Then Φ is an equivalence of ∞ -categories.*

Proof. We use Corollary 4.7.3.16 in [Lur17]. Following the reasoning in [PS18a, Theorem 7.11], we may assume that $C_*(\mathcal{O})$ is Σ -cofibrant in C . Now consider

$$\begin{array}{ccc}
 N(\mathrm{Alg}_{C_*(\mathcal{O})}(C)^c)[W_{\mathrm{Alg}_{C_*(\mathcal{O})}(C)}^{-1}] & \xrightarrow{\Phi} & \mathrm{Alg}_{N^\otimes(\mathcal{O})}(N(C^c)[W^{-1}]) \\
 & \searrow G \quad \swarrow G' & \\
 & (N(C^c)[W^{-1}])^{[\mathcal{O}]} &
 \end{array}$$

Steps (a)-(c) in [Lur17, Corollary 4.7.3.16] can be proven exactly like in [Lur17, Theorem 4.5.4.7], by just replacing the commutative operad by \mathcal{O} . For step (d), it is clear that G is conservative since the weak equivalences in $\mathrm{Alg}_{C_*(\mathcal{O})}(C)$ are transferred from the ones in C via the forgetful functor. To show that G preserves geometric realization of simplicial objects, it suffices to show that it preserves homotopy sifted colimits. This is shown in [PS18a, Proposition 7.9]. Finally, we need to show that the canonical transformation $G' \circ F' \rightarrow G \circ F$ is an equivalence, where F and F' are the left adjoints of G and G' respectively. This boils down to showing that for any cofibrant object $X \in C^{[\mathcal{O}]}$, the strict free $C_*(\mathcal{O})$ -algebra generated by X is also a free $N^\otimes(\mathcal{O})$ -algebra in the sense of [Lur17, Definition 3.1.3.1]. In [Hin15, Lemma 4.3.4], Hinich has described an analog of [Lur17, Proposition 3.1.3.13] for free algebras generated by objects of different colors. The arguments described there go through if we replace the category of chain complexes by C . This finishes the argument. \square

Proposition 4.29. *Let $A : \mathbb{E}_2^\otimes \rightarrow N_{\mathrm{dg}}(C^\circ)^\otimes$ be an \mathbb{E}_2 -algebra in the dg nerve of a symmetric monoidal dg model category. Let A^{str} be a homotopy preimage of A under Φ . Without loss of generality, assume that the underlying object of A^{str} is fibrant. Then there is a chain homotopy h*

$$\begin{array}{ccc}
 A^{\mathrm{str} \otimes 2} & \xrightarrow{A^{\mathrm{str}}(m_0)} & A^{\mathrm{str}} \\
 \simeq \downarrow & \Downarrow h & \downarrow \simeq \\
 A^{\otimes 2} & \xrightarrow{A(m_0)} & A
 \end{array}$$

and a chain homotopy of chain homotopies filling the cylinder

$$\begin{array}{ccc}
 & \xrightarrow{A^{\mathrm{str}}(m_0)} & \\
 A^{\mathrm{str} \otimes 2} & \Downarrow A^{\mathrm{str}}(\gamma_0) & A^{\mathrm{str}} \\
 \simeq \downarrow & \xrightarrow{A^{\mathrm{str}}(m_0)} & \downarrow \simeq \\
 & \xrightarrow{A(m_0)} & \\
 A^{\otimes 2} & \Downarrow A(\gamma_0) & A \\
 & \xrightarrow{A(m_0)} &
 \end{array}$$

In particular, the induced Gerstenhaber algebra from A^{str} in the derived category agrees with the one constructed directly from A in Section 3.1.

Proof. There is a natural isomorphism $\eta : \Delta^1 \times \mathbb{E}_2^\otimes \rightarrow N_{\mathrm{dg}}(C^\circ)^\otimes$ between $\Phi(A^{\mathrm{str}})$ and A . In particular, we get isomorphisms $\Phi(A^{\mathrm{str}})(\langle n \rangle) \rightarrow A(\langle n \rangle)$, that correspond to isomorphisms $\Phi(A^{\mathrm{str}})^{\otimes n} \rightarrow A^{\otimes n}$. All the (higher) chain homotopies in the statement now correspond to the evaluation of η at the appropriate simplices in $\Delta^1 \times \mathbb{E}_2^\otimes$. For example, the 1-simplex (e, m_0) yields a map $\Phi(A^{\mathrm{str}})^{\otimes 2} \rightarrow A$ in C , and the 2-simplex $(s_0 e, s_1 m_0)$ yields a chain homotopy making $\eta(e, m_0)$ into a composition of $\Phi(A^{\mathrm{str}})(m_0)$ and $\eta(e, s_0 \langle 1 \rangle)$. \square

4.3. The Hochschild complex. Let A be an associative \mathbb{k} -algebra. We want to use the equivalence constructed in Section 4.2 to view A as an \mathbb{E}_1 -algebra in the symmetric monoidal ∞ -category $\mathcal{C}^\otimes := N_{\text{dg}}(\text{Ch}(\mathbb{k}))^\otimes$. To this end, let $\phi : C_*(\mathbb{E}_1) \xrightarrow{\sim} \mathbf{Assoc}$ be the projection map. Then we get $\phi^* A \in \text{Alg}_{C_*(\mathbb{E}_1)}(\text{Ch}(\mathbb{k}))$. If $\tilde{A} \xrightarrow{\sim} \phi^* A$ is a cofibrant replacement, we can then use Theorem 4.28 to get an object $\tilde{A} \in \text{Alg}_{\mathbb{E}_1}(N_{\text{dg}}(\text{Ch}(\mathbb{k})))$. The rest of this section will go into proving the following theorem.

Theorem 4.30. *For any projective resolution $P \xrightarrow{\sim} A$ of A as an A^e -module, the evaluation map*

$$\text{Map}_{\text{Ch}(A^e)}(P, P) \otimes_{\mathbb{k}} P \rightarrow P$$

makes the Hochschild complex of A into a center of $\tilde{A} \in \text{Alg}_{\mathbb{E}_1}(\text{Ch}(\mathbb{k}))$. In particular, this makes $\text{Map}_{\text{Ch}(A^e)}(P, P)$ into an element of $\text{Alg}_{\mathbb{E}_1}(\text{Alg}_{\mathbb{E}_1}(N_{\text{dg}}(\text{Ch}(\mathbb{k})))) \simeq \text{Alg}_{\mathbb{E}_2}(N_{\text{dg}}(\text{Ch}(\mathbb{k})))$.

Corollary 4.31. *Take $P = B(A) \xrightarrow{\sim} A$ to be the Bar complex. The strictification of the Hochschild complex $\text{Hom}_{\mathbb{k}}(A^{\otimes*}, A) \in \text{Ch}(\mathbb{k})$ as the center of \tilde{A} naturally carries the structure of a $C_*(\mathbb{E}_2)$ -algebra. This $C_*(\mathbb{E}_2)$ -algebra structure recovers the classical Gerstenhaber algebra structure in cohomology.*

The proof of the Corollary is given at the end of the next section.

4.4. The center as endomorphism object of bimodules. To show that the Hochschild complex indeed yields a center for \tilde{A} , we need some methods to compute the center in algebra categories. Throughout this chapter, let \mathcal{O}^\otimes be a coherent ∞ -operad; we will mostly be interested in the case $\mathcal{O}^\otimes = \mathbb{E}_1^\otimes$. Let $\mathcal{C}^\otimes \rightarrow \text{Fin}_*$ be a symmetric monoidal ∞ -category and consider the unique bifunctor of ∞ -operads $\mathcal{O}^\otimes \times \mathcal{LM}^\otimes \rightarrow \text{Fin}_* \times \text{Fin}_* \xrightarrow{\wedge} \text{Fin}_*$. We get a coCartesian fibration $\text{Alg}_{\mathcal{O}}(\mathcal{C})^\otimes \rightarrow \mathcal{LM}^\otimes$ with fibers over \mathfrak{a} and \mathfrak{m} respectively both equivalent to $\text{Alg}_{\mathcal{O}}(\mathcal{C})$. This comes from the fact that $\text{Alg}_{\mathcal{O}}(\mathcal{C})$ admits the structure of a symmetric monoidal ∞ -category and is hence left tensored over itself. Let $A \in \text{Alg}_{\mathcal{O}}(\mathcal{C})_{\mathfrak{m}}$. If A admits a center, it has the structure of an \mathbb{E}_1 algebra object in $\text{Alg}_{\mathcal{O}}(\mathcal{C})_{\mathfrak{a}}$. In particular, if $\mathcal{O}^\otimes = \mathbb{E}_k^\otimes$ is a little k -cubes operad, by Dunn Additivity we get

$$\mathfrak{Z}(A) \in \text{Alg}_{\mathbb{E}_1}(\text{Alg}_{\mathbb{E}_k}(\mathcal{C})) \simeq \text{Alg}_{\mathbb{E}_{k+1}}(\mathcal{C}).$$

To better understand the situation, consider the 1-categorical case. Let A be an associative algebra over \mathbb{k} . Then there is an equivalence of 1-categories between algebra objects in the monoidal category of A -bimodules and associative algebras under A . In particular, an A -bimodule structure on an associative algebra B is the same as an algebra morphism $A \rightarrow B$, and the equivalence is given by sending the bimodule structure to its unit morphism. The centralizer of an algebra morphism $f : A \rightarrow B$ is defined as

$$\mathfrak{Z}(f) = \{b \in B : \forall a \in A : f(a)b = bf(a)\}.$$

Viewing the data of f as an A -bimodule structure on B , we can see that this agrees with the set of A -bimodule maps from A to B :

$$\mathfrak{Z}(f) \cong \text{Hom}_{\text{Bimod}_A}(A, B).$$

To generalize this relationship between the center and bimodule morphisms to ∞ -operads, one needs to first recover the statement on algebra objects in the monoidal category of bimodules. To

this end, let $A \in \text{Alg}_{\mathcal{O}}(\mathcal{C})_{\mathfrak{m}}$ and let $\bar{A} \in \text{LMod}_{\mathbb{1}}(\text{Alg}_{\mathcal{O}}(\mathcal{C})_{\mathfrak{m}})$ be a lift of A as a module over the trivial algebra. We get a coCartesian \mathcal{LM}^{\otimes} -family of \mathcal{O} -operads

$$\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \mathcal{LM}^{\otimes}) \rightarrow \mathcal{O}^{\otimes} \times \mathcal{LM}^{\otimes},$$

and we can regard \bar{A} as a coCartesian \mathcal{LM}^{\otimes} -family of \mathcal{O} -algebras by noting that there is a bijection

$$\text{Fun}_{\mathcal{LM}^{\otimes}}(\mathcal{LM}^{\otimes}, \text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \mathcal{LM}^{\otimes}))) \simeq \text{Fun}_{\mathcal{LM}^{\otimes}}(\mathcal{LM}^{\otimes}, \text{Alg}_{\mathcal{O}}(\mathcal{C})^{\otimes}).$$

We can view A as an \mathcal{O} -module over itself, and hence \bar{A} also determines a coCartesian \mathcal{LM}^{\otimes} -family of \mathcal{O} -algebras in the coCartesian \mathcal{LM}^{\otimes} -family of \mathcal{O} -operads

$$\bar{\mathcal{C}}^{\otimes} := \text{Mod}_{\bar{A}}^{\mathcal{O}, \mathcal{LM}^{\otimes}}(\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \mathcal{LM}^{\otimes}))^{\otimes} \rightarrow \mathcal{O}^{\otimes} \times \mathcal{LM}^{\otimes}.$$

This notation allows us to identify algebra objects in the category of A -bimodules as algebras under A .

Proposition 4.32 (Proposition 5.3.1.27 [Lur17]). *The forgetful functor*

$$\theta : \text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\text{Mod}_{\bar{A}}^{\mathcal{O}, \mathcal{LM}^{\otimes}}(\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \mathcal{LM}^{\otimes})))^{\bar{A}_{\mathcal{LM}^{\otimes}/}} \rightarrow \text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \mathcal{LM}^{\otimes}))^{\bar{A}_{\mathcal{LM}^{\otimes}/}}$$

is an equivalence of ∞ -categories. Note that for all $s \in \mathcal{LM}^{\otimes}$, the algebra $\bar{A}_s \in \text{Alg}_{/\mathcal{O}}(\text{Mod}_{\bar{A}_s}^{\mathcal{O}}(\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \{s\})))$ is a trivial algebra, so in particular for $s = \mathfrak{m}$ we get an equivalence

$$\theta_{\mathfrak{m}} : \text{Alg}_{/\mathcal{O}}(\text{Mod}_{\bar{A}}^{\mathcal{O}}(\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \{\mathfrak{m}\}))) \rightarrow \text{Alg}_{/\mathcal{O}}(\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \{\mathfrak{m}\}))^{A/} \simeq \text{Alg}_{\mathcal{O}}(\mathcal{C})_{\mathfrak{m}}^{A/}.$$

Since $\bar{A}(\mathfrak{a})$ is the trivial algebra, we also have an equivalence $\bar{\mathcal{C}}_{\mathfrak{a}}^{\otimes} \simeq \mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \{\mathfrak{a}\})$, and therefore $\text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\bar{\mathcal{C}})_{\mathfrak{a}} \simeq \text{Alg}_{\mathcal{O}}(\mathcal{C})_{\mathfrak{a}}$. To find the centralizer of id_A in $\text{Alg}_{\mathcal{O}}(\mathcal{C})_{\mathfrak{m}}$, it hence suffices to find the centralizer of id_A in $\text{Alg}_{/\mathcal{O}}(\text{Mod}_{\bar{A}}^{\mathcal{O}}(\mathcal{C}^{\otimes} \times_{\mathcal{F}\text{in}_*} (\mathcal{O}^{\otimes} \times \{\mathfrak{m}\})))$, in which A is the trivial algebra. After that, we can use Proposition 2.13 to get the center $\mathfrak{Z}(A) \in \text{Alg}_{\mathbb{E}_1}(\text{Alg}_{\mathcal{O}}(\mathcal{C})_{\mathfrak{a}})$ of A .

Theorem 4.33 (Proposition 5.3.1.29 [Lur17]). *Suppose that for all $X \in \mathcal{O}$, there exists a morphism object $\text{Mor}_{\bar{\mathcal{C}}_{X, \mathfrak{m}}}(A(X), A(X)) \in \bar{\mathcal{C}}_{X, \mathfrak{a}}$. Then there exists a centralizer $\mathfrak{Z}(\text{id}_A) \in \text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\bar{\mathcal{C}})_{\mathfrak{a}}$. Furthermore, if $Z \in \text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\bar{\mathcal{C}})_{\mathfrak{a}}$, then a commutative diagram*

$$\begin{array}{ccc} & Z \otimes A & \\ A \swarrow & & \searrow A \\ & A & \end{array} \quad \text{with } \text{id}_A \text{ on the bottom arrow.}$$

exhibits Z as the centralizer of id_A if and only if for all $X \in \mathcal{O}$, the induced map $Z(X) \otimes A(X) \rightarrow A(X)$ exhibits $Z(X)$ as a morphism object of $A(X)$ and $A(X)$.

Proof. By definition, the centralizer is a final object of the ∞ -category

$$\mathcal{A} := (\text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\bar{\mathcal{C}})_{\mathfrak{a}})_{\mathbb{1}} \times_{(\text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\bar{\mathcal{C}})_{\mathfrak{m}})_{A/}} (\text{Alg}_{/\mathcal{O}}^{\mathcal{LM}^{\otimes}}(\bar{\mathcal{C}})_{\mathfrak{m}})_{A//A}.$$

Since \bar{A}_s is the trivial algebra in $\bar{\mathcal{C}}_s^{\otimes}$ for $s \in \{\mathfrak{a}, \mathfrak{m}\}$, we can use [Lur17, Theorem 2.2.2.4] to get an \mathcal{O}^{\otimes} -monoidal ∞ -category

$$\mathcal{E}^{\otimes} := (\bar{\mathcal{C}}_{\mathfrak{a}}^{\otimes})_{\mathbb{1}_{\mathcal{O}/}} \times_{(\bar{\mathcal{C}}_{\mathfrak{m}}^{\otimes})_{A_{\mathcal{O}/}}} (\bar{\mathcal{C}}_{\mathfrak{m}}^{\otimes})_{A_{\mathcal{O}///A_{\mathcal{O}}}} \rightarrow \mathcal{O}^{\otimes}$$

such that $\text{Alg}_{/\mathcal{O}}(\mathcal{E}) \simeq \mathcal{A}$. Finally, use that limits in algebra categories are computed object-wise by [Lur17, Corollary 3.2.2.5] to argue that we are reduced to showing that for each $X \in \mathcal{O}$, the fiber \mathcal{E}_X admits a final object. But a final object in

$$\mathcal{E}_X \simeq (\overline{\mathcal{C}}_{X,\mathfrak{a}})_{\mathbb{1}(X)/} \times_{(\overline{\mathcal{C}}_{X,\mathfrak{m}})_{A(X)/}} (\overline{\mathcal{C}}_{X,\mathfrak{m}})_{A(X)//A(X)}$$

is equivalent to a morphism object $\text{Mor}_{\overline{\mathcal{C}}_{X,\mathfrak{m}}}(A(X), A(X))$ by Proposition 2.4. \square

Corollary 4.34. *Let $\mathcal{O}^\otimes = \mathbb{E}_1^\otimes$, and let $A \in \text{Alg}_{\mathbb{E}_1}(\mathcal{C})$ be an \mathbb{E}_1 -algebra in a symmetric monoidal ∞ -category \mathcal{C}^\otimes . Assume that the morphism object $\text{Mor}_{\text{Mod}_A^{\mathbb{E}_1}(\mathcal{C}^\otimes \times_{\mathcal{F}\text{in}_*} \mathbb{E}_1^\otimes)}(A(\mathfrak{a}), A(\mathfrak{a})) \in \mathcal{C}$ exists. Then there exists a centralizer $\mathfrak{Z}(\text{id}_A) \in \text{Alg}_{\mathbb{E}_1}(\mathcal{C})$ with underlying object*

$$\mathfrak{Z}(\text{id}_A)(\mathfrak{a}) \simeq \text{Mor}_{\text{Mod}_A^{\mathbb{E}_1}(\mathcal{C}^\otimes \times_{\mathcal{F}\text{in}_*} \mathbb{E}_1^\otimes)}(A(\mathfrak{a}), A(\mathfrak{a})),$$

and the action of the centralizer has underlying map given by the evaluation α of the morphism object on $A(\mathfrak{a})$. Further, the multiplication of the \mathbb{E}_1 -algebra structure on $\mathfrak{Z}(\text{id}_A)$ is induced by the action of the tensor product $\mathfrak{Z}(\text{id}_A)(\mathfrak{a}) \otimes \mathfrak{Z}(\text{id}_A)(\mathfrak{a})$ on $A(\mathfrak{a})$ given by the tensor product $\alpha \otimes \alpha$ in $\mathcal{E}_\mathfrak{a}$:

$$* : A(\mathfrak{a}) \xrightarrow{\simeq} A(\mathfrak{a}) \otimes A(\mathfrak{a}) \rightarrow (\mathfrak{Z}(\text{id}_A)(\mathfrak{a}) \otimes A(\mathfrak{a})) \otimes (\mathfrak{Z}(\text{id}_A)(\mathfrak{a}) \otimes A(\mathfrak{a})) \rightarrow A(\mathfrak{a}) \otimes A(\mathfrak{a}) \xrightarrow{\text{mult.}} A(\mathfrak{a}).$$

Proof. The first part follows directly from Theorem 4.33. For the claim about the multiplication, note that by the proof of Theorem 4.33 the algebra structure on the centralizer is induced from $\mathfrak{Z}(\text{id}_A)(\mathfrak{a})$ being final in $\mathcal{E}_\mathfrak{a}$. In particular, in $\mathcal{E}_\mathfrak{a}$ we have a unique-up-to-contractible-choice map

$$\mathfrak{Z}(\text{id}_A)(\mathfrak{a}) \otimes \mathfrak{Z}(\text{id}_A)(\mathfrak{a}) \rightarrow \mathfrak{Z}(\text{id}_A).$$

Now recall the construction of the tensor product in operadic slice categories in [Lur17, Theorem 2.2.2.4]. This shows that up to homotopy, the monoidal product in $\mathcal{E}_\mathfrak{a}$ is given in the first component by

$$\begin{array}{ccc} \begin{array}{c} \mathbb{1} \\ \downarrow \\ X \end{array} & \otimes & \begin{array}{c} \mathbb{1} \\ \downarrow \\ Y \end{array} \\ & & = \\ & & \begin{array}{c} \mathbb{1} \\ \downarrow \scriptstyle \simeq \\ \mathbb{1} \otimes \mathbb{1} \\ \downarrow \\ X \otimes Y \end{array} \end{array}$$

and in the second component by

$$\begin{array}{ccc}
 \begin{array}{c} A(\mathfrak{a}) \\ \simeq \downarrow \\ \mathbb{1} \otimes A(\mathfrak{a}) \\ \downarrow \\ X \otimes A(\mathfrak{a}) \end{array} & \xrightarrow{\text{id}_{A(\mathfrak{a})}} & A(\mathfrak{a}) \\
 & \searrow & \\
 & & A(\mathfrak{a})
 \end{array}
 \quad \otimes \quad
 \begin{array}{c} A(\mathfrak{a}) \\ \simeq \downarrow \\ \mathbb{1} \otimes A(\mathfrak{a}) \\ \downarrow \\ Y \otimes A(\mathfrak{a}) \end{array}
 \xrightarrow{\text{id}_{A(\mathfrak{a})}}
 A(\mathfrak{a})$$

$$\begin{array}{c}
 A(\mathfrak{a}) \\
 \simeq \downarrow \\
 A(\mathfrak{a}) \otimes A(\mathfrak{a}) \\
 \simeq \downarrow \\
 (\mathbb{1} \otimes A(\mathfrak{a})) \otimes (\mathbb{1} \otimes A(\mathfrak{a})) \\
 \downarrow \\
 (X \otimes A(\mathfrak{a})) \otimes (Y \otimes A(\mathfrak{a})) \longrightarrow A(\mathfrak{a}) \otimes A(\mathfrak{a}) \xrightarrow{\text{mult}} A(\mathfrak{a})
 \end{array}
 \xrightarrow{\text{id}_{A(\mathfrak{a})}}
 A(\mathfrak{a})$$

This corresponds to the tensor product action given in the statement. \square

Corollary 4.35. *Let again $\mathcal{O}^\otimes = \mathbb{E}_1^\otimes$ and $A \in \text{Alg}_{\mathbb{E}_1}(\mathcal{C})$ as above, and assume that the morphism object exists. Then there exists a center $\mathfrak{Z}(A) \in \text{Alg}_{\mathbb{E}_1}(\text{Alg}_{\mathbb{E}_1}(\mathcal{C})) \simeq \text{Alg}_{\mathbb{E}_2}(\mathcal{C})$ with underlying object $\mathfrak{Z}(A)(\mathfrak{a}) \simeq \text{Mor}_{\text{Mod}_A^{\mathbb{E}_1}(\mathcal{C}^\otimes \times_{\mathcal{F}\text{in}_*} \mathbb{E}_1^\otimes)_\mathfrak{a}}(A(\mathfrak{a}), A(\mathfrak{a}))$. The outer multiplication is given by the composition product*

$$\circ : (\mathfrak{Z}(A)(\mathfrak{a}) \otimes \mathfrak{Z}(A)(\mathfrak{a})) \otimes A(\mathfrak{a}) \xrightarrow{\text{id} \otimes \alpha} \mathfrak{Z}(A)(\mathfrak{a}) \otimes A(\mathfrak{a}) \xrightarrow{\alpha} A(\mathfrak{a}).$$

The inner multiplication is given by the convolution product described in Corollary 4.34. View the center as an object in $\text{Alg}_{\mathbb{E}_1}(\text{Alg}_{/\mathbb{E}_1}(\mathcal{E}))$ with the monoidal structure on $\text{Alg}_{/\mathbb{E}_1}(\mathcal{E})$ given by [Lur17, Definition 4.7.1.1]. The two multiplications assemble into a square

$$\begin{array}{ccc}
 (\mathfrak{Z}(A)(\mathfrak{a}) \otimes \mathfrak{Z}(A)(\mathfrak{a}))(\mathfrak{a}) \otimes (\mathfrak{Z}(A)(\mathfrak{a}) \otimes \mathfrak{Z}(A)(\mathfrak{a}))(\mathfrak{a}) & \longrightarrow & \mathfrak{Z}(A)(\mathfrak{a})(\mathfrak{a}) \otimes \mathfrak{Z}(A)(\mathfrak{a})(\mathfrak{a}) \\
 \downarrow & & \downarrow_* \\
 (\mathfrak{Z}(A)(\mathfrak{a}) \otimes \mathfrak{Z}(A)(\mathfrak{a}))(\mathfrak{a}) & \xrightarrow{\circ} & \mathfrak{Z}(A)(\mathfrak{a})(\mathfrak{a})
 \end{array}$$

in $\mathcal{E}_\mathfrak{a}$. Since $\mathfrak{Z}(A)(\mathfrak{a})(\mathfrak{a})$ is the morphism object and thus final, there is a contractible choice of 2-simplices filling this square.

Proof. By Proposition 2.13, the action \circ above is the one induced by the monoidal structure on $\text{Alg}_{/\mathbb{E}_1}(\mathcal{E})$. The square is given by

$$\begin{array}{ccc}
 \mathfrak{Z}(A)(\{\mathfrak{a}, \mathfrak{a}\})(\{\mathfrak{a}, \mathfrak{a}\}) & \longrightarrow & \mathfrak{Z}(A)(\mathfrak{a})(\{\mathfrak{a}, \mathfrak{a}\}) \\
 \downarrow & & \downarrow \\
 \mathfrak{Z}(A)(\{\mathfrak{a}, \mathfrak{a}\})(\mathfrak{a}) & \longrightarrow & \mathfrak{Z}(A)(\mathfrak{a})(\mathfrak{a})
 \end{array}$$

\square

Remark 4.36. Note that the square in the proof of Corollary 4.35 is precisely the one inducing the bracket in 3.21. Hence, the above shows that indeed the two multiplications on the center of an associative algebra are up to homotopy given by the convolution and composition product, and the bracket is up to homotopy given by the "circle product" from the classical Gerstenhaber algebra structure.

Proof of Corollary 4.31. By Theorem 4.30, the classical Hochschild complex is a center of \tilde{A} and thus inherits a \mathbb{E}_2 -algebra structure in the derived ∞ -category. The first part of the Corollary follows directly from the Rectification Theorem 4.28. For the second part, we use Corollary 3.22. By Corollary 4.35, the two multiplications on $\mathfrak{Z}(\tilde{A})$ are given by composition and convolution respectively. But both of these recover the formula for the cup product for the Bar resolution Hochschild cochain complex. Hence, we get the classical cup product in cohomology. Now the circle product yields a homotopy of the square

$$\begin{array}{ccc} \mathrm{Hom}_{\mathbb{k}}(A^{\otimes*}, A)^{\otimes 4} & \xrightarrow{\smile \otimes \smile} & \mathrm{Hom}_{\mathbb{k}}(A^{\otimes*}, A)^{\otimes 2} \\ (\smile \otimes \smile) \circ (\mathrm{id} \otimes \tau \otimes \mathrm{id}) \downarrow & \swarrow h & \downarrow \smile \\ \mathrm{Hom}_{\mathbb{k}}(A^{\otimes*}, A)^{\otimes 2} & \xrightarrow{\smile} & \mathrm{Hom}_{\mathbb{k}}(A^{\otimes*}, A) \end{array}$$

by setting $h : \mathrm{Hom}_{\mathbb{k}}(A^{\otimes*}, A)^{\otimes 4} \rightarrow \mathrm{Hom}_{\mathbb{k}}(A^{\otimes*-1}, A)$,

$$f_1 \otimes g_1 \otimes f_2 \otimes g_2 \mapsto (-1)^{|f_1|+|g_1|} f_1 \smile (f_2 \{g_1\}) \smile g_2,$$

where $f\{g\}$ denotes the circle product. Restricting along $\iota_{1,4}$ just yields the trivial homotopy, and restricting along $\iota_{2,3}$ yields the homotopy

$$\begin{aligned} H : \mathrm{Hom}_{\mathbb{k}}(A^{\otimes p}, A) \otimes \mathrm{Hom}_{\mathbb{k}}(A^{\otimes q}, A) &\rightarrow \mathrm{Hom}_{\mathbb{k}}(A^{\otimes p+q-1}, A) \\ f \otimes g &\mapsto (-1)^p g \{f\} \end{aligned}$$

between \smile and $\smile \circ \tau$. For the opposite multiplications, we can just whisker with τ . Hence by the Remark after Corollary 4.35, this yields the bracket as

$$\begin{aligned} [f, g] &= H(f \otimes g) + H\tau(f \otimes g) = (-1)^{|f|} g \{f\} + (-1)^{|g|+|f||g|} f \{g\} \\ &= (-1)^{|g|+|f||g|} (f \{g\} - (-1)^{(|f|+1)(|g|+1)} g \{f\}) \\ &= (-1)^{|g|+|f||g|} [f, g]_G. \end{aligned}$$

But this agrees with the classical Gerstenhaber bracket up to a sign, proving the claim. \square

4.5. The category of \mathbb{E}_1 -modules in chain complexes. As explained above, we want to apply Corollary 4.35 to the symmetric monoidal ∞ -category $\mathcal{C}^{\otimes} = N_{\mathrm{dg}}(\mathrm{Ch}(\mathbb{k}))^{\otimes}$ and $\tilde{A} \in \mathrm{Alg}_{\mathbb{E}_1}(\mathcal{C})$ obtained from a \mathbb{k} -algebra A . Note that \mathbb{E}_2^{\otimes} only has a single color \mathfrak{a} , and by the above theorem the underlying object of the center $\mathfrak{Z}(\tilde{A})$ at this color is equivalent to the morphism object $\mathrm{Mor}_{\bar{\mathcal{C}}_{\mathfrak{a}, \mathfrak{m}}}(\tilde{A}, \tilde{A}) \in \bar{\mathcal{C}}_{\mathfrak{a}, \mathfrak{a}}$. Here \tilde{A} is viewed as a module over itself, i.e. as an object of

$$\bar{\mathcal{C}}_{\mathfrak{a}, \mathfrak{m}} = \mathrm{Mod}_{\tilde{A}}^{\mathbb{E}_1}(\mathcal{C} \times_{N(\mathrm{Fin}_*)} \mathbb{E}_1)_{\mathfrak{a}}.$$

Note that $\bar{\mathcal{C}}_{\mathfrak{a}, \mathfrak{a}} = \mathrm{Mod}_1^{\mathbb{E}_1}(\mathcal{C} \times_{N(\mathrm{Fin}_*)} \mathbb{E}_1)_{\mathfrak{a}} \simeq (\mathcal{C}^{\otimes} \times_{N(\mathrm{Fin}_*)} \mathbb{E}_1^{\otimes})_{\mathfrak{a}} \simeq \mathcal{C}$, so

$$\mathfrak{Z}(\tilde{A})(\mathfrak{a}) \simeq \mathrm{Mor}_{\mathrm{Mod}_{\tilde{A}}^{\mathbb{E}_1}(\mathcal{C} \times_{N(\mathrm{Fin}_*)} \mathbb{E}_1)_{\mathfrak{a}}}(\tilde{A}, \tilde{A}) \in \mathcal{C}.$$

We are hence reduced to showing that $\text{Map Ch}(A^e)(P, P)$ together with the evaluation map is such a morphism object, meaning that it satisfies the universal property of 2.3.

To show this, we must understand the ∞ -category $\text{Mod}_{\tilde{A}}^{\mathbb{E}_1}(N_{\text{dg}}(\text{Ch}(k)) \times_{N(\mathcal{F}\text{in}_*)} \mathbb{E}_1)_{\mathfrak{a}}$. By [Hin15, Proposition B.1.2], we have an equivalence of ∞ -categories

$$\text{Mod}_{\tilde{A}}^{\mathbb{E}_1}(N_{\text{dg}}(\text{Ch}(\mathbb{k})) \times_{N(\mathcal{F}\text{in}_*)} \mathbb{E}_1)_{\mathfrak{a}} \simeq \text{Alg}_{M\mathbb{E}_1}(N_{\text{dg}}(\text{Ch}(\mathbb{k}))) \times_{\text{Alg}_{\mathbb{E}_1}(N_{\text{dg}}(\text{Ch}(\mathbb{k})))} \{\tilde{A}\}.$$

Here $M\mathbb{E}_1$ is defined as in [Hin15, 5.2]. We can now use Hinich's Rectification Theorem for modules [Hin15, Theorem 5.2.3] to get an equivalence of ∞ -categories

$$N(\text{Mod}_{\tilde{A}}^{C_*(\mathbb{E}_1)}(\text{Ch}(\mathbb{k}))^c)[W_{\text{Mod}}^{-1}] \xrightarrow{\simeq} \text{Mod}_{\tilde{A}}^{\mathbb{E}_1}(N_{\text{dg}}(\text{Ch}(\mathbb{k})) \times_{N(\mathcal{F}\text{in}_*)} \mathbb{E}_1)_{\mathfrak{a}}.$$

Note here that since \tilde{A} is cofibrant, by [BM09, Theorem 2.6] the module category $\text{Mod}_{\tilde{A}}^{C_*(\mathbb{E}_1)}(\text{Ch}(\mathbb{k}))$ indeed carries a model category structure transferred via the forgetful functor to $\text{Ch}(\mathbb{k})$. Similarly, by [BM09, Proposition 2.7], the category $\text{LMod}_{U_{C_*(\mathbb{E}_1)}(\tilde{A})}(\text{Ch}(\mathbb{k}))$ can be made into a model category via transfer from the forgetful functor, where $U_{C_*(\mathbb{E}_1)}(\tilde{A})$ is the universal enveloping algebra of \tilde{A} viewed as an $C_*(\mathbb{E}_1)$ -algebra. By [BM09, Theorem 1.10] we have an isomorphism of categories making the following diagram commute

$$\begin{array}{ccc} \text{Mod}_{\tilde{A}}^{C_*(\mathbb{E}_1)}(\text{Ch}(\mathbb{k})) & \xrightarrow{\cong} & \text{LMod}_{U_{C_*(\mathbb{E}_1)}(\tilde{A})}(\text{Ch}(\mathbb{k})) \\ & \searrow & \swarrow \\ & \text{Ch}(\mathbb{k}) & \end{array}.$$

In particular, this isomorphism yields a Quillen equivalence between these two model categories.

4.6. The trouble with universal enveloping algebras. We now want to relate this category of left modules over $U_{C_*(\mathbb{E}_1)}(\tilde{A})$ to the category $\text{Ch}(A^e)$. To this end, we have the following result.

Proposition 4.37. *There exists a zig-zag of quasi-isomorphisms between $U_{C_*(\mathbb{E}_1)}(\tilde{A})$ and $A \otimes A^{\text{op}}$.*

Proof. Let $\theta : \mathbb{A}_{\infty} \xrightarrow{\simeq} \mathbf{Assoc}$ be a cofibrant replacement of dg operads, and note that we have a diagram

$$\begin{array}{ccc} & & C_*(\mathbb{E}_1) \\ & \nearrow \psi, \simeq & \downarrow \phi, \simeq \\ \mathbb{A}_{\infty} & \xrightarrow{\theta, \simeq} & \mathbf{Assoc} \end{array}$$

Since we work in characteristic zero, all of the involved operads are admissible and Σ -cofibrant. In particular, all the above weak equivalences are strong equivalences of operads, and thus induce a Quillen equivalence between their respective algebra categories. Let $A' \xrightarrow{\simeq} A$ be a cofibrant replacement in associative algebras, and let $\hat{A} \xrightarrow{\simeq} \theta^* A'$ be a cofibrant replacement in \mathbb{A}_{∞} -algebras. Then in particular, the unit map $\hat{A} \rightarrow \psi^* \psi_! \hat{A}$ is a weak equivalence, and hence using [Fre09, Theorem 17.4.A, 17.4.B] we get the following diagram of weak equivalences of dg algebras

$$\begin{array}{ccccc}
 & & U_{\mathbb{A}_\infty}(\hat{A}) & & \\
 & \swarrow \simeq & & \searrow \simeq & \\
 & U_{\mathbb{A}_\infty}(\theta^* A') & & U_{\mathbb{A}_\infty}(\psi^* \psi_! \hat{A}) & \\
 & \swarrow \theta_\flat & & \searrow \psi_\flat & \\
 A' \otimes A'^{\text{op}} = U_{\text{Assoc}}(A') & & & & U_{C_*(\mathbb{E}_1)}(\psi_! \hat{A}) \\
 \swarrow \simeq & & & & \\
 A \otimes_k A^{\text{op}} & & & &
 \end{array}$$

It hence suffices to show that $U_{C_*(\mathbb{E}_1)}(\psi_! \hat{A})$ is quasi-isomorphic to $U_{C_*(\mathbb{E}_1)}(\tilde{A})$. To this end, let $A_1 \xrightarrow{\simeq} \phi^* A'$ be any cofibrant replacement. Note that $\psi_! \hat{A}$ is again cofibrant, and we hence have a lift $f : \psi_! \tilde{A} \rightarrow A_1$ in the diagram

$$\begin{array}{ccc}
 \emptyset & \longrightarrow & A_1 \\
 \downarrow & \nearrow f & \downarrow \simeq \\
 \psi_! \tilde{A} & \xrightarrow{\simeq} & \phi^* A'
 \end{array}$$

and by 2-out-of-3, f must be a weak equivalence. This is a weak equivalence between cofibrant objects, so again by [Fre09, Theorem 17.4.A], we get a quasi-isomorphism $U_{C_*(\mathbb{E}_1)}(\psi_! \hat{A}) \xrightarrow{\simeq} U_{C_*(\mathbb{E}_1)}(A_1)$. The map $\phi^* A' \rightarrow \phi^* A$ is again a trivial fibration as ϕ^* is right Quillen, and in particular the composition

$$A_1 \xrightarrow{\simeq} \phi^* A' \xrightarrow{\simeq} \phi^* A$$

is again a cofibrant replacement. We now again find a lift $g : A_1 \rightarrow \tilde{A}$ in the diagram

$$\begin{array}{ccc}
 \emptyset & \longrightarrow & \tilde{A} \\
 \downarrow & \nearrow g & \downarrow \simeq \\
 A_1 & \xrightarrow{\simeq} & \phi^* A
 \end{array}$$

which again is a weak equivalence, and thus finally induces a quasi-isomorphism

$$U_{C_*(\mathbb{E}_1)}(A_1) \xrightarrow{\simeq} U_{C_*(\mathbb{E}_1)}(\tilde{A}).$$

Summarizing, we get the following zig-zag

$$A \otimes A^{\text{op}} \xleftarrow{\simeq} U_{\mathbb{A}_\infty}(\hat{A}) \xrightarrow{\simeq} U_{C_*(\mathbb{E}_1)}(\tilde{A}).$$

□

Corollary 4.38. *Let $\tilde{A} \xrightarrow{\simeq} \phi^* A$ be a cofibrant replacement of the associative algebra A as an $C_*(\mathbb{E}_1)$ -algebra. Then the category $\text{LMod}_{A \otimes A^{\text{op}}}(\text{Ch}(\mathbb{k}))$ is Quillen equivalent to the category $\text{LMod}_{U_{C_*(\mathbb{E}_1)}(\tilde{A})}(\text{Ch}(\mathbb{k}))$.*

Finally, we have an isomorphism

$$\begin{array}{ccc} \mathrm{LMod}_{A \otimes A^{\mathrm{op}}}(\mathrm{Ch}(\mathbb{k})) & \xrightarrow{\cong} & \mathrm{Ch}(A \otimes A^{\mathrm{op}}) \\ & \searrow & \swarrow \\ & \mathrm{Ch}(\mathbb{k}) & \end{array}$$

which again induces a Quillen equivalence. In particular, the two model categories $\mathrm{Ch}(A \otimes A^{\mathrm{op}})$ and $\mathrm{Mod}_{\tilde{A}}^{C_*(\mathbb{E}_1)}(\mathrm{Ch}(\mathbb{k}))$ are Quillen equivalence and therefore present the same ∞ -category.

Corollary 4.39. *There is an equivalence of ∞ -categories*

$$N_{\mathrm{dg}}(\mathrm{Ch}(A \otimes A^{\mathrm{op}})^{\circ}) \xrightarrow{\simeq} \mathrm{Mod}_{\tilde{A}}^{\mathbb{E}_1}(N_{\mathrm{dg}}(\mathrm{Ch}(\mathbb{k})) \times_{N(\mathcal{F}\mathrm{in}_*)} \mathbb{E}_1)_{\mathbf{a}}.$$

Proof. This follows from the above discussion together with the fact that $\mathrm{Ch}(A \otimes A^{\mathrm{op}})$ satisfies the conditions of Corollary 4.24. \square

4.7. Morphism objects in dg model categories. To proof Theorem 4.30, we want to argue that $\mathrm{MapCh} A^e(P, P)$ is an endomorphism object of \tilde{A} in the \mathbb{E}_1 -module category over \tilde{A} . We have already seen that this module category is equivalent to the ∞ -category $N_{\mathrm{dg}}(\mathrm{Ch}(A^e)^{\circ})$, and we know that $\mathrm{MapCh} A^e(P, P) \in \mathrm{Ch}(\mathbb{k})$ is indeed an endomorphism object for $P \simeq A$ in the dg category $\mathrm{Ch}(A^e)$. To finish the argument, we will now proof the following general theorem connecting morphism objects in a dg category and its underlying ∞ -category.

Lemma 4.40. *Let C be a monoidal dg model category with underlying monoidal product $\otimes : C_0 \times C_0 \rightarrow C_0$. Then the induced monoidal product $N_{\mathrm{dg}}(C^{\circ}) \times N_{\mathrm{dg}}(C^{\circ}) \rightarrow N_{\mathrm{dg}}(C^{\circ})$ sends $A, B \in C^{\circ}$ to an object equivalent to $R(A \otimes B)$. A similar statement holds for dg model categories left tensored over a monoidal dg model category.*

Proof. This follows directly from the description of the monoidal structure in Proposition 4.26. \square

Theorem 4.41. *Let C be a monoidal dg model category and let M be a dg model category that is left tensored over C . In particular, we have a dg functor $\otimes : C \boxtimes M \rightarrow M$ whose underlying functor is a left Quillen bifunctor. Assume that for $A, B \in M^{\circ}$ we have a dg morphism object $\mathrm{Mor}_M(A, B) \in C$ together with map $\alpha : \mathrm{Mor}_M(A, B) \otimes A \rightarrow B$ in M such that composition with α induces an isomorphism*

$$(2) \quad \mathrm{Map}_C(C, \mathrm{Mor}_M(A, B)) \cong \mathrm{Map}_M(C \otimes A, B).$$

Then

- (1) *The induced map $\tilde{\alpha} \in \mathrm{Map}_{N_{\mathrm{dg}}(M^{\circ})}(R(Q \mathrm{Mor}_M(A, B) \otimes A), B)$ makes $Q \mathrm{Mor}_M(A, B) \in N_{\mathrm{dg}}(C^{\circ})$ into a morphism object for $A, B \in N_{\mathrm{dg}}(M^{\circ})$ in the sense of Definition 2.3.*
- (2) *If $\beta : R(M \otimes A) \rightarrow B$ is another morphism object for A and B in $N_{\mathrm{dg}}(M^{\circ})$, then $f : M \xrightarrow{\simeq} Q \mathrm{Mor}_M(A, B)$ is a weak equivalence in C , and $\tilde{\alpha} \circ R(f \otimes \mathrm{id}_A) \simeq \beta$ are chain homotopic.*

Proof. For (1), note that if $C \in N_{\mathrm{dg}}(C^{\circ})$ is bifibrant, $Q \mathrm{Mor}_M(A, B) \xrightarrow{\simeq} \mathrm{Mor}_M(A, B)$ is the cofibrant replacement map, and $C \otimes A \xrightarrow{\simeq} R(C \otimes A)$ is the fibrant replacement map, we get a weak equivalence

$$\mathrm{Map}_C(C, Q \mathrm{Mor}_M(A, B)) \xrightarrow{\simeq} \mathrm{Map}_C(C, \mathrm{Mor}_M(A, B)) \cong \mathrm{Map}_M(C \otimes A, B) \xrightarrow{\simeq} \mathrm{Map}_M(R(C \otimes A), B)$$

of chain complexes. Applying $\mathrm{DK}_{\bullet, \tau \geq 0}$, we get

$$\mathrm{Map}_{N_{\mathrm{dg}}(C^{\circ})}(C, Q \mathrm{Mor}_M(A, B)) \simeq \mathrm{Map}_{N_{\mathrm{dg}}(M^{\circ})}(R(C \otimes A), B).$$

Together with Lemma 4.40 this yields the result.

For (2), we automatically get $M \simeq \text{Mor}_M(A, B)$ in the ∞ -category $N_{\text{dg}}(C^\circ)$ since morphism objects are unique up to equivalence. Now recall that $N_{\text{dg}}(C^\circ) \simeq N(C^e)[W^{-1}]$, and since model categories are saturated this implies the result. \square

This finally allows us to proof the main theorem of this section.

Proof of Theorem 4.30. By Theorem 4.33, it suffices to show that $\text{ev} : \text{Map}_{\text{Ch}(A^e)}(P, P) \otimes_k P \rightarrow P$ makes $\text{Map}_{\text{Ch}(A^e)}(P, P)$ into an endomorphism object for \tilde{A} in the ∞ -category $\text{Mod}_{\tilde{A}}^{\mathbb{E}_1}(\mathcal{C} \times_{N(\mathcal{F}\text{in}_*)} \mathbb{E}_1)_a$. By Corollary 4.39, this ∞ -category is equivalent to the ∞ -category $N_{\text{dg}}(\text{Ch}(A^e)^\circ)$. Note that $\mathcal{M} = \text{Ch}(A^e)$ satisfies the conditions of Theorem 4.41 with $C = \text{Ch}(\mathbb{k})$, and $P \in \mathcal{M}$ is indeed cofibrant (and thus bifibrant). Therefore, the evaluation map above indeed makes $\text{Map}_{\text{Ch}(A^e)}(P, P)$ into a morphism object for $P \in N_{\text{dg}}(\text{Ch}(A^e)^\circ)$. Clearly, P is equivalent to \tilde{A} viewed in the \mathbb{E}_1 -module category, and we hence get our result. \square

5. THE HOCHSCHILD COMPLEX OF A SCHEME

We would now like to globalize the above results and consider a quasi-compact separable scheme X over \mathbb{k} . The structure sheaf \mathcal{O}_X can be viewed as an $C_*(\mathbb{E}_1)$ -algebra in the category of sheaves of \mathbb{k} -vector spaces on X . We argue that the \mathbb{E}_1 -center of the structure sheaf is a good model of the Hochschild cochain complex of X . Note that this does not require X to be smooth.

5.1. The ∞ -category of sheaves of \mathbb{k} -modules. Let X be a quasi-compact separable scheme over \mathbb{k} . Then \mathcal{O}_X is an associative algebra object in the category of sheaves of \mathbb{k} -modules on X . We again want to view \mathcal{O}_X as a \mathbb{E}_1 -algebra in the associated ∞ -category. To this end, we consider a model structure on the category of dg presheaves presenting this ∞ -category.

Proposition 5.42 ([Hin05], Theorem 1.3.1). *Let S be a site. There is a cofibrantly generated model structure on the category $\text{dgPSh}(S)$ of presheaves of \mathbb{k} -module complexes on S with*

- *weak equivalences the maps $f : \mathcal{F} \rightarrow \mathcal{G}$ such that the degreewise sheafification $f^a : \mathcal{F}^a \rightarrow \mathcal{G}^a$ is a quasi-isomorphism of complexes of sheaves,*
- *cofibrations generated by maps $f : \mathcal{F} \rightarrow \mathcal{F}\langle x; dx = z \in \mathcal{F}(U) \rangle$ corresponding to adding a section to kill a cycle z over $U \in S$, and*
- *fibrations the maps $f : \mathcal{F} \rightarrow \mathcal{G}$ such that $f(U) : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is surjective for all $U \in S$ and for any hypercover $\epsilon : V_\bullet \rightarrow U$ the diagram*

$$\begin{array}{ccc} \mathcal{F}(U) & \longrightarrow & \check{C}(V_\bullet, \mathcal{F}) \\ f(U) \downarrow & & \downarrow \\ \mathcal{G}(U) & \longrightarrow & \check{C}(V_\bullet, \mathcal{G}) \end{array}$$

is a homotopy pullback.

This is the left Bousfield localization of the projective model structure on $\text{dgPSh}(S)$ with respect to the Čech complexes of hypercoverings. We therefore call it the local projective model structure. In particular, acyclic fibrations in the local projective model structure are just acyclic fibrations in the global projective model structure. Note that if S has enough points, weak equivalences can be detected at stalks.

Definition 5.43. Let X be a scheme over \mathbb{k} . Let $\text{Aff}(X)$ be the site of affine open subsets of X , and let $\text{Open}(X)$ be the site of all open subsets of X . Call the associated categories of dg presheaves $\text{dgPSh}^{\text{aff}}(X)$ and $\text{dgPSh}(X)$ respectively. We have a natural inclusion $\iota : \text{Aff}(X) \rightarrow \text{Open}(X)$ that induces to a restriction functor

$$\iota_* : \text{dgPSh}(X) \rightarrow \text{dgPSh}^{\text{aff}}(X)$$

Proposition 5.44. *The restriction functor ι_* admits a left adjoint ι^{-1} and the pair $\iota^{-1} \dashv \iota_*$ forms a Quillen equivalence. Both ι_* and ι^{-1} preserve weak equivalences, and ι^{-1} preserves acyclic fibrations. The unit $\text{id} \Rightarrow \iota_* \iota^{-1}$ is an isomorphism, and the counit $\iota^{-1} \iota_* \Rightarrow \text{id}$ is a component-wise weak equivalence.*

Proof. The left adjoint ι^{-1} is given by $\iota^{-1}\mathcal{F}(V) = \text{colim}_{V \subseteq U \in \text{Aff}(X)} \mathcal{F}(U)$. The direct image ι_* clearly preserves acyclic fibrations, since these are pointwise. The sites of all opens and of affine opens have the same points, namely points in the topological space X . This follows because affine opens form a basis of the Zariski topology. Even more, ι_* and ι^{-1} preserve stalks at these points. Taking stalks is a left adjoint, so this follows trivially for the inverse image, and for the direct image we note that small enough neighborhoods of a point $x \in X$ always contain an affine open neighborhood of x . This shows that both adjoints preserve weak equivalences, and in particular ι^{-1} preserves acyclic cofibrations. The fact that ι^{-1} preserves acyclic fibrations follows from the fact that filtered colimits are exact in Grothendieck categories. Finally, note that if U is affine, then $\text{colim}_{U \subseteq W \in \text{Aff}(X)} \mathcal{F}(W) \cong \mathcal{F}(U)$ since U is final in the index category. This shows that the unit is an isomorphism. The fact that the counit is a component-wise weak equivalence again follows from the fact that both adjoints preserve weak equivalences. \square

Definition 5.45. We call the underlying ∞ -category of the local projective model structure on $\text{dgPSh}(X)$ is the ∞ -category of sheaves of \mathbb{k} -modules on X

$$\text{Sh}_{\infty}(X) := N(\text{dgPSh}(X)^c)[W^{-1}].$$

By Proposition 5.44, we have an equivalence of ∞ -categories

$$\iota_* : N(\text{dgPSh}^{\text{aff}}(X)^c)[W^{-1}] \rightarrow \text{Sh}_{\infty}(X)$$

with quasi-inverse $\iota^{-1} : \text{Sh}_{\infty}(X) \rightarrow N(\text{dgPSh}^{\text{aff}}(X)^c)[W^{-1}]$.

Remark 5.46. Even though the model categories of presheaves on affine opens and general opens yield the same ∞ -category, the above model category structure depends on the choice of site. On affine open subsets, all quasi-coherent sheaves on X are automatically fibrant, which is not true for general opens. In particular, on affine opens the structure sheaf \mathcal{O}_X itself is fibrant.

Proposition 5.47. *If the topos on S has enough points and S admits finite products, then the local projective model structure yields a closed symmetric monoidal model category. If in addition S admits a final object, then $\text{dgPSh}(S)$ is a dg symmetric monoidal model category.*

Proof. By [PS18a, Proposition 7.9], the global projective model structure on $\text{dgPSh}(S)$ inherits the structure of a (symmetric) monoidal model category since S admits finite products. By [Whi22, Theorem 4.6], to show that this monoidal model structure descends to the local projective model structure, it suffices to argue that for f a local weak equivalence and \mathcal{F} a cofibrant object, the map $f \otimes \text{id}_{\mathcal{F}}$ is again a local weak equivalence. But this is clear if the topos has enough points, since we can then check local weak equivalences on stalks. Since the presheaf category admits an internal hom, this shows the first part. For the second part, note that presheaves of chain complexes are

the same as chain complexes of presheaves of \mathbb{k} -modules. Since the later is an abelian category, this automatically admits a dg enrichment. Recall that if $*$ $\in S$ is terminal, we have the constant presheaf functor

$$\begin{aligned} \text{const}_* : \text{Ch}(\mathbb{k}) &\rightarrow \text{dgPSh}(S) \\ C &\mapsto (U \mapsto C) \end{aligned}$$

By the argument below, this functor preserves cofibrations. We can hence define a tensoring

$$\text{Ch}(\mathbb{k}) \times \text{dgPSh}(S) \rightarrow \text{dgPSh}(S), \quad (C, \mathcal{F}) \mapsto \text{const}_*(C) \otimes \mathcal{F}$$

as well as a powering

$$\text{Ch}(\mathbb{k})^{\text{op}} \times \text{dgPSh}(S) \rightarrow \text{dgPSh}(S), \quad (C, \mathcal{F}) \mapsto \mathcal{H}om(\text{const}_*(C), \mathcal{F}).$$

One easily checks that these indeed satisfy the correct adjointness properties. It hence suffices to check the pushout-product axiom. But if $i : C \rightarrow D$ is a cofibration in $\text{Ch}(\mathbb{k})$, then $\text{const}_*(i) : \text{const}_*(C) \rightarrow \text{const}_*(D)$ is a cofibration in $\text{dgPSh}(S)$, and therefore this follows directly from the pushout-product axiom in $\text{dgPSh}(S)$. \square

Corollary 5.48. *The category $\text{dgPSh}(X)$ admits the structure of a dg symmetric monoidal model category.*

Let $U \subseteq X$ be an affine open. Then we have adjoint functors

$$\text{Ch}(\mathbb{k}) \xrightleftharpoons[\Gamma_U]{C_U} \text{dgPSh}^{\text{aff}}(X)$$

where Γ_U sends a complex of presheaves \mathcal{F} to $\mathcal{F}(U)$ and C_U is the constant presheaf functor sending C to the presheaf

$$V \mapsto \begin{cases} C & \text{if } V \subseteq U \\ 0 & \text{otherwise} \end{cases}.$$

If we equip $\text{dgPSh}^{\text{aff}}(X)$ with the projective model structure, then Γ_U preserves fibrations and weak equivalences by construction. Therefore we obtain a Quillen adjunction. We can compose this with the Quillen adjunction

$$\text{dgPSh}^{\text{aff}}(X)^{\text{proj}} \xrightleftharpoons[\text{id}]{\text{id}} \text{dgPSh}^{\text{aff}}(X)^{\text{loc}}$$

of the Bousfield localization to obtain a Quillen adjunction

$$\text{Ch}(\mathbb{k}) \xrightleftharpoons[\Gamma_U]{C_U} \text{dgPSh}^{\text{aff}}(X)^{\text{loc}}.$$

Since C_U is left Quillen, it preserves weak equivalences between cofibrant objects. But every object in $\text{Ch}(\mathbb{k})$ is cofibrant, so C_U preserves weak equivalences.

The same argument works if we instead consider the site of all opens $\text{Open}(X)$. In this case, for any open $V \subseteq X$ we obtain a Quillen adjunction $C_V \dashv \Gamma_V$. Note that in particular we can then take $U = X$. If V is affine, this agrees with the above construction.

The functors C_V and Γ_V are both strong monoidal, since the tensor product of presheaves is taken section-wise. In particular, we obtain lax monoidal functors of ∞ -categories $C_V : \mathcal{D}_{\infty}(\mathbb{k})^{\otimes} \rightarrow \text{Sh}_{\infty}(X)^{\otimes}$ and $\mathbb{R}\Gamma_V : \text{Sh}_{\infty}(X)^{\otimes} \rightarrow \mathcal{D}_{\infty}(\mathbb{k})^{\otimes}$.

Definition 5.49. Let \mathcal{O} be a dg operad. The corresponding operad in $\mathrm{dgPSh}(X)$ is given by $C_X(\mathcal{O})$. By abuse of notation, we will usually denote the operad $C_X(\mathcal{O})$ just by \mathcal{O} .

Lemma 5.50. *The functor C_X preserves cofibrancy and Σ -cofibrancy of operads, as well as weak equivalences of operads. Every operad in $\mathrm{dgPSh}(X)$ is admissible, and even strongly admissible if it is in the image of C_X .*

Proof. The Quillen adjunction $C_X \dashv \Gamma_X$ induces adjunctions between the respective categories of symmetric collections and symmetric operads, since both are strong symmetric monoidal. The model structure on symmetric collection is transferred from the underlying model category, and hence the adjunction is again Quillen. Similarly, fibrations and weak equivalences of operads are pointwise, and hence Γ_X preserves fibrations and trivial fibrations of operads. To see that C_X preserves weak equivalences, note that these are point-wise in operads, and C_X preserves weak equivalences on the underlying model categories. To see that operads in $\mathrm{dgPSh}(X)$ are admissible, use [PS18a, Theorem 5.11] and Section 8 of [PS18b]. Now to see that every operad in the image of C_X is even strongly admissible, use [PS18a, Proposition 6.3] together with the fact that any operad in $\mathrm{Ch}(\mathbb{k})$ is Σ -cofibrant. \square

This shows that we get a diagram of admissible Σ -cofibrant operads

$$\begin{array}{ccc} & C_X(C_*(\mathbb{E}_1)) & \\ C_X(\psi), \simeq \nearrow & \downarrow C_X(\phi), \simeq & \\ C_X(\mathbb{A}_\infty) & \xrightarrow{C_X(\theta), \simeq} & C_X(\mathbf{Assoc}) \end{array}$$

and $C_X(\mathbb{A}_\infty)$ is still cofibrant.

We can hence form the $C_*(\mathbb{E}_1)$ -algebra $\phi^*\mathcal{O}_X$ and choose a cofibrant replacement $\tilde{\mathcal{O}}_X \xrightarrow{\simeq} \phi^*\mathcal{O}_X$. Then we have $\tilde{\mathcal{O}}_X \in \mathrm{Alg}_{\mathbb{E}_1}(\mathrm{Sh}_\infty(X))$ and can consider the center

$$\mathfrak{Z}(\tilde{\mathcal{O}}_X) \in \mathrm{Alg}_{\mathbb{E}_1}(\mathrm{Alg}_{\mathbb{E}_1}(\mathrm{Sh}_\infty(X))) \simeq \mathrm{Alg}_{\mathbb{E}_2}(\mathrm{Sh}_\infty(X)).$$

The Rectification Theorem 4.28 applies to \mathbb{E}_2 algebras in the ∞ -category of sheaves on X . In particular, $\mathrm{dgPSh}(X)$ is symmetrically flat by the proof of Lemma 5.50. This means that we can strictify this \mathbb{E}_2 -algebra structure to the category of complexes of presheaves.

The remainder of this chapter is dedicated to arguing that this center is the correct Hochschild complex of X . In particular, we will show that for a smooth scheme this recovers the sheaf of polydifferential operators.

5.2. The Hochschild complex of a scheme is local. In this section we will prove the following theorem, showing that the center of a scheme glues together the affine Hochschild complexes.

Theorem 5.51. *Let $U = \mathrm{Spec}(A) \subseteq X$ be an affine open. The map $\mathbb{R}\Gamma_U : \mathrm{Sh}_\infty(X) \rightarrow \mathcal{D}_\infty(\mathbb{k})$ is lax symmetric monoidal and hence induces a map $\mathbb{R}\Gamma_U : \mathrm{Alg}_{\mathbb{E}_2}(\mathrm{Sh}_\infty(X)) \rightarrow \mathrm{Alg}_{\mathbb{E}_2}(\mathcal{D}_\infty(\mathbb{k}))$. We have*

$$\mathbb{R}\Gamma_U(\mathfrak{Z}(\tilde{\mathcal{O}}_X)) \simeq \mathfrak{Z}(\tilde{A}) \in \mathrm{Alg}_{\mathbb{E}_2}(\mathcal{D}_\infty(\mathbb{k}))$$

for any cofibrant replacement \tilde{A} of A .

Just like in the affine case, we have

$$\mathfrak{Z}(\tilde{\mathcal{O}}_X)(\mathfrak{a}) \simeq \text{Mor}_{\text{Mod}_{\tilde{\mathcal{O}}_X}^{\mathbb{E}_1}(\text{Sh}_\infty(X) \times_{\mathcal{F}\text{in}_*} \mathbb{E}_1)_\mathfrak{a}}(\tilde{\mathcal{O}}_X, \tilde{\mathcal{O}}_X) \in \text{Sh}_\infty(X),$$

and it hence suffices to understand this endomorphism object. To this end, note that we can adapt Hinich's Rectification Theorem for modules [Hin15, Theorem 5.2.3] to the local projective model structure on complexes of presheaves to get an equivalence of ∞ -categories

$$N(\text{Mod}_{\tilde{\mathcal{O}}_X}^{C_*(\mathbb{E}_1)}(\text{dgPSh}(X)^c)[W_{\text{Mod}}^{-1}] \simeq \text{Mod}_{\tilde{\mathcal{O}}_X}^{\mathbb{E}_1}(\text{Sh}_\infty(X) \times_{\mathcal{F}\text{in}_*} \mathbb{E}_1)_\mathfrak{a}.$$

Again following the affine case, we have a Quillen equivalence

$$\text{Mod}_{\tilde{\mathcal{O}}_X}^{C_*(\mathbb{E}_1)}(\text{dgPSh}(X)) \cong \text{LMod}_{U_{C_*(\mathbb{E}_1)}(\tilde{\mathcal{O}}_X)}(\text{dgPSh}(X)).$$

Proposition 5.52. *There exists a zig-zag of weak equivalences between $U_{C_*(\mathbb{E}_1)}(\tilde{\mathcal{O}}_X)$ and $\mathcal{O}_X \otimes \mathcal{O}_X$ in the category of associative algebras in $\text{dgPSh}(X)$.*

Proof. We adapt the proof of Proposition 4.37. Let $\mathcal{O}'_X \xrightarrow{\sim} \mathcal{O}_X$ be a cofibrant resolution of **Assoc**-algebras in $\text{dgPSh}(X)$. Then

$$(\mathcal{O}'_X \otimes \mathcal{O}'_X)_x \xrightarrow{\cong} \mathcal{O}'_{X,x} \otimes \mathcal{O}'_{X,x} \xrightarrow{\sim} \mathcal{O}_{X,x} \otimes \mathcal{O}_{X,x} \xrightarrow{\cong} (\mathcal{O}_X \otimes \mathcal{O}_X)_x,$$

showing that $\mathcal{O}'_X \otimes \mathcal{O}'_X$ is weakly equivalent to $\mathcal{O}_X \otimes \mathcal{O}_X$. The rest of the argument goes through exactly like before with the following amendment: Let $\hat{\mathcal{O}}_X \xrightarrow{\sim} \theta^* \mathcal{O}'_X$ be a cofibrant replacement of \mathbb{A}_∞ -algebras. To obtain the zig-zag

$$\begin{array}{ccc} & U_{\mathbb{A}_\infty}(\hat{\mathcal{O}}_X) & \\ \swarrow \simeq & & \searrow \simeq \\ U_{\mathbb{A}_\infty}(\theta^* \mathcal{O}'_X) & & U_{\mathbb{A}_\infty}(\psi^* \psi_! \hat{\mathcal{O}}_X) \end{array}$$

we have to argue that the underlying complexes of presheaves of these \mathbb{A}_∞ -algebras are cofibrant. To this end, recall from Lemma 5.50 that \mathbb{A}_∞ and **Assoc** are both strongly admissible, meaning that the forgetful functor from algebras preserves cofibrant objects. In particular, the underlying complexes of presheaves of \mathcal{O}'_X and $\hat{\mathcal{O}}_X$ are both cofibrant. Now recall that $\psi_!$ is left Quillen and hence preserves cofibrancy, and the restriction of scalars functors θ^* and ψ^* do not alter the underlying complex. This finishes the argument. \square

In particular, the category $\text{LMod}_{U_{C_*(\mathbb{E}_1)}(\tilde{\mathcal{O}}_X)}(\text{dgPSh}(X))$ is Quillen equivalent to the category $\text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}(X))$.

Lemma 5.53. *The category $\text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}(X))$ is a dg model category that is left tensored over $\text{dgPSh}(X)$. For $\mathcal{M}, \mathcal{N} \in \text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}(X))$, we have a morphism object*

$$\text{Hom}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{M}, \mathcal{N})(V) = \text{Hom}_{\mathcal{O}_V \otimes \mathcal{O}_V}(\mathcal{M}|_V, \mathcal{N}|_V) \in \text{dgPSh}(X).$$

Proof. Since $\text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}(X))$ is the category of complexes of presheaves of $\mathcal{O}_X \otimes \mathcal{O}_X$ -modules, the dg enrichment is clear. Now note that $\text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}(X))$ is tensored and powered over $\text{dgPSh}(X)$: If $\mathcal{F} \in \text{dgPSh}(X)$ and $\mathcal{M} \in \text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{Ch}(\text{PSh}(\text{Open}(X))_\mathbb{k}))$, we obtain an $\mathcal{O}_X \otimes \mathcal{O}_X$ -module structure on the tensor product in complexes of presheaves by

$$(\mathcal{O}_X \otimes \mathcal{O}_X) \otimes (\mathcal{F} \otimes \mathcal{M}) \cong \mathcal{F} \otimes \mathcal{O}_X \otimes \mathcal{O}_X \otimes \mathcal{M} \rightarrow \mathcal{F} \otimes \mathcal{M}.$$

We obtain a module structure on $\mathcal{H}om(\mathcal{F}, \mathcal{M})$ by the pointwise module structure

$$\mathcal{O}_X \otimes \mathcal{O}_X \otimes \mathcal{H}om(\mathcal{F}, \mathcal{M}) \rightarrow \mathcal{H}om(\mathcal{M}, \mathcal{M}) \otimes \mathcal{H}om(\mathcal{F}, \mathcal{M}) \xrightarrow{\text{comp}} \mathcal{H}om(\mathcal{F}, \mathcal{M}).$$

One easily checks that these indeed yield a tensoring and powering respectively. Now to make the module category into a dg category, we precompose these operations with the constant presheaf functor $C_X : \text{Ch}(\mathbb{k}) \rightarrow \text{dgPSh}(X)$. The pushout-product axiom is checked in [Hin05, Lemma 1.6.3], and Hinich also shows in the same section that the above yields a morphism object. \square

We can now again use Theorem 4.41 to conclude that the center of $\tilde{\mathcal{O}}_X$ in the ∞ -category $\text{Alg}_{\mathbb{E}_1}(\text{Sh}_{\infty}(X))$ is given by

$$\mathfrak{Z}(\tilde{\mathcal{O}}_X)(\mathfrak{a}) \simeq Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{O}, \mathcal{O})$$

for a bifibrant model \mathcal{O} of \mathcal{O}_X as an $\mathcal{O}_X \otimes \mathcal{O}_X$ -module. The center action is given by the evaluation map

$$R(Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{O}, \mathcal{O}) \otimes \mathcal{O}) \rightarrow \mathcal{O}.$$

This is in contrast to the affine case, where A was already fibrant as an $A \otimes A$ -module. The reason for this is that \mathcal{O}_X is not fibrant in the local projective model structure on $\text{dgPSh}(X)$. It is however fibrant in the local projective model structure for the site of affine opens on X , and we have already seen that presheaves on this smaller site present the same ∞ -category.

Proposition 5.54. *We have an equivalence*

$$\iota_*(Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{O}, \mathcal{O})) \simeq Q\mathcal{H}om_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\mathcal{P}, \mathcal{P})$$

for any cofibrant resolution $\mathcal{P} \xrightarrow{\simeq} \iota_*\mathcal{O}_X$ in $\text{LMod}_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\text{dgPSh}^{\text{aff}}(X))$.

Lemma 5.55. *We have a Quillen equivalence*

$$\text{LMod}_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\text{dgPSh}^{\text{aff}}(X)) \xrightleftharpoons[\iota_*]{\iota^*} \text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}(X)).$$

Both adjoints preserve weak equivalences.

Proof. The Quillen equivalence $\iota^{-1} \dashv \iota_*$ induces a Quillen equivalence

$$\text{LMod}_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\text{dgPSh}^{\text{aff}}(X)) \xrightleftharpoons[\iota_*]{\iota^{-1}} \text{LMod}_{\iota^{-1}\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\text{dgPSh}(X)).$$

The counit yields a weak equivalence $\epsilon : \iota^{-1}\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X) \xrightarrow{\simeq} \mathcal{O}_X \otimes \mathcal{O}_X$, and hence we get a Quillen equivalence

$$\text{LMod}_{\iota^{-1}\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\text{dgPSh}(X)) \xrightleftharpoons[\epsilon_*]{\epsilon^*} \text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}(X)).$$

composing these yields the Quillen equivalence in the statement. Clearly, ϵ_* preserves weak equivalences. But ϵ is an isomorphism on stalks, and hence ϵ^* also preserves weak equivalences. \square

Lemma 5.56. *For $\mathcal{F} \in \text{LMod}_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\text{dgPSh}^{\text{aff}}(X))$ and $\mathcal{G} \in \text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}(X))$, we have*

$$\iota_*\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\iota^*\mathcal{F}, \mathcal{G}) \cong \mathcal{H}om_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\mathcal{F}, \iota_*\mathcal{G}).$$

Proof. Let $U \in \text{Aff}(X)$, and let $\iota' : \text{Aff}(U) \rightarrow \text{Open}(U)$ be the inclusion. Then

$$\begin{aligned} \text{Hom}_{\mathcal{O}_U \otimes \mathcal{O}_U}(\iota^* \mathcal{F}|_U, \mathcal{G}|_U) &\cong \text{Hom}_{\mathcal{O}_U \otimes \mathcal{O}_U}(\iota'^*(\mathcal{F}|_U), \mathcal{G}|_U) \\ &\cong \text{Hom}_{\iota'_*(\mathcal{O}_U \otimes \mathcal{O}_U)}(\mathcal{F}|_U, \iota'_*(\mathcal{G}|_U)) \\ &\cong \text{Hom}_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)|_U}(\mathcal{F}|_U, \iota_* \mathcal{G}|_U). \end{aligned}$$

□

Proof of Proposition 5.54. By definition, we have a diagram

$$\begin{array}{c} \mathcal{P}' \xrightarrow{\simeq} \mathcal{O}_X \\ \simeq \downarrow \\ \mathcal{O} \end{array}$$

with \mathcal{P}' cofibrant and \mathcal{O} bifibrant. If $\mathcal{P} \xrightarrow{\simeq} \iota_* \mathcal{O}_X$ is a cofibrant resolution, we get a weak equivalence $\iota^* \mathcal{P} \xrightarrow{\simeq} \iota^* \iota_* \mathcal{O}_X$ and $\iota^* \mathcal{P}$ is still cofibrant. We can hence solve the following lifting problem

$$\begin{array}{ccc} 0 & \xrightarrow{\quad} & \mathcal{P}' \\ \downarrow & \nearrow \text{dashed} & \downarrow \simeq \\ \iota^* \mathcal{P} & \xrightarrow{\simeq} \iota^* \iota_* \mathcal{O}_X \xrightarrow{\simeq} & \mathcal{O}_X \end{array}$$

and by 2-out-of-3, the lift $\iota^* \mathcal{P} \rightarrow \mathcal{P}'$ is again a weak equivalence. Let $\iota^* \mathcal{P} \xrightarrow{\simeq} \mathcal{R}$ be a fibrant resolution. We can also solve the lifting problem

$$\begin{array}{ccc} \iota^* \mathcal{P} & \xrightarrow{\simeq} \mathcal{P}' \xrightarrow{\simeq} & \mathcal{O} \\ \simeq \downarrow & \nearrow \text{dashed} & \downarrow \\ \mathcal{R} & \xrightarrow{\quad} & 0 \end{array}$$

to obtain a weak equivalence $\mathcal{R} \xrightarrow{\simeq} \mathcal{O}$. In particular, this is a weak equivalence between bifibrant objects. Therefore,

$$\begin{aligned} \iota_* Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{O}, \mathcal{O}) &\simeq \iota_* Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{R}, \mathcal{R}) \\ &\simeq \iota_* Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\iota^* \mathcal{P}, \mathcal{R}) \\ &\simeq Q\mathcal{H}om_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\mathcal{P}, \iota_* \mathcal{R}) \\ &\simeq Q\mathcal{H}om_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\mathcal{P}, \mathcal{P}), \end{aligned}$$

where in the last step we used that $\iota_* \iota^*(\mathcal{P}) \cong \mathcal{P}$ and that hence $\iota_* \mathcal{R} \leftarrow \iota_* \iota^* \mathcal{P}$ is a weak equivalence between fibrant objects. □

Since ι_* is symmetric monoidal, we get an induced \mathbb{E}_2 -algebra structure on $Q\mathcal{H}om_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\mathcal{P}, \mathcal{P})$, and in view of Theorem 5.51, the above argument shows that if $U \subseteq X$ is affine, then

$$\mathbb{R}\Gamma_U(\mathfrak{Z}(\tilde{\mathcal{O}}_X)) \simeq \mathbb{R}\Gamma_U(Q\mathcal{H}om_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\mathcal{P}, \mathcal{P}))$$

as \mathbb{E}_2 -algebras. It now suffices to show that

$$\mathbb{R}\Gamma_U(Q\mathcal{H}om_{\iota_*(\mathcal{O}_X \otimes \mathcal{O}_X)}(\mathcal{P}, \mathcal{P})) \simeq \text{Hom}_{A \otimes A}(P, P)$$

for $U = \text{Spec}(A)$ and $P \xrightarrow{\sim} A$ a cofibrant replacement. Since we will solely work with the affine open site from now on, we will denote the restriction of a presheaf \mathcal{F} on $\text{Open}(X)$ to a presheaf on $\text{Aff}(X)$ simply by \mathcal{F} for the remainder of this section.

Definition 5.57. Let Diag_X be the site of affine opens on $X \times_{\mathbb{k}} X$ of the form $W \times_{\mathbb{k}} W$ for $W \subseteq X$ affine open. Of course, this site is isomorphic to the affine open site on X , but it better conceptualizes sheaves coming from A -bimodules.

Lemma 5.58. (1) *The maps*

$$\Delta_* : \text{dgPSh}^{\text{aff}}(X) \rightarrow \text{dgPSh}(\text{Diag}_X), \quad \mathcal{F} \mapsto (W \times_{\mathbb{k}} W \mapsto \mathcal{F}(\Delta^{-1}(W \times_{\mathbb{k}} W)) = \mathcal{F}(W)) \text{ and}$$

$$\Delta^{-1} : \text{dgPSh}(\text{Diag}_X) \rightarrow \text{dgPSh}^{\text{aff}}(X), \quad \mathcal{G} \mapsto (U \mapsto \text{colim}_{\Delta(U) \subseteq W \times_{\mathbb{k}} W} \mathcal{G}(W \times_{\mathbb{k}} W) \cong \mathcal{G}(U \times_{\mathbb{k}} U))$$

form an isomorphism of categories.

(2) *We have $\Delta_*(\mathcal{O}_X \otimes \mathcal{O}_X) \cong \mathcal{O}_{X \times_{\mathbb{k}} X}$ and $\Delta^{-1}(\mathcal{O}_{X \times_{\mathbb{k}} X}) \cong \mathcal{O}_X \otimes \mathcal{O}_X$. In particular, the above isomorphism yields an isomorphism*

$$\text{LMod}_{\mathcal{O}_{X \times_{\mathbb{k}} X}}(\text{dgPSh}(\text{Diag}_X)) \cong \text{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\text{dgPSh}^{\text{aff}}(X)).$$

(3) *Both Δ_* and Δ^{-1} preserve all three classes of fibrations, cofibrations and weak equivalences in the presheaf categories as well as the left module categories.*

(4) *For any commutative k -algebra A , the adjunction $(\tilde{-}) \vdash \Gamma_{\text{Spec}(A)}$ between complexes of A -modules and complexes of presheaves of $\mathcal{O}_{\text{Spec}(A)}$ -modules is a Quillen adjunction. In addition, $(\tilde{-})$ preserves acyclic fibrations.*

(5) *The previous statement remains true if we consider $(\tilde{-})$ as a functor from complexes of $A \otimes_{\mathbb{k}} A$ -modules to complexes of presheaves of $\mathcal{O}_{\text{Spec}(A)}$ -modules on the site $\text{Diag}_{\text{Spec}(A)}$.*

Proof. Statement 1. is true by construction of the functors.

For 2., simply note that $\mathcal{O}_{X \times_{\mathbb{k}} X}(W \times_{\mathbb{k}} W) \cong \mathcal{O}_X(W) \otimes_{\mathbb{k}} \mathcal{O}_X(W) \cong (\mathcal{O}_X \otimes \mathcal{O}_X)(W)$.

For 3., note that we get two adjoint equivalences $\Delta^{-1} \dashv \Delta_*$ and $\Delta_* \dashv \Delta^{-1}$. Clearly both Δ^{-1} and Δ_* preserve acyclic fibrations, and hence both also preserve cofibrations. Since Diag_X is isomorphic as a site to $\text{Aff}(X)$, the sheaf topos $\text{Sh}(\text{Diag}_X)_{\mathbb{k}}$ also has enough points and hence we can check weak equivalences at stalks. But at the same time, the only points in the sheaf topos on Diag_X are of the form $\Delta(x)$ for $x \in X$, and $(\Delta_* \mathcal{F})_{\Delta(x)} \cong \mathcal{F}_x$ and $(\Delta^{-1} \mathcal{G})_x \cong \mathcal{G}_x$, proving that both Δ^{-1} and Δ_* preserve weak equivalences.

For 4., first note that this is indeed an adjunction. To see this, let M be a complex of A -modules and consider a map $M \rightarrow \mathcal{F}(\text{Spec}(A))$. If $U = \text{Spec}(B)$ is an affine open of $X = \text{Spec}(A)$, then we get a restriction map $\mathcal{F}(X) \rightarrow \mathcal{F}(\text{Spec}(B))$ which is a map of A -modules. But $\mathcal{F}(\text{Spec}(B))$ is a B -module, and hence we get a map $\mathcal{F}(X) \otimes_A B \rightarrow \mathcal{F}(\text{Spec}(B))$. We can hence construct a map

$$\tilde{M}(\text{Spec}(B)) \cong M \otimes_A B \rightarrow \mathcal{F}(X) \otimes_A B \rightarrow \mathcal{F}(\text{Spec}(B)).$$

of B -modules. Now note that $(\tilde{-})$ sends quasi-isomorphisms to pointwise weak equivalences: If $M \rightarrow N$ is a quasi-isomorphism of complexes of A -modules and $U = \text{Spec}(B) \subseteq \text{Spec}(A)$ is an affine open, then in particular B is flat over A and therefore $- \otimes_A B$ preserves quasi-isomorphisms. Hence $(\tilde{M})(U) = M \otimes_A B \rightarrow N \otimes_A B = \tilde{N}(U)$ is again a quasi-isomorphism. Further, we already know that the global sections functor preserves acyclic fibrations. This shows that the above adjunction is Quillen. Now if $M \rightarrow N$ is an acyclic fibration, then so is $M \otimes_A B \rightarrow N \otimes_A B$. This finishes the proof.

For 5., just note that everything in the proof of 4. still works. \square

Proof of Theorem 5.51. We work over the affine open site. Let $\mathcal{P} \xrightarrow{\simeq} \mathcal{O}_X$ be a cofibrant resolution in $\mathcal{O}_X \otimes \mathcal{O}_X$ modules. The $\mathcal{O}_X \otimes \mathcal{O}_X$ -module \mathcal{P} is bifibrant and $\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(-, -)$ is a right Quillen bifunctor, implying that $Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P})$ is again bifibrant in $\mathrm{dgPSh}^{\mathrm{aff}}(X)$. We hence get a weak equivalence

$$\mathbb{R}\Gamma_U(Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P})) \simeq \mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P})(U) \cong \mathrm{Hom}_{\mathcal{O}_U \otimes \mathcal{O}_U}(\mathcal{P}|_U, \mathcal{P}|_U)$$

We then have the following chain of weak equivalences

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_U \otimes \mathcal{O}_U}(\mathcal{P}|_U, \mathcal{P}|_U) &\cong \mathrm{Hom}_{\mathcal{O}_{U \times_{\mathbb{k}} U}}((\Delta_U)_*(\mathcal{P}|_U), (\Delta_U)_*(\mathcal{P}|_U)) \\ &\cong \mathrm{Hom}_{\mathcal{O}_{X \times_{\mathbb{k}} X}|_{U \times_{\mathbb{k}} U}}(\Delta_*(\mathcal{P})|_{U \times_{\mathbb{k}} U}, \Delta_*(\mathcal{P})|_{U \times_{\mathbb{k}} U}). \end{aligned}$$

By the above lemma $\Delta_*\mathcal{P}$ is again bifibrant, and $(\Delta_*\mathcal{P})|_{U \times_{\mathbb{k}} U}$ is fibrant. We can therefore use proposition [Hin05, 1.7.3] with a choice of cofibrant resolution $\mathcal{P}' \xrightarrow{\simeq} (\Delta_*(\mathcal{P}))|_{U \times_{\mathbb{k}} U}$ to get weak equivalences

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_{X \times_{\mathbb{k}} X}|_{U \times_{\mathbb{k}} U}}(\Delta_*(\mathcal{P})|_{U \times_{\mathbb{k}} U}, \Delta_*(\mathcal{P})|_{U \times_{\mathbb{k}} U}) &\xrightarrow{\simeq} \mathrm{Hom}_{\mathcal{O}_{X \times_{\mathbb{k}} X}|_{U \times_{\mathbb{k}} U}}(\mathcal{P}', \Delta_*(\mathcal{P})|_{U \times_{\mathbb{k}} U}) \\ &\xleftarrow{\simeq} \mathrm{Hom}_{\mathcal{O}_{X \times_{\mathbb{k}} X}|_{U \times_{\mathbb{k}} U}}(\mathcal{P}', \mathcal{P}') \end{aligned}$$

Note that $(\Delta_*\mathcal{P})|_{U \times_{\mathbb{k}} U} \xrightarrow{\simeq} (\Delta_U)_*\mathcal{O}_U \cong \tilde{A}$ is again a trivial fibration, and therefore $\mathcal{P}' \xrightarrow{\simeq} \tilde{A}$ is a cofibrant resolution in $\mathcal{O}_{U \times_{\mathbb{k}} U}$ -modules. Now let $P \xrightarrow{\simeq} A$ be a cofibrant resolution of A as an A^e -module. Then \tilde{P} is a cofibrant $\mathcal{O}_{U \times_{\mathbb{k}} U}$ -module, and we hence get a weak equivalence $\tilde{P} \xrightarrow{\simeq} \mathcal{P}'$ between bifibrant objects. Therefore,

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_{U \times_{\mathbb{k}} U}}(\mathcal{P}', \mathcal{P}') &\xrightarrow{\simeq} \mathrm{Hom}_{\mathcal{O}_{U \times_{\mathbb{k}} U}}(\tilde{P}, \mathcal{P}') \\ &\xleftarrow{\simeq} \mathrm{Hom}_{\mathcal{O}_{U \times_{\mathbb{k}} U}}(\tilde{P}, \tilde{P}) \\ &\cong \mathrm{Hom}_{A \otimes A}(P, P). \end{aligned}$$

This proves that $\mathbb{R}\Gamma_U(\mathfrak{Z}(\tilde{\mathcal{O}}_X)(\mathfrak{a})) \simeq \mathfrak{Z}(\tilde{A})(\mathfrak{a})$ as complexes of \mathbb{k} -modules. Recall that $\mathbb{R}\Gamma_U$ is lax symmetric monoidal, and therefore we get an induced evaluation map

$$Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P})(U) \otimes \mathcal{P}(U) \rightarrow R(Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P})(U) \otimes \mathcal{P}(U)) \rightarrow \mathcal{P}(U)$$

But $\mathcal{P}(U) \simeq \mathcal{O}_X(U) \cong A \simeq P$, so this is in fact equivalent to the evaluation map

$$\mathrm{Hom}_{A \otimes A}(P, P) \otimes P \rightarrow P$$

of the center of A . This shows that the above equivalence is indeed an equivalences of \mathbb{E}_2 -algebras. \square

5.3. Recovering polydifferential operators as the center of \mathcal{O}_X . Now suppose that X is also smooth. Recall that in this case the Hochschild cohomology of X is given by the hypercohomology of the sheaf of polydifferential operators $\mathcal{D}_{\mathrm{poly}}(X) \in \mathrm{dgSh}(X)$. If $U = \mathrm{Spec}(A) \subseteq X$ is an affine open, then

$$\begin{aligned} \mathcal{D}_{\mathrm{poly}}(X)_n(U) &= \{f \in \mathrm{Hom}_{\mathbb{k}}(A^{\otimes n}, A) : f \text{ is a differential operator in each factor}\} \\ &\subseteq \mathrm{Hom}_{\mathbb{k}}(A^{\otimes n}, A). \end{aligned}$$

The sheaf of polydifferential operators is quasi-coherent as an \mathcal{O}_X -module, and therefore fibrant in the local projective model structure on affine opens. We want to show that the sheaf of polydifferential operators is indeed a model of the center of \mathcal{O}_X .

Theorem 5.59. *Let X be a smooth, quasi-compact, separated scheme of finite type over \mathbb{k} . We have an equivalence*

$$Q\mathcal{D}_{\text{poly}}(X) \simeq \mathfrak{Z}(\tilde{\mathcal{O}}_X)(\mathfrak{a})$$

in the ∞ -category $\text{LMod}_{\tilde{\mathcal{O}}_X}(\text{Sh}_{\infty}(X))$ of $\tilde{\mathcal{O}}_X$ -modules.

It suffices to show this equivalence for the sites of affine opens, since they yield an equivalent ∞ -category. Since $\iota_*\mathfrak{Z}(\tilde{\mathcal{O}}_X)(\mathfrak{a}) \simeq Q\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P})$, it suffices to show

$$\iota_*\mathcal{D}_{\text{poly}}(X) \simeq \mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P})$$

as presheaves of \mathcal{O}_X -modules. In the following we will suppress the restriction to affine opens.

Let \mathcal{O} be an associative algebra in complexes of sheaves. If \mathcal{F}, \mathcal{G} are sheaves of left \mathcal{O} -modules, recall that $\mathbb{R}\mathcal{H}om_{\mathcal{O}}(\mathcal{F}, \mathcal{G}) = \mathcal{H}om_{\mathcal{O}}(\mathcal{F}, \mathcal{J})$ for some K -injective resolution \mathcal{J} of \mathcal{G} in the category of sheaves of left \mathcal{O} -modules. Let $\Delta : X \rightarrow X \times_{\mathbb{k}} X$ be the diagonal. We have already used the adjunction $\Delta^{-1} \dashv \Delta_*$ induced by this in Lemma 5.58 above, but we now want to consider the full site of affine opens on $X \times_{\mathbb{k}} X$ instead of the smaller site Diag_X , and we also consider sheaves instead of presheaves. In particular, the map $\Delta^{-1} : \text{dgSh}^{\text{aff}}(X \times_{\mathbb{k}} X) \rightarrow \text{dgSh}(X)$ is now given by the presheaf version followed by sheafification. We then have $\Delta^{-1}\Delta_* \cong \text{id}$ since X is separated. Denote by $\overset{a}{\otimes}$ the tensor product of sheaves.

Lemma 5.60. (1) *We have a local quasi-isomorphism of complexes of presheaves*

$$\mathcal{H}om_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P}) \simeq \mathbb{R}\mathcal{H}om_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X)$$

(2) *If \mathcal{F} and \mathcal{G} are sheaves, then $\Delta_*\mathcal{H}om_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\Delta^{-1}\mathcal{F}, \mathcal{G}) \cong \mathcal{H}om_{\mathcal{O}_X \times_{\mathbb{k}} X}(\mathcal{F}, \Delta_*\mathcal{G})$.*

(3) *If $\mathcal{O}_X \xrightarrow{\sim} \mathcal{I}$ is a K -injective resolution in sheaves of $\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X$ -modules, then $\Delta_*\mathcal{O}_X \rightarrow \Delta_*\mathcal{I}$ is a K -injective resolution in $\mathcal{O}_X \times_{\mathbb{k}} X$ -modules.*

Assuming this lemma, we can prove the theorem as follows.

Proof of Theorem 5.59. By [Yek02, Corollary 2.9] we have a local weak equivalence

$$\Delta_*\mathcal{D}_{\text{poly}}(X) \simeq \mathbb{R}\mathcal{H}om_{\mathcal{O}_X \times_{\mathbb{k}} X}(\Delta_*\mathcal{O}_X, \Delta_*\mathcal{O}_X).$$

We then get the following chain of local weak equivalences

$$\begin{aligned} \Delta_*\mathbb{R}\mathcal{H}om_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X) &= \Delta_*\mathcal{H}om_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\mathcal{O}_X, \mathcal{I}) \\ &\cong \Delta_*\mathcal{H}om_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\Delta^{-1}\Delta_*\mathcal{O}_X, \mathcal{I}) \\ &\cong \mathcal{H}om_{\mathcal{O}_X \times_{\mathbb{k}} X}(\Delta_*\mathcal{O}_X, \Delta_*\mathcal{I}) \\ &\simeq \mathbb{R}\mathcal{H}om_{\mathcal{O}_X \times_{\mathbb{k}} X}(\Delta_*\mathcal{O}_X, \Delta_*\mathcal{O}_X) \\ &\simeq \Delta_*\mathcal{D}_{\text{poly}}(X) \end{aligned}$$

where in the second to last step we used 5.60(3.). Now note that Δ^{-1} preserves local weak equivalences, and therefore

$$\mathbb{R}\mathcal{H}om_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X) \simeq \mathcal{D}_{\text{poly}}(X).$$

Together with 5.60(1.) this finishes the proof. \square

Proof of Lemma 5.60. For 1., let $\alpha : \mathcal{O}_X \otimes \mathcal{O}_X \xrightarrow{\sim} \mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X$ be the unit of the sheafification adjunction. This is a weak equivalence of dg algebras in presheaves, and therefore induces a Quillen equivalence

$$\mathrm{LMod}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathrm{dgPSh}^{\mathrm{aff}}(X)) \xrightleftharpoons[\alpha_*]{\alpha^*} \mathrm{LMod}_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\mathrm{dgPSh}^{\mathrm{aff}}(X)) .$$

We therefore get the following chain of weak equivalences

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P}) &\simeq \mathrm{Hom}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \alpha_* \mathcal{O}_X) \\ &\cong \mathrm{Hom}_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\alpha^* \mathcal{P}, \mathcal{O}_X) \\ &\simeq \mathrm{Hom}_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\alpha^* \mathcal{P}, \mathcal{I}) \\ &\cong \mathrm{Hom}_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}((\alpha^* \mathcal{P})^a, \mathcal{I}) \\ &\simeq \mathrm{Hom}_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\mathcal{O}_X, \mathcal{I}) \\ &= \mathbb{R}\mathrm{Hom}_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X). \end{aligned}$$

The 2. statement is standard.

For the 3. statement, note first that \mathcal{O}_X and \mathcal{I} are both fibrant in the local projective model structure, and the presheaf version of the $\Delta^{-1} \dashv \Delta_*$ adjunction is Quillen for this model structure on the affine open sites, so $\Delta_* \mathcal{O}_X \rightarrow \Delta_* \mathcal{I}$ is again a local weak equivalence. Further Δ^{-1} is exact, and therefore preserves acyclic complexes. Therefore, if \mathcal{S} is an acyclic $\mathcal{O}_{X \times_k X}$ -module, then

$$\mathrm{Hom}_{\mathcal{O}_{X \times_k X}}(\mathcal{S}, \Delta_* \mathcal{I}) \cong \mathrm{Hom}_{\mathcal{O}_X \overset{a}{\otimes} \mathcal{O}_X}(\Delta^{-1} \mathcal{S}, \mathcal{I})$$

is acyclic, proving that $\Delta_* \mathcal{I}$ is K-injective. \square

Let $\mathcal{B}_\bullet(\mathcal{O}_X)$ denote the $\mathcal{O}_X \otimes \mathcal{O}_X$ -module $U = \mathrm{Spec}(A) \mapsto B_\bullet(A)$. We have a surjective map $\mathcal{B}_\bullet(\mathcal{O}_X) \rightarrow \mathcal{O}_X$ given by multiplication. Hence for a projective resolution $\mathcal{P} \xrightarrow{\sim} \mathcal{O}_X$ of \mathcal{O}_X as an $\mathcal{O}_X \otimes \mathcal{O}_X$ -module, we get a lift $\mathcal{P} \rightarrow \mathcal{B}_\bullet(\mathcal{O}_X)$. We get an evaluation map

$$\iota_* \mathcal{D}_{\mathrm{poly}}(X) \otimes \mathcal{B}_\bullet(\mathcal{O}_X) \rightarrow \iota_* \mathcal{O}_X$$

coming from the fact that $\mathcal{D}_{\mathrm{poly}}(X)$ is affine locally a subcomplex of the Hochschild complex. This lifts to a map

$$\begin{array}{ccc} Q\iota_* \mathcal{D}_{\mathrm{poly}}(X) \otimes \mathcal{P} & \dashrightarrow & \mathcal{P} \\ \downarrow & & \downarrow \simeq \\ \iota_* \mathcal{D}_{\mathrm{poly}}(X) \otimes \mathcal{B}_\bullet(\mathcal{O}_X) & \longrightarrow & \mathcal{O}_X \end{array}$$

and it is clear from the proof of Theorem 5.59 that this map corresponds to the evaluation map

$$Q\mathrm{Hom}_{\mathcal{O}_X \otimes \mathcal{O}_X}(\mathcal{P}, \mathcal{P}) \otimes \mathcal{P} \rightarrow \mathcal{P}.$$

It therefore makes $Q\mathcal{D}_{\mathrm{poly}}(X)$ into a center of $\tilde{\mathcal{O}}_X$. In particular, this equips the sheaf of polydifferential operators with a new \mathbb{E}_2 -algebra structure in the ∞ -category of sheaves on X .

5.4. Comparison to the classical homotopy Gerstenhaber algebra structure on polydifferential operators. For a smooth scheme X , Tamarkin's proof of Deligne's conjecture equips $\mathcal{D}_{\text{poly}}(X)$ with a $\mathcal{G}er_\infty$ -algebra structure coming from the $\mathcal{B}races$ -algebra structure, and thus a Gerstenhaber algebra structure in the \mathbb{k} -linear derived 1-category. On the other hand, exhibiting $\mathcal{D}_{\text{poly}}(X)$ as a center of $\tilde{\mathcal{O}}_X$ equips it with an \mathbb{E}_2 -algebra structure in $\text{Sh}_\infty(X)$, and therefore another Gerstenhaber algebra structure in the derived 1-category, which is just the homotopy category of $\text{Sh}_\infty(X)$. We want to compare these two Gerstenhaber algebra structures. For the remainder of this section, let \otimes denote the tensor product of sheaves instead of presheaves.

To this end, note that by Corollary 4.35, the multiplication in the center Gerstenhaber algebra structure is given equivalently by the convolution product or the composition product, and the bracket is induced by any filling of the appropriate square in the action category.

par

By [Yek02], we have an isomorphism of sheaves

$$\mathcal{D}_{\text{poly}}(X) \cong \mathcal{H}om_{\mathcal{O}_X \times_{\mathbb{k}} X}^{\text{cont}}(\widehat{\mathcal{B}}(X), \mathcal{O}_X)$$

where $\widehat{\mathcal{B}}_n(X) = \mathcal{O}_{X^{n+2}}$ is the complete Bar complex with \mathfrak{X}^n the formal completion of X^n along the diagonal. From this presentation it is easy to compute the convolution and composition product. Note that for $U = \text{Spec}(A) \subseteq X$, we have

$$\Gamma_U(\mathcal{H}om_{\mathcal{O}_X \times_{\mathbb{k}} X}^{\text{cont}}(\widehat{\mathcal{B}}(X), \mathcal{O}_X)) \cong \mathcal{H}om_{A \otimes A}^{\text{cont}}(\widehat{B}(A), A)$$

for $\widehat{B}_n(A)$ the adic completion of $B_n(A)$ at the kernel of the multiplication map $B_n(A) \rightarrow A$. In particular, the flat resolution

$$\widehat{\mathcal{B}}(X) \rightarrow \mathcal{O}_X$$

admits a section $s : \mathcal{O}_X \rightarrow \widehat{\mathcal{B}}_0(X)$ that is glued together from the sections of the resolutions $B(A) \rightarrow A$. We can hence build a diagonal

$$\Delta : \widehat{\mathcal{B}}(X) \rightarrow \mathcal{O}_X \xrightarrow{\cong} \mathcal{O}_X \otimes_{\mathcal{O}_X} \mathcal{O}_X \xrightarrow{s \otimes s} \widehat{\mathcal{B}}(X) \otimes_{\mathcal{O}_X} \widehat{\mathcal{B}}(X).$$

Lemma 5.61. *The convolution product on $\mathcal{H}om_{\mathcal{O}_X \times_{\mathbb{k}} X}^{\text{cont}}(\widehat{\mathcal{B}}(X), \mathcal{O}_X)$ is homotopic to the cup product on $\mathcal{D}_{\text{poly}}(X)$ which locally agrees with the classical cup product on Hochschild cochains.*

Proof. The diagonal is zero on $\widehat{\mathcal{B}}_n(X)$ for $n > 0$, and for $n = 0$ it is given locally by the formula

$$a_0 \otimes a_1 \mapsto (1 \otimes 1) \otimes_A (1 \otimes a_0 a_1).$$

The diagonal on $B(A)$ coming from its universal property is given for $n = 0$ by

$$a_0 \otimes a_1 \mapsto (a_0 \otimes 1) \otimes_A (1 \otimes a_1),$$

and for $n > 0$ by

$$a_0 \otimes a_1 \otimes \cdots \otimes a_n \otimes a_{n+1} \mapsto \sum_{i=0}^n (a_0 \otimes a_1 \otimes \cdots \otimes a_i \otimes 1) \otimes_A (1 \otimes a_{i+1} \otimes \cdots \otimes a_n \otimes a_{n+1}).$$

Locally on $B(A)$, a homotopy between these two maps is given by

$$H : B(A) \rightarrow (B(A) \otimes_A B(A))[1]$$

$$a_0 \otimes a_1 \otimes \cdots \otimes a_n \otimes a_{n+1} \mapsto \sum_{i=0}^{n+1} (1 \otimes a_0 \otimes \cdots \otimes a_{i-1} \otimes 1) \otimes_A (1 \otimes a_i \otimes \cdots \otimes a_{n+1}).$$

Recall that the convolution product of two continuous maps f and g is given by

$$\widehat{B}(X) \xrightarrow{\Delta} \widehat{B}(X) \otimes_{\mathcal{O}} \widehat{B}(X) \xrightarrow{f \otimes g} \mathcal{O}_X \otimes_{\mathcal{O}_X} \mathcal{O}_X \rightarrow \mathcal{O}_X.$$

Locally, it suffices to consider the restriction to $B(A)$. We can then locally use the above homotopy H to obtain a homotopy between the cup product and the above formula for the convolution product with our new diagonal. Inspecting the formula for H we see that these glue together to yield a global homotopy between the global convolution product and the cup product. \square

We already know that the local circle products coming from the *Braces*-algebra structure glue together to give a homotopy for the square

$$\begin{array}{ccc} \mathcal{D}_{\text{poly}}(X)^{\otimes 4} & \xrightarrow{\smile \otimes \smile} & \mathcal{D}_{\text{poly}}(X)^{\otimes 2} \\ (\smile \otimes \smile) \circ (\text{id} \otimes \tau \otimes \text{id}) \downarrow & & \downarrow \smile \\ \mathcal{D}_{\text{poly}}(X)^{\otimes 2} & \xrightarrow{\smile} & \mathcal{D}_{\text{poly}}(X) \end{array}$$

like in the affine case.

Proposition 5.62. *The Gerstenhaber structure on $\mathcal{D}_{\text{poly}}(X)$ coming from the center agrees with the classical one in the \mathbb{k} -linear derived 1-category.*

Proof. By the above Lemma 5.61, the product agrees with the cup product in the derived category. Then using again Corollary 4.35 and the above square, we see that the bracket from the center locally agrees with the classical Gerstenhaber bracket, and hence agrees with the classical one on polydifferential operators. \square

APPENDIX A. THE ENDOMORPHISM ∞ -CATEGORY

We want to show that our endomorphism ∞ -category $\mathcal{C}_a^\otimes \times_{\mathcal{M}} \mathcal{M}_{/M}$ agrees with Lurie's definition [Lur17, Definition 4.7.1.1]. This will show that our endomorphism ∞ -category is the underlying category of a monoidal ∞ -category.

Let $q : \mathcal{C}^\otimes \rightarrow \mathcal{LM}^\otimes$ be a coCartesian fibration of ∞ -operads. In particular, q exhibits $\mathcal{M} := \mathcal{C}_m$ as left-tensored over the monoidal ∞ -category \mathcal{C}_a^\otimes . Construct an ∞ -category \mathcal{M}^\otimes as the fiber product

$$\begin{array}{ccc} \mathcal{M}^\otimes := \mathcal{C}^\otimes \times_{\mathcal{LM}^\otimes} (N(\Delta^{\text{op}}) \times \Delta^1) & \longrightarrow & \mathcal{C}^\otimes \\ \downarrow & & \downarrow q \\ N(\Delta^{\text{op}}) \times \Delta^1 & \xrightarrow{\gamma} & \mathcal{LM}^\otimes \end{array} .$$

In particular, \mathcal{M}^\otimes comes equipped with a coCartesian fibration $p : \mathcal{M}^\otimes \rightarrow N(\Delta^{\text{op}}) \times \Delta^1$. We have

$$\mathcal{M}_{[0],0}^\otimes = \mathcal{C}^\otimes \times_{\mathcal{LM}^\otimes} \{[0], 0\} \simeq \mathcal{C}_m = \mathcal{M},$$

since the functor $\text{LCut} : N(\Delta^{\text{op}}) \rightarrow \mathcal{LM}^\otimes$ sends $[0]$ to $(\langle 1 \rangle, \{1\}) = \mathbf{m}$. Consider the diagram

$$\begin{array}{ccccc} \mathcal{M}^\otimes \times_{\Delta^1} \{1\} & \longrightarrow & \mathcal{C}^\otimes \times_{\mathcal{LM}^\otimes} (N(\Delta^{\text{op}}) \times \Delta^1) & \longrightarrow & \mathcal{C}^\otimes \\ \downarrow & & \downarrow & & \downarrow q \\ N(\Delta^{\text{op}}) \times \{1\} & \hookrightarrow & N(\Delta^{\text{op}}) \times \Delta^1 & \xrightarrow{\gamma} & \mathcal{LM}^\otimes \\ \downarrow & & \downarrow & & \\ \{1\} & \hookrightarrow & \Delta^1 & & \end{array} .$$

Then the upper right hand side square is a pullback by definition, the lower left hand side square is a pullback and the left hand side rectangle is a pullback, again by definition. By the pasting law, the upper left hand side square is a pullback, and hence, again by the pasting law, the large upper rectangle is a pullback. The lower horizontal arrow of this rectangle agrees with the map $\text{Cut} : N(\Delta^{\text{op}}) \rightarrow \mathcal{LM}^\otimes$, so

$$\mathcal{M}^\otimes \times_{\Delta^1} \{1\} \simeq \mathcal{C}^\otimes \times_{\mathcal{LM}^\otimes} N(\Delta^{\text{op}}).$$

But in the diagram

$$\begin{array}{ccccc} \mathcal{C}_a^\otimes \times_{\mathcal{Assoc}^\otimes} N(\Delta^{\text{op}}) & \longrightarrow & \mathcal{C}_a^\otimes & \longrightarrow & \mathcal{C}^\otimes \\ \downarrow & & \downarrow & & \downarrow q \\ N(\Delta^{\text{op}}) & \xrightarrow{\text{Cut}} & \mathcal{Assoc}^\otimes & \hookrightarrow & \mathcal{LM}^\otimes \end{array}$$

both squares are pullbacks, so the rectangle is as well, and we get

$$\mathcal{M}^\otimes \times_{\Delta^1} \{1\} \simeq \mathcal{C}_a^\otimes \times_{\mathcal{Assoc}^\otimes} N(\Delta^{\text{op}}),$$

which is the \mathbb{A}_∞ -monoidal ∞ -category corresponding to the monoidal ∞ -category \mathcal{C}_a^\otimes . Call this \mathbb{A}_∞ -monoidal category \mathcal{C}_a^\otimes . Then p exhibits \mathcal{M} as left-tensored over \mathcal{C}_a^\otimes in the planar sense.

Proposition A.63. *Let $q : \mathcal{C}^\otimes \rightarrow \mathcal{LM}^\otimes$ be a coCartesian fibration of ∞ -operads. Let $p : \mathcal{M}^\otimes \rightarrow N(\Delta^{\text{op}}) \times \Delta^1$ as above. Then the ∞ -category $\mathcal{C}_a[M]$ from Definition 2.5 is equivalent to the endomorphism ∞ -category of M as defined in [Lur17, Definition 4.7.1.1].*

Proof. Under $\gamma : N(\mathbb{A}^{\text{op}}) \times \Delta^1 \rightarrow \mathcal{LM}^\otimes$, the map

$$a : ([0], 0) \rightarrow ([1], 0)$$

sending the point in $[0]$ to $0 \in [1]$ maps to

$$\begin{aligned} \text{LCut}(a) : (\langle 2 \rangle, \{2\}) &\rightarrow (\langle 1 \rangle, \{1\}) \\ 1 &\mapsto 1 \\ 2 &\mapsto 1. \end{aligned}$$

Interpreting $(\langle 2 \rangle, \{2\})$ as $(\mathfrak{a}, \mathfrak{m})$ and $(\langle 1 \rangle, \{1\})$ as \mathfrak{m} , this map corresponds to the unique element

$$\phi \in \text{Mul}_{\mathcal{LM}}(\{\mathfrak{a}, \mathfrak{m}\}, \mathfrak{m}).$$

Similarly, the map

$$b : ([0], 0) \rightarrow ([1], 0)$$

sending the point in $[0]$ to $1 \in [1]$ maps to

$$\begin{aligned} \text{LCut}(b) : (\langle 2 \rangle, \{2\}) &\rightarrow (\langle 1 \rangle, \{1\}) \\ 1 &\mapsto * \\ 2 &\mapsto 1. \end{aligned}$$

This map corresponds to the unique element

$$\psi \in \text{Mul}_{\mathcal{LM}}(\{\mathfrak{m}\}, \mathfrak{m}).$$

Therefore, to give an enriched morphism of \mathcal{M} is equivalent to giving a diagram

$$M \xleftarrow{\alpha} X \xrightarrow{\beta} N$$

in \mathcal{C}^\otimes such that

- (1) $q(\alpha) = \text{LCut}(a)$,
- (2) $q(\beta) = \text{LCut}(b)$, and
- (3) β is inert, i.e. q -coCartesian.

Unpacking this, M and N are objects in \mathcal{M} , and $X = (C, M')$ is an object in $\mathcal{C}_{(\mathfrak{a}, \mathfrak{m})}^\otimes \simeq \mathcal{C}_{\mathfrak{a}} \times \mathcal{C}_{\mathfrak{m}}^1$, while $\alpha : (C, M') \rightarrow M$ and $\beta : (C, M') \rightarrow N$ are morphisms in \mathcal{C}^\otimes lifting ϕ and ψ respectively. Since q is coCartesian, there is a q -coCartesian lift for ϕ and $X = (C, M')$, namely the map $(C, M') \rightarrow C \otimes M'$. Hence, the data of α is equivalent to a map $C \otimes M' \rightarrow M$ in \mathcal{M} . Similarly, there is q -coCartesian lift for ψ and $X = (C, M')$, namely the map $(C, M') \rightarrow M'$. Hence the data of β is equivalent to a map $M' \rightarrow N$ in \mathcal{M} , and since β is supposed to be q -coCartesian as well, this map has to be an equivalence. Hence, the ∞ -category $\mathcal{C}_{\mathfrak{a}}[M]$ as defined in [Lur17, Definition 4.7.1.1] is equivalent to the ∞ -category with objects given by pairs $(C \in \mathcal{C}_{\mathfrak{a}}, \eta : C \otimes M \rightarrow M)$, which is better known as

$$\mathcal{C}_{\mathfrak{a}} \times_{\mathcal{M}} \mathcal{M}_{/M}.$$

□

¹This holds because of [Lur17, Proposition 2.1.2.12]

REFERENCES

- [ABA00] Didier Arnal and Nabiha Ben Amar. Kontsevich’s wheels and invariant polynomial functions on the dual of Lie algebras. *Lett. Math. Phys.*, 52(4):291–300, 2000.
- [AT12] Anton Alekseev and Charles Torossian. The Kashiwara-Vergne conjecture and Drinfeld’s associators. *Ann. of Math. (2)*, 175(2):415–463, 2012.
- [BM09] Clemens Berger and Ieke Moerdijk. On the derived category of an algebra over an operad. *Georgian Math. J.*, 16(1):13–28, 2009.
- [BN98] Dror Bar-Natan. On associators and the Grothendieck-Teichmüller group. I. *Selecta Math. (N.S.)*, 4(2):183–212, 1998.
- [BR23] Christopher Brav and Nick Rozenblyum. The cyclic deligne conjecture and calabi-yau structures, 2023.
- [CVdB10] Damien Calaque and Michel Van den Bergh. Hochschild cohomology and Atiyah classes. *Adv. Math.*, 224(5):1839–1889, 2010.
- [DRW15] V. A. Dolgushev, C. L. Rogers, and T. H. Willwacher. Kontsevich’s graph complex, GRT, and the deformation complex of the sheaf of polyvector fields. *Ann. of Math. (2)*, 182(3):855–943, 2015.
- [Fra13] John Francis. The tangent complex and hochschild cohomology of \mathcal{E}_n -rings. *Compositio Mathematica*, 149(3):430–480, 2013.
- [Fre09] Benoit Fresse. *Modules over operads and functors*, volume 1967 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009.
- [Ger63] Murray Gerstenhaber. The cohomology structure of an associative ring. *Ann. of Math. (2)*, 78:267–288, 1963.
- [Hin05] Vladimir Hinich. Deformations of sheaves of algebras. *Adv. Math.*, 195(1):102–164, 2005.
- [Hin15] Vladimir Hinich. Rectification of algebras and modules. *Doc. Math.*, 20:879–926, 2015.
- [HKV06] Po Hu, Igor Kriz, and Alexander A. Voronov. On Kontsevich’s Hochschild cohomology conjecture. *Compos. Math.*, 142(1):143–168, 2006.
- [HV02] Vladimir Hinich and Arkady Vaintrob. Cyclic operads and algebra of chord diagrams. *Selecta Math. (N.S.)*, 8(2):237–282, 2002.
- [Iwa20] Isamu Iwanari. Differential calculus of Hochschild pairs for infinity-categories. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 16:Paper No. 097, 57, 2020.
- [Kon03] Maxim Kontsevich. Deformation quantization of Poisson manifolds. *Lett. Math. Phys.*, 66(3):157–216, 2003.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [Lur17] J. Lurie. Higher algebra, 2017. <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [Mar09] Nikita Markarian. The Atiyah class, Hochschild cohomology and the Riemann-Roch theorem. *J. Lond. Math. Soc. (2)*, 79(1):129–143, 2009.
- [MS02] James E. McClure and Jeffrey H. Smith. A solution of Deligne’s Hochschild cohomology conjecture. In *Recent progress in homotopy theory (Baltimore, MD, 2000)*, volume 293 of *Contemp. Math.*, pages 153–193. Amer. Math. Soc., Providence, RI, 2002.
- [PS18a] Dmitri Pavlov and Jakob Scholbach. Admissibility and rectification of colored symmetric operads. *J. Topol.*, 11(3):559–601, 2018.
- [PS18b] Dmitri Pavlov and Jakob Scholbach. Homotopy theory of symmetric powers. *Homology Homotopy Appl.*, 20(1):359–397, 2018.
- [Ram08] Ajay C. Ramadoss. The big Chern classes and the Chern character. *Internat. J. Math.*, 19(6):699–746, 2008.
- [RW10] Justin Roberts and Simon Willerton. On the Rozansky-Witten weight systems. *Algebr. Geom. Topol.*, 10(3):1455–1519, 2010.
- [Tam98] D. Tamarkin. Another proof of M. Kontsevich formality theorem, 1998.
- [Tam02] Dimitri Tamarkin. Action of the Grothendieck-Teichmüller group on the operad of Gerstenhaber algebras, 2002.
- [Tam03] Dmitry E. Tamarkin. Formality of chain operad of little discs. *Lett. Math. Phys.*, 66(1-2):65–72, 2003.
- [Vor00] Alexander A. Voronov. Homotopy Gerstenhaber algebras. In *Conférence Moshé Flato 1999, Vol. II (Dijon)*, volume 22 of *Math. Phys. Stud.*, pages 307–331. Kluwer Acad. Publ., Dordrecht, 2000.
- [Whi22] David White. Monoidal Bousfield localizations and algebras over operads. In *Equivariant topology and derived algebra*, volume 474 of *London Math. Soc. Lecture Note Ser.*, pages 180–240. Cambridge Univ. Press, Cambridge, 2022.

- [Yek02] Amnon Yekutieli. The continuous Hochschild cochain complex of a scheme. *Canad. J. Math.*, 54(6):1319–1337, 2002.

DEPARTMENT OF MATHEMATICS & STATISTICS, UNIVERSITY OF NEVADA, RENO.

Email address: `sonjaf@unr.edu`