PURAMINO
DHA & ARA
Hypoallergenic Infant Formula
amino acid-based powder with iron
0-24 months
NON-GMO
For oral and tube feeding

DHA ARA

HYPOALLERGENIC

MCT OIL

Explore our patient savings programs

Request Samples

View Product Resources

PurAmino®

Print

Hypoallergenic amino acid-based formula for infants with severe cow's milk protein allergy and multiple food allergies or elemental diet needs

Indication

PurAmino is a 20 cal/fl oz, iron-fortified, hypoallergenic, amino acid-based infant formula for the dietary management of infants with severe cow's milk protein allergy, not effectively managed by an extensively hydrolyzed formula. PurAmino is also indicated for the dietary management of infants with multiple food protein allergies. It may also be suitable for conditions requiring an elemental diet, such as fat malabsorption, multiple food protein allergies, food protein-induced enterocolitis syndrome (FPIES), eosinophilic esophagitis (EoE),* and short bowel syndrome. Appropriate for both oral and tube feeding.

*EoE is the most common type of eosinophilic gastrointestinal disorders (EGID).

Product Benefits

PurAmino provides relief and supports growth so infants can grow and thrive

- Demonstrated to support healthy weight gain and improvement in allergic manifestations in infants not responding to extensively hydrolyzed formulas
- Hypoallergenic, 100% free amino acids as nitrogen source:
  - 100% of infants allergic to cow's milk had no adverse reaction when fed PurAmino[1]
  - 2.8 g protein equivalent/100 cal
  - 33% of fat as medium chain triglycerides (MCT) oil to help facilitate fat absorption
- Significant improvement in GI and skin symptoms by the next physician visit, when fed PurAmino™[2,†] (see study below). **Current guidelines recommend extensively hydrolyzed formula (eHF) as the first choice for CMPA, with amino acid formulas (AAF) for more severe cases or those not responding to eHF.**
- Significantly reduced incidence and severity[‡] of allergic symptoms at 12 weeks[3] (see study below)
- 87% catch-up weight gain in 12 weeks[8]

Additional Product Features

- Nutritionally complete:
  - Can be sole source of nutrition up to age 6 months
  - Can be major source of nutrition through 24 months
  - Dietary reference intakes include a recommendation for fiber for >12 months of age
- Lactose-free
- Kosher Dairy (OU-D), Halal
- Clinically proven to help infants stay on their growth curve[1] (see study below)
- Expert-recommended DHA at 0.32% of total fatty acids, the same amount as the worldwide breastmilk average[9,10¶]

[†] Based on HCP reported user experience in infants with suspected CMPA.
[‡] 15.8 GI Symptom Score improvement and reduced scoring atopic dermatitis (SCORAD).
[¶] The World Health Organization recommended amount of DHA is 0.2%-0.36% of total fatty acids. Average DHA in worldwide breast milk is 0.32% ±0.22% (mean ± standard deviation of total fatty acids) based on an analysis of 65 studies of 2474 women.

Significant improvement in GI and skin symptoms by the next physician visit[2]



STUDY DESIGN (Wilsey et al. 2023 ↗): Prospective cohort analysis of de-identified patient data collected by HCPs who analyzed the clinical profile of infants displaying symptoms consistent with suspected CMPA or diagnosed CMPA. Inclusion criteria included patients less than 6 months of age prescribed an amino acid formula at the enrollment visit (Visit 1) whose HCPs provided symptom data at Visit 1 and the next visit (Visit 2) at 3 to 6 weeks (n = 104). Study suggests that AAF may be an effective option for CMPA management for infants 6 months of age and under with suspected CMPA, often by the next physician visit.

The percent improvement is calculated by determining the percent difference in the incidence of total presented symptoms (at different severity levels: low, moderate, and severe) between Visit 1 and Visit 2.

*Number of patients who presented with the reported issue at the first visit and were assessed at the second visit.

PurAmino™ promotes catch-up weight gain

In babies with faltering growth,[†] mean growth percentiles improved from 5th to 14th.



STUDY DESIGN (Vanderhoof et al. 2016 ↗): Observational, prospective, multicenter study of infants (n = 30; 1-12 months of age [mean age 6.6 months]) with a birth weight of ≥1500 g, suspected CMPA (experiencing at least one persistent CMPA symptom), and a history of weight loss (≥0.5 z score) who were not responding to an extensively hydrolyzed formula. Eligible infants received an amino acid-based formula for 12 weeks. Z score improved from -1.6 (5th percentile) to -1.1 (14th percentile) ($P < 0.001$).

Chart adapted from WHO growth charts 0 to 5 years, boys and girls combined.[2]

[§] Defined by a GI Symptom Score >16.

Savings and Assistance Programs

Long-Term Usage

PurAmino is designed to provide a sole source of nutrition for infants up to age 6 months and provide a major source of nutrition through 24 months of age, when indicated. Normally, in feeding infants, gradual introduction of solid foods after 4-6 months of age is an important developmental as well as nutritional step. In cases of severe and multiple food allergies or intolerances, PurAmino is sometimes continued in the diet of children.

Extended use of medical foods like PurAmino as a sole source of nutrition should be used with medical supervision.

PurAmino Recipe Book

To support the nutritional status of patients identified as having pediatric food allergies, Registered Dietitian and pediatric allergy expert Raquel Durban MS, RD, LD/N developed the recipes and nutrient profiles as a paid consultant to Mead Johnson Nutrition.

Download Recipes ↗

Ingredients

Ingredients: Corn syrup solids (46%), amino acids (potassium aspartate, L-leucine, L-lysine hydrochloride, L-proline, L-alanine, L-valine, L-isoleucine, L-serine, L-threonine, L-glutamine, L-tyrosine, L-aspartic acid, L-arginine, L-phenylalanine, L-histidine, L-cystine, glycine, L-tryptophan, L-methionine) (18%), high oleic sunflower oil (9%), medium chain triglycerides (MCT) oil (9%), soy oil (8%), modified tapioca starch (4%), and less than 2%: Mortierella alpina oil,[||] Schizochytrium sp. oil,[¶] choline chloride, inositol, ascorbic acid, niacinamide, calcium pantothenate, riboflavin, thiamin hydrochloride, vitamin B_6 hydrochloride, vitamin D_3, folic acid, vitamin K_1, biotin, taurine, vitamin E acetate, L-carnitine, vitamin A palmitate, vitamin B_{12}, calcium phosphate, potassium citrate, sodium citrate, magnesium phosphate, ferrous sulfate, zinc sulfate, cupric sulfate, manganese sulfate, sodium chloride, sodium iodide, sodium selenite.

[||] A source of arachidonic acid (ARA).
[¶] A source of docosahexaenoic acid (DHA).

POTENTIAL ALLERGENS

PurAmino™ contains soy oil. PurAmino™ is hypoallergenic.

WARNING: Not for parenteral (I.V.) use.

Ingredient Transparency

Explore the function behind our ingredients.

Learn More ↗

Nutrients

Normal Dilution	Per 100 kcal (5 fl oz)	Per 100 mL	Per 100 grams Powder (500 kcal)
Protein equivalent, g	2.8	1.9	14
Fat, g	5.3	3.6	27
Linoleic acid, mg	860	580	4300
Carbohydrate, g	10.6	7.2	53
Water, g	133	90	1.76
Vitamins/Other Nutrients			
Vitamin A, IU	300	200	1500
Vitamin D, IU	60	41	300
Vitamin E, IU	2	1.36	10
Vitamin K, mcg	9	6.1	45
Thiamin (Vitamin B_1), mcg	80	54	400
Riboflavin (Vitamin B_2), mcg	90	61	450
Vitamin B_6, mcg	60	41	300
Vitamin B_{12}, mcg	0.3	0.2	1.5
Niacin, mcg	1000	680	5000
Folic acid (Folacin), mcg	16	10.9	80
Pantothenic acid, mcg	500	340	2500
Biotin, mcg	3	2	15
Vitamin C (Ascorbic acid), mg	12	8.2	60
Choline, mg	24	16.3	120
Inositol, mg	24	16.3	120
Minerals			
Calcium, mg	116	79	580
Phosphorus, mg	64	44	320
Magnesium, mg	8	5.4	40
Iron, mg	1.8	1.22	9
Zinc, mg	1	0.68	5
Manganese, mcg	25	17	125
Copper, mcg	75	51	380
Iodine, mcg	15	10.2	75
Selenium, mcg	2.8	1.9	14
Sodium, mg	47	32	240
Potassium, mg	110	75	550
Chloride, mg	86	58	430

Product nutrient values and ingredients are subject to change. Please see product label for current information.

Nutrient Density	20 kcal/fl oz
Protein (% Calories)	11
Fat (% Calories)	47
Carbohydrate (% Calories)	42
Potential Renal Solute Load (mOsm/100 kcal)[§]	25
Potential Renal Solute Load (mOsm/100 mL)[§]	17.2
Osmolality (mOsm/kg water)	350
Osmolarity (mOsm/L formula)	320
Lactose-free	Yes

AMINO ACID COMPOSITION

Essential and Nonessential Amino Acids

Download tables ↗

Dilutions and Preparation

DILUTION INFORMATION FOR PURAMINO INFANT

Mixing instruction by the scoop

Download tables ↗

Mixing instruction by the quart

Download tables ↗

POWDER

Refer to the product label for the most accurate information.

A baby's health depends on carefully following the instructions below. Use only as directed by a medical professional. Proper hygiene, preparation, dilution, use, and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should **NOT** be fed to premature infants or infants who might have immune problems unless directed and supervised by a doctor.

Discuss with parents which formula is appropriate for the baby.

Discuss with parents whether they need to use cooled, boiled water for mixing and whether they need to boil clean utensils, bottles and nipples in water before use.

Wash hands thoroughly with soap and water before preparing formula.Pour the desired amount of water into the bottle. Add powder. Cap bottle and **SHAKE WELL**.

Use the charts below for correct amounts of water and powder. Use scoop in can to measure powder. Store **DRY** scoop in its original can.

WARNING: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 35-40 °F (2-4 °C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within 1 hour or discard.

For every 1 fl oz of water, add 1 unpacked level scoop of powder (4.5 g). Each scoop adds about 0.1 fl oz to the amount of prepared formula. For example, adding 3 unpacked level scoops of powder to 3 fl oz of water will make about 3.3 fl oz of formula.

	Water	Powder
For every	1 fl oz	add 1 unpacked level scoop (4.5 g)

Per 1 Scoop*

Calories Desired (kcal)		Add 1 scoop powder to water	Formula Yield
per fl oz[†]	per mL	(fl oz)	Formula Yield
20	0.68	1.0	1.1
22	0.74	0.9	1.0
24	0.81	0.8	0.9
26	0.88	0.7	0.9
27	0.91	0.7	0.8
28	0.95	0.7	0.8
30	1.01	0.6	0.8

PurAmino formula should be measured with unpacked, level scoops.

*One scoop of powder provides 23 calories.
[†] Fluid ounce measures in the above table are rounded to the nearest 0.1 fl oz.

Powder Storage

Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

USE BY DATE ON BOTTOM OF CAN.

Product Forms and Ordering Information

PURAMINO IS AVAILABLE IN POWDER

Coding systems and reimbursement allowable rates vary by payer:

- Medicare Part B uses HCPCS (Healthcare Common Procedure Coding System) to group products
- Medicaid systems vary by state; some use HCPCS while others use NDC format code or systems of their own—contact your state provider for more information
- Private health insurance and managed care companies may use HCPCS, NDC format code or their own system—contact your provider for more information

PurAmino, Powder

Item #:
179101

Description:
Powder

Unit Size:
14.1 oz can

Calories/Unit:
1960

Product Yield/Unit (fl oz):
98

Case:
4 cans per case

HCPCS Code:
B4161

Reimbursement Code (NDC format):
00087-5104-80

Related Products

PURAMINO Jr
AGES 1 AND UP
Medical Food
Amino Acid-Based Nutrition Powder
VANILLA
DHA
NON-GMO

PurAmino™ Jr Vanilla

Nutritionally complete*, hypoallergenic amino acid-based medical food to help support growth and development of children 1 year and up

More Information

PURAMINO Jr
AGES 1 AND UP
Medical Food
Amino Acid-Based Nutrition Powder
UNFLAVORED
DHA
NON-GMO

PurAmino™ Jr

Nutritionally complete,* hypoallergenic amino acid-based medical food to help support growth and development of children 1 year and up

More Information

Patient Savings & Support Programs

Helping Hands™ provides support for families:

Helping Hands Sample Assistance

FREE one-time shipment of PurAmino product, per child, per household

Learn more about Helping Hands Sample Assistance here.

Helping Hands Reimbursement Support

PurAmino may be eligible for Insurance coverage through Medicald, Medi-Cal, TRICARE, and most commercial health Insurance plans when prescribed. We know that navigating insurance coverage is not always easy. Through Helping Hands" Reimbursement Support, our team of dedicated reimbursement experts can help guide parents through the process from start to finish. From benefits verification to prior authorization assistance, we handle all the details so patients can get quick access to Mead Johnson Nutrition products when prescribed.

While we can't guarantee insurance reimbursement, our dedicated staff will help parents navigate the reimbursement process.

Learn more about helping Hands Reimbursement Support here.

PurAmino is WIC* eligible in all 50 states:

Learn more about WIC here.*

*WIC is a registered trademark of the United States Department of Agriculture (USDA) for the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC). No endorsement of any brand or product by the USDA is implied or intended.

Resources

Clinical Study

Hypoallergenicity, growth, and tolerance with amino acid-based formula and DHA.

See Study ↗

PurAmino Letter of Medical Necessity

Letters of medical necessity are available to help make accessing the Mead Johnson formula you recommend easier for your patients. Use this form for patients prescribed PurAmino or PurAmino™ Jr.

See Study ↗

Contact Mead Johnson

Contact a Mead Johnson representative to request information.

Contact Us

PurAmino Infant formula is WIC®*-eligible in 46 states

Access your state's medical documentation form to help your WIC® infants receive the formula they need.

Read More

Empower families to manage food allergies

Understand the key facts, considerations, and hypoallergenic solutions for cow's milk allergy.

Read More

*WIC® is a registered service mark of the US Department of Agriculture for USDA's Special Supplemental Nutrition Program for Women, Infants, and Children. No endorsement of any brand or product by the USDA is implied or intended. PurAmino Infant is WIC®-eligible in 46 states. It is not approved in Alaska, New York, Washington, or Wisconsin.

References

1. Burks W, Jones SM, Berseth CL, Harris C, Sampson HA, Scalabrin SMF. Hypoallergenicity and effects on growth and tolerance of a new amino acid-based formula with docosahexaenoic acid and arachidonic acid. *J Pediatr.* 2008;153(2):266-271.
2. Vanderhoof J, Moore N, de Boissieu D. Evaluation of an amino acid-based formula in infants not responding to extensively hydrolyzed protein formula. *J Pediatr Gastroenterol Nutr.* 2016;63(5):531-533.
3. Birch EE, Hoffman DR, Castañeda YS, Fawcett SL, Birch DG, Uauy RD. A randomized controlled trial of long-chain polyunsaturated fatty acid supplementation of formula in term infants after weaning at 6 wk of age. *Am J Clin Nutr.* 2002;75(3):570-580.
4. Hoffman DR, Birch EE, Castañeda YS, et al. Maturation of visual and mental function in 18-month old infants receiving dietary long-chain polyunsaturated fatty acids [abstract]. *FASEB J.* 2003;17:A727-A728. Abstract 445.1.
5. Hoffman DR, Birch EE, Castañeda YS, et al. Visual function in breast-fed term infants weaned to formula with or without long-chain polyunsaturates at 4 to 6 months: a randomized clinical trial. *J Pediatr.* 2003;142(6):669-677.
6. Hoffman DR, Birch EE, Birch DG, et al. Impact of early dietary intake and blood lipid composition of long-chain polyunsaturated fatty acids on later visual development. *J Pediatr Gastroenterol Nutr.* 2000;31(5):540-553.
7. Birch EE, Castañeda YS, Wheaton DH, et al. Visual maturation of term infants fed long-chain polyunsaturated fatty acid-supplemented or control formula for 12 mo. *Am J Clin Nutr.* 2005;81(4):871-879.
8. Morale SE, Hoffman DR, Castañeda YS, Wheaton DH, Burns RA, Birch EE. Duration of long-chain polyunsaturated fatty acids availability in the diet and visual acuity. *Early Hum Dev.* 2005;81(2):197-203.

Intended for U.S. Healthcare Professionals Only. ©2004, 2026 Mead Johnson & Company, LLC. All Rights Reserved.