

```

 $\mathbf{x}^T \mathbf{y}$  ( $\mathbf{x}^H \mathbf{y}$ ): dot = psb_gedot(x,y,desc_a,info)
 $\mathbf{y} \leftarrow \alpha \mathbf{x} + \beta \mathbf{y}$ : call psb_geaxpby(alpha,x,beta,y,desc_a,info)
 $\max_i |x_i|$ : amax = psb_geamax(x,desc_a,info)
 $\sum_i |x_i|$ : asum = psb_geasum(x,desc_a,info)
 $\|\mathbf{x}\|_2$ : nrm2 = psb_genrm2(x,desc_a,info)
 $\|\mathbf{A}\|_\infty$  nrmi = psb_spnrm1(A,desc_a,info)
 $\mathbf{y} \leftarrow \alpha \mathbf{A}^T \mathbf{x} + \beta \mathbf{y}$ : call psb_spmv(alpha,A,x,beta,y,desc_a,info,trans='T')
 $\mathbf{y} \leftarrow \alpha \mathbf{D} \mathbf{T}^T \mathbf{x} + \beta \mathbf{y}$ : call psb_spsv(alpha,T,x,beta,y,desc_a,info,trans='T',[,unitd])

```

Note: T is a triangular **AND** block diagonal matrix (i.e.: Block-Jacobi or Hybrid GS type preconditioners)