


Tools for Characterizing a Population of Massive Black Hole Binaries (Imaging gravitational wave driven binaries?)

Daniel J. D'Orazio


NASA Einstein Postdoctoral Fellow, Harvard University

Testing Gravity 2017
Saturday January 28, 2017

The Gravitational Wave Spectrum




Massive Black Hole Binaries (MBHBs)



- * Primary (known) sources of low frequency gravitational waves
- * Can teach us about the mutual evolution of galaxies and MBHs
- * Characterization of MBHB population by EM signatures can inform/compliment characterization by **gravitational wave background...**
- * How do we find them? They don't (necessarily) merge in vacuum!




$$M_{\text{bin}} \sim 10^6 \rightarrow 10^{10} M_\odot$$

Quasar Anatomy: where MBHBs hang out





Electromagnetic MBHB evidence/searches

Separation



Grav waves




\sim kpc

\sim pc


\sim sub-pc

Jet morphology




NGC 326


Broad line monitoring




The Event Horizon Telescope: Milky Way Black Hole



Doeleman+2009, Fish+2012, Lu+2014



$$\theta_{EH} \sim \frac{R_S}{8\text{kpc}} \sim 10\mu\text{as}$$





Diffraction limited resolution

$$\theta \sim \frac{\lambda}{D} = \frac{1\text{mm}}{10^4\text{km}} \rightarrow 20\mu\text{as}$$

*Can push astrometric precision to
 $\sim 1\mu\text{as}$

(Broderick, Loeb, Reid 2011)

The Event Horizon Telescope: MBHBs



Could resolve binaries in GW driven regime
(work with Avi Loeb)

How many binary's could we Image with the EHT?

*Assume a fraction of all AGN are triggered by binaries:

Requirements:

- * Bright in (sub)-mm
(preferably both BHs)

- * Minimum binary separation
set by EHT resolution
- * Maximum binary separation
set by observable orbital
period (<10 years)

Ingredients:

$$\frac{N_{\text{EHT}}(z)}{\Delta\Omega\Delta z} \approx \frac{dV}{dzd\Omega} \int_{L_{mm}^{\min}}^{\infty} \frac{dN}{dL_{mm}dV} \mathcal{F}(L_{mm}, z) dL_{mm}$$

sub-mm luminosity function:

From observationally constructed AGN
radio luminosity functions


Binary Probability:

Calculated from residence
time at required separations


$$t_{\text{res}} = \frac{\dot{a}}{a}$$

Assume:
Gas + Gravitational waves

How many binary's could we Image with the EHT?



Demographics



- * Orbital velocities could allow large Doppler-boost modulations
- * In PTA band, but strain below current GWB limits

Implications

- * Aid in narrowing down mechanisms which drive MBHBs to merger -> constrain GWB
- * Assume cosmology: precise measurement of the binary mass

$$GM = \left(\frac{\Omega}{1+z} \right)^2 (\theta_{\max} D_A(z))^3$$

- * Measure binary mass: measurement of Hubble constant

Summary

- * Characterization of MBHB population with EM signatures can constrain expected GWB as well as astrophysics (accretion, mutual growth of BHs and galaxies)
- * The EHT could resolve the orbital separations of MBHBs on short enough periods to watch entire orbits
- * Beyond imaging such elusive objects, this could allow precise binary mass measurements, a measurement of the Hubble constant, test of the speed of gravitational waves, or...