

Modeling and Constraining the Cluster Mass Function to Test Gravity at Large Scales

David Rapetti

University of Colorado Boulder / NASA Ames

Work presented here in collaboration with:

Matteo Cataneo (ROE), Fabian Schmidt (MPA), Baojiu Li (Durham), Lucas Lombriser (ROE), Adam Mantz (KIPAC), Steve Allen (KIPAC), Douglas Applegate (KICP), Patrick Kelly (Berkeley), Anja von der Linden (Stony Brook), Glenn Morris (KIPAC)

Constraining alternative theories of gravity

1. **Strength** from cluster observations: constraining **gravity at large scales**, either using **concrete models** or **general parameterizations**.
2. Worked example: viable **$f(R)$ gravity** models with the **chameleon screening mechanism**.
3. For $f(R)$, **current prospects** for improvement are primarily based on **analyses including less massive and low- z objects**.

Hu-Sawicki $f(R)$ gravity model (departing from GR)

$$f(R) = -2\Lambda - \frac{f_{R0}}{n} \frac{\bar{R}_0^{n+1}}{R^n}$$

Approximated Hu-Sawicki model in the high curvature regime. In the limit $|f_{R0}| \ll 10^{-2}$ closely mimics the Λ CDM expansion history

$$\lambda_{C0} \approx 29.9 \sqrt{\frac{|f_{R0}|}{10^{-4}} \frac{n+1}{4-3\Omega_m}} h^{-1} \text{Mpc}$$

Compton wavelength; scales below this present modified gravity until GR is recovered when the fifth force is screened by non-linear effects

$$g(a, k) \equiv -\frac{1}{3} \frac{k^2}{k^2 + m_{f_R}^2 a^2}$$

$$m_{f_R}^{-2} = \lambda_C^2 = 3f_{RR} \quad f_R \equiv \frac{\partial f}{\partial R}$$

Linear growth is different than GR+ Λ CDM

Halo mass function modeling

$$n_{\Delta_v} \equiv \frac{dn}{d \ln M_v} = \frac{\bar{\rho}_m}{M_v} \frac{d \ln \nu}{d \ln M_v} \nu f(\nu) \quad \text{Sheth-Tormen mass function}$$

$$\nu f(\nu) = A \sqrt{\frac{2}{\pi}} a \nu^2 [1 + (a \nu^2)^{-p}] \exp [-a \nu^2 / 2] \quad \nu = \delta_c / \sigma(M_v) \quad \text{Peak height}$$

$$\delta_c(\Omega_m, z) = \mathcal{A} \left(1 - \mathcal{B} \log_{10} \left[1 + \frac{\Omega_m^{-1} - 1}{(1+z)^3} \right] \right) \quad \text{Density threshold fitting formula}$$


$$n_{\Delta} = \left(\frac{n_{\Delta}^{f(R)}}{n_{\Delta}^{\text{GR}}} \right) \Big|_{\text{ST}} n_{\Delta} \Big|_{\text{Tinker}}$$

$$\mathcal{A} = 1.6865 \text{ and } \mathcal{B} = 0.0123 \text{ for GR}$$

$$\mathcal{A} = 1.7063 \text{ and } \mathcal{B} = 0.0136 \text{ for } f(R)$$

$$n_{\text{ST}}^{f(R)} / n_{\text{ST}}^{\text{GR}} \quad \text{We set this equal to 1 when becomes } < 1$$


Halo mass function modeling



- Using N-body simulations including the **Chameleon screening mechanism** (Schmidt et al 2009).

- Our **modeling** is based on the **bottom line** of the blue shaded area, which as shown here is **conservative**.

Halo mass function modeling




- Using N-body simulations including the **Chameleon screening mechanism** (Schmidt et al 2009).

- Our **modeling** is based on the **bottom line** of the blue shaded area, which as shown here is **conservative**.

HS n=1 f(R) growth + flat Λ CDM

Cataneo et al 15 (PRD 2015, 92, 044009)



$$|f_{R0}| < 1.62 \times 10^{-5} (All)$$

Clusters: XLF: BCS+REFLEX+MACS (z<0.5) 224 survey with 94 X-ray follow-up (Mantz et al 2015) + cluster f_{gas} (Mantz et al 2014)

CMB (Planck collaboration 2014; SPT, Story et al 2013; ACT, Das et al 2014) + SNIa (Union 2.1, Suzuki et al 2012) + BAO (6dF, Beutler et al 2011; SDSS, Padmanabhan et al 2012, Anderson et al 2014; WiggleZ, Blake et al 2011)


Gold: all data sets combined

For General Relativity |f_{R0}|=0

Correlation between σ₈ and |f_{R0}|

$$\rho = 0.73 (All) \quad \rho = 0.90 (CMB)$$

Halo mass function modeling




- For **Chameleon gravity models** using medium-resolution N-body simulations for $f(R)$ as limiting case (Lombriser et al 2014).

- **Modeling** the effect of the **Chameleon screening mechanism** to the mass function to obtain a more accurate mass function and **stronger constraints** (Cataneo et al 2017, in prep.).

- The **small-field regime** is **potentially** constrainable by **lower-mass cluster data** (with a **$\sim 5\%$ mass calibration precision**). The curves depend on the **environmental density**.

Chameleon screening refinement

Cataneo et al 16, (JCAP 2016, 12, 024)




- Modeling the effect of the Chameleon screening mechanism on the mass function using high-resolution N-body simulations to obtain a more accurate mass function (Cataneo, Rapetti, Lombriser, Li, 2016)
- Spherical collapse thresholds at $z=0$ ($\Omega_m=0.281$). Lombriser et al 2014 calculations at the peak of the environmental density distribution in blue and our corrected/calibrated δ_c in red to account for self-screening and environmental screening mechanisms.

Chameleon screening refinement

$$\delta_c^{\text{eff}} = \epsilon(M \mid M_{\text{th}}^{(1)}, \mu, M_{\text{th}}^{(2)}, \nu, \alpha) \times \delta_c(\delta_{\text{env}}^{\text{peak}})$$


$$\epsilon = \frac{1 + (M/M_{\text{th}}^{(1)})^\eta (\delta_c^\Lambda / \delta_c^{f(R)})^\chi + (M/M_{\text{th}}^{(2)})^\vartheta (\delta_c^{f(R)} / \delta_c^\Lambda)}{1 + (M/M_{\text{th}}^{(1)})^\eta + (M/M_{\text{th}}^{(2)})^\vartheta}$$



GR/LCDM environmental density probability distribution (Lam & Sheth 08); $z=0$, 0.2 , 0.5 ; $\Omega_m=0.281$

New halo mass function

Cataneo et al 16




- High-resolution: $L_{\text{box}}=1024 \text{ Mpc}/h$; $N_{\text{particles}}=1024$; force resolution = $15.3 \text{ kpc}/h$; $N_{\text{realizations}} = 1$; $z = 0$ (blue), 0.1, 0.2 (red), 0.3, 0.4, 0.5 (green); $n_s=0.971$, $\Omega_m=0.281$, $H_0=69.7$, $T_{\text{CMB}}=2.7255$, $Y_{\text{He}}=0.24$, $N_{\nu}=3$, $\sigma_8=0.82$ (flat Λ CDM background). Current fits:
 5% for $10^{13.5} \leq M_{300m}(M_{\odot}/h)^{-1} \leq 10^{15.5}$
 $10^{-6} \leq |f_{R0}| \leq 10^{-4}$ and $0 \leq z \leq 0.5$
- Rockstar halo finder with spherical overdensity masses with average density equals $300\rho_m$; simulations divided in octants; uncertainties on the HMF $f(R)/\text{GR}$ ratios propagated with the jackknife method with halos in mass bins of $\Delta\log_{10}(M)=0.15$. (We only keep mass bins with $N_{\text{halos}}>20$.)

Jan 26, 2017

Testing Gravity, SFU, Vancouver

New halo mass function

Cataneo et al 16




- Low-resolution: $L_{\text{box}}=1.5$ Gpc/h; $N_{\text{particles}}=1024$; force resolution = 22.9 kpc/h; $N_{\text{realizations}} = 6$; $z = 0$ (blue), 0.11, 0.25 (red), 0.43 (green); $n_s=0.958$, $\Omega_m=0.24$, $H_0=73$, $T_{\text{CMB}}=2.7255$, $Y_{\text{He}}=0.24$, $N_{\nu}=3$, $\sigma_8=0.796$ (flat Λ CDM background).
- **Rockstar halo finder** with spherical overdensity masses with average density equals $300\rho_m$; uncertainties on the HMF $f(R)/GR$ ratios propagated with **the jackknife method using the six realizations** with halos in mass bins of $\Delta\log_{10}(M)=0.15$. (We only keep mass bins with $N_{\text{halos}}>20$.)

Jan 26, 2017

Testing Gravity, SFU, Vancouver

New halo mass function



$$\mathcal{R}(M, z, \bar{f}_{R0}, \boldsymbol{\theta}) = \frac{n_{\ln M}^{f(R)}}{n_{\ln M}^{\text{GR}}}$$


- Dashed lines show the effect of neglecting the correction,
- which is only weakly dependent on **standard cosmological parameters**
- and potentially insensitive to **baryonic physics and massive neutrinos**

CATANEO ET AL 2016

Jan 26, 2017

Testing Gravity, SFU, Vancouver


Comparison of current and prospective constraints



$$|\bar{f}_{R0}| \lesssim 5 \times 10^{-6}$$

Gray band, 7% mass calibration from WtG with the corresponding lower mass limit; red line, current 2-sigma level from the WtG constraints using the previous mass function; blue line, projected constraints for the next WtG constraints with the new $f(R)$ HMF (\sim a factor of 2 improvement).

Prospective constraints for a lower mass/z survey




$\log_{10} (M_{300} \text{ h}/M_{\odot})$

Cataneo et al 16

$$|\bar{f}_{R0}| \lesssim 10^{-6}$$

Gray band, 5% mass calibration from DES with the corresponding lower mass limit; blue line, projected constraints for DES constraints with the new $f(R)$ HMF (~an order of magnitude improvement).

Comparison of current and prospective constraints



Adapted from Joyce,
Jain, Khoury & Trodden
2014 (review)

Cataneo et al 15
 $\lambda_c < \sim 10 \text{ Mpc}$ $|f_{R0}| < \sim 10^{-5}$

Cataneo et al 17 (in pr.),
a factor of ~ 2 better

DES forecast
 $\lambda_c < \sim 3 \text{ Mpc}$ $|f_{R0}| < \sim 10^{-6}$
(Cataneo et al 2016)