# Geographic Variation in Childhood Obesity

The effect of a healthier school environment

Samuele Giambra

December 3, 2017

Brown University

# Introduction

#### **US** Obesity Epidemics

- Obesity in the US has increased dramatically in the last decades
  - ♦ In late 1970s, 12.7% of men and 17% of women were obese
  - ♦ During 2011-2014, 34.3% of mean and 38.3% percent of women were obese
- Important health implications
  - Obesity increases inpatient and outpatient spending by 36% (Sturm, 2002)
  - ♦ Estimated annual medical cost of obesity is \$147 billion in 2008 US dollars (CDC)
  - Negative health outcomes related to obesity include heart disease, stroke, type 2 diabetes and certain type of cancer
- Childhood obesity is also a pervasive phenomenon
  - ♦ Obesity rates are 8.9% among 2- to 5-year-olds, 17.5% for 6to 11-year-olds and 20.5% of 12- to 19-year-olds

### **Definition of Obesity**

- Based on Body Mass Index (BMI)
  - ♦ Normal weight is a BMI between 18.5 and 24.9
  - ♦ Overweight is a BMI between 25.0 and 29.9
  - ♦ Obesity is a BMI over 30
- BMI measured as body mass (kg) divided by square of body height  $(m^2)$
- Different use of BMI for children aged 2 to 20
  - Comparison against percentile for children of same sex and age
  - ♦ BMI above the 95th percentile is considered obese


#### **School Nutrition Policies**

- National School Lunch Act of 1946
  - ⋄ Created the National School Lunch Program (NSLP)
  - ♦ Grown from 3,368 million lunches served in 1969 to 5,052 million in 2016
- Child Nutrition and WIC Reauthorization Act of 2004
  - Requires that school districts design and implement wellness policies from the 2006/2007 school year
- Healthy Hunger-Free Kids Act of 2010
  - Gives USDA the authority to set new standards for food sold in lunches (e.g. reduced portion sizes, minimum on fruit and vegetables per serving)
  - ♦ Increases access to NSLP
  - School districts audited every 3 years to see if they meet nutrition standards

#### Literature


- Eid, Overman, Puga and Turner (2008)
  - ⋄ Relationship between urban sprawl and obesity
  - Use NLSY79 data to track change in BMI for young adults moving into more sprawling neighborhoods
- Cutler, Glaeser and Shapiro (2003)
  - Increase in obesity since 1980 mostly driven by increase in calories consumed instead of decrease in calories expended
  - $\diamond$  Propose a theory based on technical change in food preparation
- Large descriptive literature on adult and childhood obesity
  - ♦ Li et al. (2015), Ng et al. (2014), Ogden et al. (2016)
- Impact of school nutrition environment
  - Anderson et al. (2017), Bauhoff (2014), Campbell et al. (2011), Schanzenbach (2009)

#### BMI Over Lifetime



- Source: National Health Interview Survey (NHIS)
- Data on adults age 18+ from 1976 through 2016, young adults age 12-17 from 2008

#### Secular Trend in BMI



- BMI in NHIS based on interviews:
  - ♦ Tendency to underreport weight and overreport height

#### Research Questions


How effective are school wellness policies in addressing childhood obesity?

Are students affected by the health status of their peers?

Potential Identification Strategies

## **School Eating Environment**


- Large time and geographic variation in school eating environment
- Example: strength of policies regulating standard for a la carte beverages in 2003, elementary school



Source: CLASS, 2003

## School Eating Environment

 Policies regulating standard for a la carte beverages in 2015, elementary school




Source: CLASS, 2015

## School Eating Environment: Data

- Classification of Laws Associated with School Students (CLASS)
  - Scoring system used to evaluate state-level school physical education and nutrition policies from 2003 through 2015
  - Developed by the National Cancer Institute
  - High correlation with other state law evaluation data (e.g. Bridging the Gap)
- National Health and Nutrition Examination Survey (NHANES)
  - ♦ Cross sectional surveys conducted biennially from 1999
  - 24-hour dietary recalls and physical examinations
  - ♦ Geocode variables are restricted use and require on-site access
- National Survey of Children's Health (NSCH)
  - ♦ State-level BMI data for kids 10-17 years
  - Interviews conducted in 2004, 2007, 2011 and 2016

#### Movers

- Non-trivial fraction of students change school
  - Bradbury et al. (2013) measure that 10.6% of non-high-school
    Boston Public School students have "nonstructural" school changes
- Different childhood obesity rates at the school district level (likely at the school level as well)



#### Movers: Data

- Early Childhood Longitudinal Study
  - ♦ ECLS-K follows kindergarten class of 1988-99 through 8th grade
    - ightarrow 22,700 children sampled from 1,300 schools
    - ightarrow 2,542 movers from spring of 1st grade and spring of 3rd grade
    - ightarrow Geolocation of the students' school and home in restricted use data
  - ♦ ECLS-K:2011 tracks cohort of 2010-2011 through 5th grade (Spring 3rd grade will be released in Jan 2018)
- Alternative: Massachusetts requires BMI screening for students in grades 1, 4, 7, and 10 since the 2010-2011 school year
- New York has similar requirement for grades K, 2, 4, 7, and 10
  - Administrative data on students moving within one of these states would probably give more power to the analysis

# References

Jean, Eid, Henry G. Overman, Diego Puga and Matthew A. Turner. 2008. Fat city: Questioning the relationship between urban sprawl and obesity. *Journal of Urban Economics* 63: 385-404.

David, M. Cutler, Edward L. Glaeser and Jesse M. Shapiro. 2003. Why have Americans become more obese? *Journal of Economic Perspectives* 17(3): 93-118.

Sturm, Roland. 2002. The effects of obesity, smoking, and drinking on medical problems and costs. *Health affairs* 21(2): 245-253.

Bradbury, Katharine, Mary A. Burke and Robert K. Triest. 2013. The effect of foreclosure on Boston Public School student academic performance. Working Papers, Federal Reserve Bank of Boston, No.13-12.