On recovering induced polarization information from airborne time domain EM data

Seogi Kang, Douglas W. Oldenburg, Dikun Yang, David Marchant, Geophysical Inversion Facility, University of British Columbia*

SUMMARY

XXX

XXX

Seigel (1959) Oldenburg and Li (1994) Wynn and Zonge (1975) Smith et al. (1988) Marchant et al. (2014) Flis et al. (1989) Pelton et al. (1978) Kang and Oldenburg (2016) Weidelt (1982)

REFERENCES

Flis, M. F., G. A. Newman, and G. W. Hohmann, 1989, Inducedpolarization effects in timedomain electromagnetic measurements: Geophysics, **54**, 514–523.

Kang, S., and D. W. Oldenburg, 2016, On recovering distributed IP information from inductive source time domain electromagnetic data (in revision): Geophysical Journal International.

Marchant, D., E. Haber, and D. Oldenburg, 2014, Three-dimensional modeling of IP effects in time-domain electromagnetic data: Geophysics, **79**, E303–E314.

Oldenburg, D., and Y. Li, 1994, Inversion of induced polarization data: Geophysics, **59**, 1327–1341.

Pelton, W., S. Ward, P. Hallof, W. Sill, and P. Nelson, 1978, Mineral discrimination and removal of inductive coupling with multifrequency IP: Geophysics, **43**, 588–609.

Seigel, H., 1959, Matehmatical formulation and type curves for induced polarization: Geophysics, **24**, 547–565.

Smith, R. S., P. Walker, B. Polzer, and G. F. West, 1988, The time-domain electromagnetic response of polarizable bodies: an approximate convolution algorithm: Geophysical Prospecting, **36**, 772–785.

Weidelt, P., 1982, Response characteristics of coincident loop transient electromagnetic systems: **47**, 1325–1330.

Wynn, J. C., and K. L. Zonge, 1975, EM coupling, its intrinsic value, its removal and the cultural coupling problem: Geophysics, **40**, 831–850.