

Cyclic difference families, perfect difference families and their applications

Tao Feng

Beijing Jiaotong University, Beijing
tfeng@bjtu.edu.cn

Let v be a positive odd integer. A (v, k, λ) -perfect difference family (PDF) is a collection \mathcal{F} of k -subsets of $\{0, 1, \dots, v-1\}$ such that the multiset $\bigcup_{F \in \mathcal{F}} \{x - y : x, y \in F, x > y\}$ covers each element of $\{1, 2, \dots, (v-1)/2\}$ exactly λ times. Perfect difference families are a special class of perfect systems of difference sets. This talk shows that a $(v, 4, \lambda)$ -PDF exists if and only if $\lambda(v-1) \equiv 0 \pmod{12}$, $v \geq 13$, and $(v, \lambda) \notin \{(25, 1), (37, 1)\}$. This result resolves a nearly 50-year-old conjecture posed by Bermond. Perfect difference families find applications in radio astronomy, optical orthogonal codes for optical code-division multiple access systems, geometric orthogonal codes for DNA origami, difference triangle sets, additive sequences of permutations, and graceful graph labelings. Perfect difference families can be also seen as a special kind of cyclic difference families. This talk will also give a survey on the recent progress on constructions for cyclic difference families.

References

- [1] Hengrui Liu, Tao Feng, Xiaomiao Wang, Menglong Zhang, *Perfect difference families, perfect systems of difference sets and their applications*, arXiv:2510.20446.
- [2] C. Zhao, Y. Chang, and T. Feng, *The existence of optimal $(v, 4, 1)$ optical orthogonal codes achieving the Johnson bound*, IEEE Trans. Inform. Theory, 70 (2024), 8746–8757.
- [3] C. Zhao, B. Zhao, Y. Chang, T. Feng, X. Wang, and M. Zhang, *Cyclic relative difference families with block size four and their applications*, J. Combin. Theory Ser. A, 206 (2024), 105890.