

Schizophrenia GBD Analysis Results

Contents

Overall Burden Trends	1
Country And Sex Contrasts	3
Segmented Trends	3
Age-Specific Patterns	4
Full Age-Year Heatmaps	11
Decomposition	13
Annual Chained Decomposition	14
Age-Period-Cohort Analysis	16
Surface Coverage	16
Incidence and DALY Age-Period Surfaces: China and United States	17
Marginal APC Summaries: China and United States	17
Cohort Relative-Risk Patterns: China and United States	18
Comparative Summary	19

This report mirrors the breast cancer analysis structure, adapted to the schizophrenia extract. The schizophrenia cause table contains incidence, prevalence, YLDs, and DALYs for China and the United States, with female and male strata. The source age structure uses **70+ years** as the top age bin. The report focuses on the available cause-burden measures in this extract.

Overall Burden Trends


Counts are all-age values and rates are age-standardized rates per 100,000. The **Percent basis** column identifies the age basis of the GBD **Percent** metric used for each measure; in this extract, incidence and prevalence use age-standardized percent rows when available, whereas DALYs and YLDs use all-age percent rows. DALY and YLD numbers and rates are numerically identical here because the extract has no YLL component for schizophrenia; their percent values can still differ because the GBD percent denominator is measure-specific.

Selected endpoint highlights:


- **Female Incidence.** China reached 132,520 in 2023 (+14.5% since 1990; ASR 18.42 per 100,000). The United States reached 36,071 (+13.2%; ASR 23.04 per 100,000).
- **Female Prevalence.** China reached 2,469,230 in 2023 (+59.9% since 1990; ASR 278.86 per 100,000). The United States reached 767,642 (+22.7%; ASR 403.27 per 100,000).
- **Female DALYs.** China reached 1,551,859 in 2023 (+57.5% since 1990; ASR 177.61 per 100,000). The United States reached 465,494 (+20.5%; ASR 247.86 per 100,000).
- **Male Incidence.** China reached 174,592 in 2023 (+14.1% since 1990; ASR 22.77 per 100,000). The United States reached 42,426 (+18.2%; ASR 26.42 per 100,000).
- **Male Prevalence.** China reached 2,925,661 in 2023 (+51.9% since 1990; ASR 324.40 per 100,000). The United States reached 800,475 (+31.6%; ASR 428.80 per 100,000).

- **Male DALYs.** China reached 1,867,497 in 2023 (+49.5% since 1990; ASR 209.14 per 100,000). The United States reached 499,087 (+29.2%; ASR 270.16 per 100,000).


Schizophrenia all-age counts in China and the United States (1990-2023)



Schizophrenia measure-specific GBD percent metric (1990-2023)



Schizophrenia age-standardized rates (1990-2023)



Country And Sex Contrasts

Ratios compare like-for-like point estimates within each metric. The **Measure-specific GBD percent** rows should be read as ratios of the GBD **Percent** metric for the same measure and age basis, not as a single common “share of all-cause burden” denominator across incidence, prevalence, YLDs, and DALYs.

Concise contrast summary:

- **Country rate contrast.** In 2023, China/United States age-standardized rate ratios for the main measures ranged from 0.69-0.86; values below 1.00 indicate lower ASRs in China than in the United States.
- **Country count contrast.** China’s 2023 all-age numbers were 3.22-4.12 times the corresponding U.S. counts across the main measures.
- **Sex contrast.** Male/female 2023 ratios ranged from 1.06-1.24 for ASRs and 1.04-1.32 for all-age numbers.
- **Largest ratio shifts.** China vs United States, Female DALYs Measure-specific GBD percent (+110.9%); China vs United States, Male DALYs Measure-specific GBD percent (+95.1%); China vs United States, Female DALYs All-age number (+30.7%).

Segmented Trends


Age-standardized rates are fitted with a compact BIC-selected segmented log-linear model, allowing up to two joinpoints per location-sex-measure series. The single-slope EAPC is retained alongside the segmented AAPC for continuity with common GBD trend reporting.

Segmented-model summary:

- **Incidence.** AAPC ranged from -0.21% to +0.01% across 4 location-sex panels; 4 panels selected at least one joinpoint. The comparable single-slope EAPC range was -0.25% to +0.00%.

- **Prevalence.** AAPC ranged from -0.23% to +0.02% across 4 location-sex panels; 4 panels selected at least one joinpoint. The comparable single-slope EAPC range was -0.28% to +0.03%.
- **DALYs.** AAPC ranged from -0.25% to +0.05% across 4 location-sex panels; 4 panels selected at least one joinpoint. The comparable single-slope EAPC range was -0.28% to +0.04%.


Segmented log-linear trends in schizophrenia age-standardized rates




Age-Specific Patterns


The age-specific panels retain the schizophrenia-specific top bin (70+ years). The heatmap difference panels show China relative to the United States within sex, measure, age, and year. Percent panels use the GBD Percent metric for each measure.


Age-specific schizophrenia numbers in 1990 and 2023







Age-specific schizophrenia GBD percent metric in 1990 and 2023






Age-specific schizophrenia rates in 1990 and 2023







Full Age-Year Heatmaps



Age-specific schizophrenia GBD percent-metric heatmaps, 1990-2023






Decomposition

The decomposition uses adult age-specific numbers and rates from 15-19 years through 70+ years. It decomposes 1990-2023 adult count change into adult population growth, adult population aging, and age-specific rate change using the average replacement approach. Population denominators are calculated from GBD count/rate pairs.

Endpoint decomposition highlights:

- **Incidence.** The largest adult count change was in China Male (+22,089). The dominant component was population growth (+48,764; +220.8% of total change).
- **Prevalence.** The largest adult count change was in China Male (+999,958). The dominant component was population growth (+720,519; +72.1% of total change).
- **DALYs.** The largest adult count change was in China Male (+619,033). The dominant component was population growth (+463,193; +74.8% of total change).

Decomposition of adult schizophrenia burden change, 1990-2023




Annual Chained Decomposition

The annual chained decomposition repeats the same population-growth, population-aging, and rate-change replacement decomposition for adjacent years, then accumulates components from 1990. As in the breast cancer analysis, component allocations differ from the direct 1990-2023 endpoint decomposition because the decomposition is path-dependent; total observed change should still agree apart from numerical tolerance.


Annual-chain decomposition checks:

- **Total-change agreement.** Annual-chain totals matched endpoint totals within 0.000000 counts across the main panels.
- **Path dependence.** Component allocation shifted most for China Male Prevalence: endpoint rate change was +6,207 versus +3,975 under annual chaining.

Annual chained decomposition of adult schizophrenia burden change



Interval chained decomposition of adult schizophrenia burden change



Age-Period-Cohort Analysis

This section follows the breast cancer report's APC layout. Annual age-specific schizophrenia observations from 1990-2023 are collapsed into recent-aligned five-year period blocks (1990-1993, 1994-1998, ..., 2019-2023) and summarized through net drift, age curves, local drift, period relative risks, and cohort relative risks.

Surface Coverage


The APC model coverage is summarized below without the full reference-group matrix.

APC coverage summary:

- Main-measure APC panels covered 1990-2023 with 11 age groups, 7 period blocks, 22 cohort groups, and 77 age-period cells per panel.
- Reference ages, periods, and cohorts remain available in the APC summary CSV, but the full panel matrix is omitted from the report body for readability.

Incidence and DALY Age-Period Surfaces: China and United States


The age-period rate surfaces are shown with China and the United States side by side within each sex and measure panel.



Marginal APC Summaries: China and United States

The marginal panels show fitted age curves, local drift, period relative risks, and cohort relative risks by measure and sex.


Schizophrenia ridge-regularized APC marginal effects



Cohort Relative-Risk Patterns: China and United States

The cohort panels compare China and the United States within each sex and measure.

Schizophrenia APC cohort relative-risk patterns



Comparative Summary

APC comparative highlights:

- **Net drift.** Largest absolute net-drift estimates: United States of America Female DALYs (-0.28%); United States of America Female Prevalence (-0.27%); United States of America Male DALYs (-0.12%); United States of America Male Prevalence (-0.10%).
- **Local drift.** Strongest local-drift estimates: United States of America Female Incidence at 20-24 years (-0.48%); United States of America Female DALYs at 20-24 years (-0.47%); United States of America Female Incidence at 15-19 years (-0.46%); United States of America Female Prevalence at 20-24 years (-0.46%).