



Quiz 2

[Back to Week 2](#)

9/10 points
earned (90%)

Quiz passed!



1 / 1
points

1.

Consider the following data with x as the predictor and y as the outcome.

```
1 x <- c(0.61, 0.93, 0.83, 0.35, 0.54, 0.16, 0.91, 0.62, 0.62)
2 y <- c(0.67, 0.84, 0.6, 0.18, 0.85, 0.47, 1.1, 0.65, 0.36)
```

Give a P-value for the two sided hypothesis test of whether β_1 from a linear regression model is 0 or not.



0.391



2.325



0.025



0.05296

Correct Response

```
1 summary(lm(y ~ x))$coef
```

1	##	Estimate	Std. Error	t value	Pr(> t)
2	## (Intercept)	0.1885	0.2061	0.9143	0.39098
3	## x	0.7224	0.3107	2.3255	0.05296

1 / 1
points

2.

Consider the previous problem, give the estimate of the residual standard deviation.

☐ 0.05296☐ 0.3552☒ 0.223**Correct Response**

```
1 summary(lm(y ~ x))$sigma
```

```
1 ## [1] 0.223
```

☐ 0.43581 / 1
points

3.

In the **mtcars** data set, fit a linear regression model of weight (predictor) on mpg (outcome). Get a 95% confidence interval for the expected mpg at the average weight. What is the lower endpoint?

☐ -6.486☒ 18.991**Correct Response**

```
1 data(mtcars)
2 fit <- lm(mpg ~ I(wt - mean(wt)), data = mtcars)
3 confint(fit)
```

```
1 ##                2.5 % 97.5 %
2 ## (Intercept)    18.991 21.190
3 ## I(wt - mean(wt)) -6.486 -4.203
```



-4.00



21.190

1 / 1
points

4.

Refer to the previous question. Read the help file for `mtcars`. What is the weight coefficient interpreted as?



The estimated expected change in mpg per 1,000 lb increase in weight.

**Correct Response**

This is the standard interpretation of a regression coefficient. The expected change in the response per unit change in the predictor.



It can't be interpreted without further information



The estimated 1,000 lb change in weight per 1 mpg increase.



The estimated expected change in mpg per 1 lb increase in weight.

1 / 1
points

5.

Consider again the `mtcars` data set and a linear regression model with mpg as predicted by weight (1,000 lbs). A new car is coming weighing 3000 pounds. Construct a 95% prediction interval for its mpg. What is the upper endpoint?



27.57

**Correct Response**

```
1 fit <- lm(mpg ~ wt, data = mtcars)
2 predict(fit, newdata = data.frame(wt = 3), interval = "prediction")
```

```
1 ##      fit    lwr    upr
2 ## 1 21.25 14.93 27.57
```

- ☐ -5.77
- ☐ 21.25
- ☐ 14.93



1 / 1
points

6.

Consider again the `mtcars` data set and a linear regression model with `mpg` as predicted by weight (in 1,000 lbs). A “short” ton is defined as 2,000 lbs. Construct a 95% confidence interval for the expected change in `mpg` per 1 short ton increase in weight. Give the lower endpoint.



-12.973

Correct Response

```
1 fit <- lm(mpg ~ wt, data = mtcars)
2 confint(fit)[2, ] * 2
```

```
1 ## 2.5 % 97.5 %
2 ## -12.973 -8.405
```

```
1 ## Or equivalently change the units
2 fit <- lm(mpg ~ I(wt * 0.5), data = mtcars)
3 confint(fit)[2, ]
```

```
1 ## 2.5 % 97.5 %
2 ## -12.973 -8.405
```

- ☐ -6.486
- ☐ -9.000
- ☐ 4.2026



0 / 1
points

7.

If my X from a linear regression is measured in centimeters and I convert it to meters what would happen to the slope coefficient?

- ☐ It would get multiplied by 10
- ☐ It would get multiplied by 100.
- ☐ It would get divided by 10
- ☒ It would get divided by 100

**Incorrect Response**1 / 1
points

8.

I have an outcome, Y , and a predictor, X and fit a linear regression model with $Y = \beta_0 + \beta_1 X + \epsilon$ to obtain $\hat{\beta}_0$ and $\hat{\beta}_1$. What would be the consequence to the subsequent slope and intercept if I were to refit the model with a new regressor, $X + c$ for some constant, c ?

- ☐ The new slope would be $\hat{\beta}_1 + c$
- ☐ The new slope would be $c\hat{\beta}_1$
- ☐ The new intercept would be $\hat{\beta}_0 + c\hat{\beta}_1$
- ☒ The new intercept would be $\hat{\beta}_0 - c\hat{\beta}_1$

**Correct Response**

This is exactly covered in the notes. But note that if $Y = \beta_0 + \beta_1 X + \epsilon$ then $Y = \beta_0 - c\beta_1 + \beta_1(X + c) + \epsilon$ so that the answer is that the intercept gets subtracted by $c\beta_1$

1 / 1
points

9.

Refer back to the mtcars data set with mpg as an outcome and weight (wt) as the predictor. About what is the ratio of the the sum of the squared errors, $\sum_{i=1}^n (Y_i - \hat{Y}_i)^2$ when comparing a model with just an intercept (denominator) to the model with the intercept and slope (numerator)?

- ☐ 4.00
- ☐ 0.50
- ☐ 0.75
- ☒ 0.25

Correct Response

This is simply one minus the R^2 values

```
1 fit1 <- lm(mpg ~ wt, data = mtcars)
2 fit2 <- lm(mpg ~ 1, data = mtcars)
3 1 - summary(fit1)$r.squared
```

```
1 ## [1] 0.2472
```

```
1 sse1 <- sum((predict(fit1) - mtcars$mpg)^2)
2 sse2 <- sum((predict(fit2) - mtcars$mpg)^2)
3 sse1/sse2
```

```
1 ## [1] 0.2472
```



1 / 1
points

10.

Do the residuals always have to sum to 0 in linear regression?

- ☒ If an intercept is included, then they will sum to 0.

Correct Response

They do provided an intercept is included. If not they most likely won't.



The residuals must always sum to zero

The residuals must always sum to zero.

- ☐ If an intercept is included, the residuals most likely won't sum to zero.
 - ☐ The residuals never sum to zero.
-

