

Common Diuca-Finch (Diuca diuca), type L34: the model reproduces this one

resynthesis cosine +0.971 — 14 member syllables with cached audio — fitted duration 141 ms, timbre class reed




Black-faced Grosbeak (Caryothraustes poliogaster), type L07: a median type

resynthesis cosine +0.725 — 10 member syllables with cached audio — fitted duration 101 ms, timbre class reed



Bolivian Tyrannulet (Zimmerius bolivianus), type L13: the model does not reproduce this one

resynthesis cosine −0.328 — 8 member syllables with cached audio — fitted duration 107 ms, timbre class reed



Each spectrogram: 22.05 kHz cached audio, 256-sample Hann window, 64-sample hop, peak-normalised, 70 dB range. Both sides of a row carry identical settings, identical axes and one colour scale. The resynthesis is rendered by the repository's own renderer at 48 kHz and resampled to the cache's rate; 20 ms of context is kept on the recording and matched with silence on the resynthesis, so the flat band at each panel's edge is the field recording's own noise floor. Third panel, grey: the measured pitch contour of every cached member of the type; slate: the member plotted to its left; violet: the contour the fit ships. The cosine is the objective the fit was run under; docs/19 measured its correlation with audible distortion at about +0.1, so it ranks the fit's own criterion and not audible fidelity. Production store, and the audio cache holds iNaturalist recordings only.