

Fuzzy logic, fuzzy sets and fuzzy relations

Shabana K M

PhD Research Scholar
Department of Computer Science and Engineering

IIT Palakkad

15 June 2021



INDIAN INSTITUTE
OF TECHNOLOGY
PALAKKAD



Fuzzy logic vs Classical logic

Classical logic

- based on two values - True and False
- might be inadequate to represent human reasoning
- cannot handle propositions with variable answers

Fuzzy logic vs Classical logic


Classical logic

- based on two values - True and False
- might be inadequate to represent human reasoning
- cannot handle propositions with variable answers

Fuzzy logic


- the truth value of variables may be any real number between 0 and 1, both inclusive
- handles the concept of partial truth
- represents vagueness and imprecise information

Fuzzy logic vs. Crisp logic




(a) Crisp logic

Fuzzy logic vs. Crisp logic




(a) Crisp logic



(b) Fuzzy logic

Fuzzy logic vs. Crisp logic



Fuzzy sets vs. Crisp sets

Sets

- a collection of elements, also referred to as members
- can be finite or infinite

Fuzzy sets vs. Crisp sets

Sets

- a collection of elements, also referred to as members
- can be finite or infinite

Crisp sets

- an element is either a member of the set or not
- example - red belongs to the set of colors whereas apple doesn't

Fuzzy sets vs. Crisp sets

Sets

- a collection of elements, also referred to as members
- can be finite or infinite


Crisp sets

- an element is either a member of the set or not
- example - red belongs to the set of colors whereas apple doesn't

Fuzzy Sets

- allow elements to be partially in a set
- each element given a degree of membership
- membership value ranges from 0 (not an element) to 1 (a member)
- example - consider the set of young people - a 100 year old person will not be a member, but people at the age of 20, 30, or 40 years can have varying degrees of membership

Fuzzy sets vs. Crisp sets



Fuzzy sets

- A fuzzy set A is characterized by a function $\mu_A : \mathcal{X} \rightarrow [0, 1]$
 - μ_A is called the membership function of A
 - defined over the universe of discourse \mathcal{X}
 - for each $x \in \mathcal{X}$, the value $\mu_A(x)$ is called the grade of membership of x in A

Fuzzy sets

- A fuzzy set A is characterized by a function $\mu_A : \mathcal{X} \rightarrow [0, 1]$
 - μ_A is called the membership function of A
 - defined over the universe of discourse \mathcal{X}
 - for each $x \in \mathcal{X}$, the value $\mu_A(x)$ is called the grade of membership of x in A
- Let $x \in \mathcal{X}$. Then x is called
 - **not included** in the fuzzy set A if $\mu(x) = 0$ (no member)
 - **fully included** if $\mu(x) = 1$ (full member)
 - **partially included** if $0 < \mu(x) < 1$ (fuzzy member)


Fuzzy sets

- A fuzzy set A is characterized by a function $\mu_A : \mathcal{X} \rightarrow [0, 1]$
 - μ_A is called the membership function of A
 - defined over the universe of discourse \mathcal{X}
 - for each $x \in \mathcal{X}$, the value $\mu_A(x)$ is called the grade of membership of x in A
- Let $x \in \mathcal{X}$. Then x is called
 - **not included** in the fuzzy set A if $\mu(x) = 0$ (no member)
 - **fully included** if $\mu(x) = 1$ (full member)
 - **partially included** if $0 < \mu(x) < 1$ (fuzzy member)
- Crisp sets are special cases of fuzzy sets with 0 and 1 as membership degrees

Fuzzy sets


- A fuzzy set A is characterized by a function $\mu_A : \mathcal{X} \rightarrow [0, 1]$
 - μ_A is called the membership function of A
 - defined over the universe of discourse \mathcal{X}
 - for each $x \in \mathcal{X}$, the value $\mu_A(x)$ is called the grade of membership of x in A
- Let $x \in \mathcal{X}$. Then x is called
 - **not included** in the fuzzy set A if $\mu(x) = 0$ (no member)
 - **fully included** if $\mu(x) = 1$ (full member)
 - **partially included** if $0 < \mu(x) < 1$ (fuzzy member)
- Crisp sets are special cases of fuzzy sets with 0 and 1 as membership degrees
- The set of elements with a non-zero membership is called the Support of the fuzzy set

Membership function for Fuzzy sets¹




¹Image source: Google

Fuzzy set representation




Fuzzy set representation

- A fuzzy set is often represented as a pair (\mathcal{X}, μ)




Fuzzy set representation




- A fuzzy set is often represented as a pair (\mathcal{X}, μ)
- $\mathbb{F}(\mathcal{X})$ denote the set of all fuzzy sets on \mathcal{X}

Fuzzy set representation




- A fuzzy set is often represented as a pair (\mathcal{X}, μ)
- $\mathbb{F}(\mathcal{X})$ denote the set of all fuzzy sets on \mathcal{X}
- For finite universes \mathcal{X} , a fuzzy set can be represented as $A = \sum_{x \in \mathcal{X}} \mu(x)/x$

Fuzzy set representation



- A fuzzy set is often represented as a pair (\mathcal{X}, μ)
- $\mathbb{F}(\mathcal{X})$ denote the set of all fuzzy sets on \mathcal{X}
- For finite universes \mathcal{X} , a fuzzy set can be represented as $A = \sum_{x \in \mathcal{X}} \mu(x)/x$
- and for infinite universes \mathcal{X} as $A = \int_{x \in \mathcal{X}} \mu(x)/x$

Fuzzy set representation



- A fuzzy set is often represented as a pair (\mathcal{X}, μ)
- $\mathbb{F}(\mathcal{X})$ denote the set of all fuzzy sets on \mathcal{X}
- For finite universes \mathcal{X} , a fuzzy set can be represented as $A = \sum_{x \in \mathcal{X}} \mu(x)/x$
- and for infinite universes \mathcal{X} as $A = \int_{x \in \mathcal{X}} \mu(x)/x$
- In these representations, \sum, \int and $/$ have only symbolic meaning

Fuzzy sets - Examples




Figure: Consider the fuzzy set $A = \{\text{real numbers near } 0\}$ represented as $A = \int_{x \in \mathcal{X}} \mu(x)/x$ where $\mu(x) = \frac{1}{1+x^2}$

Fuzzy sets - Examples

- A fuzzy membership function is different from a statistical probability distribution

Fuzzy sets - Examples


- A fuzzy membership function is different from a statistical probability distribution
- Consider the fuzzy set A defined as $\{\text{two or so}\}$. In this case, the universal set \mathcal{X} is the set of positive integers

Fuzzy sets - Examples

- A fuzzy membership function is different from a statistical probability distribution
- Consider the fuzzy set A defined as {two or so}. In this case, the universal set \mathcal{X} is the set of positive integers
- A membership function $\mu_A(x)$ can be defined as follows:
 $\mu_A(1) = 0.5, \mu_A(2) = 1, \mu_A(3) = 0.5, \mu_A(4) = 0, \dots$

Fuzzy sets - Examples

- A fuzzy membership function is different from a statistical probability distribution
- Consider the fuzzy set A defined as {two or so}. In this case, the universal set \mathcal{X} is the set of positive integers
- A membership function $\mu_A(x)$ can be defined as follows:
 $\mu_A(1) = 0.5, \mu_A(2) = 1, \mu_A(3) = 0.5, \mu_A(4) = 0, \dots$



Fuzzy sets - Other definitions

- **Support:** $\text{supp}(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$

Fuzzy sets - Other definitions

- **Support:** $\text{supp}(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$
- The fuzzy set A is said to be **empty** ($A = \emptyset$) iff $\forall x \in \mathcal{X} : \mu_A(x) = 0$

Fuzzy sets - Other definitions

- **Support:** $\text{supp}(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$
- The fuzzy set A is said to be **empty** ($A = \emptyset$) iff $\forall x \in \mathcal{X} : \mu_A(x) = 0$
- **Height:** $\text{hgt}(A) = \sup_{x \in \mathcal{X}} \mu_A(x)$

Fuzzy sets - Other definitions

- **Support:** $\text{supp}(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$
- The fuzzy set A is said to be **empty** ($A = \emptyset$) iff $\forall x \in \mathcal{X} : \mu_A(x) = 0$
- **Height:** $\text{hgt}(A) = \sup_{x \in \mathcal{X}} \mu_A(x)$
- The fuzzy set A is said to be **unimodal** if $\text{hgt}(A)$ is the membership degree for only one element of \mathcal{X}

Fuzzy sets - Other definitions

- **Support:** $\text{supp}(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$
- The fuzzy set A is said to be **empty** ($A = \emptyset$) iff $\forall x \in \mathcal{X} : \mu_A(x) = 0$
- **Height:** $\text{hgt}(A) = \sup_{x \in \mathcal{X}} \mu_A(x)$
- The fuzzy set A is said to be **unimodal** if $\text{hgt}(A)$ is the membership degree for only one element of \mathcal{X}
- The fuzzy set A is said to be **normalized** iff $\text{hgt}(A) = 1$

Fuzzy sets - Other definitions

- **Support:** $\text{supp}(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$
- The fuzzy set A is said to be **empty** ($A = \emptyset$) iff $\forall x \in \mathcal{X} : \mu_A(x) = 0$
- **Height:** $\text{hgt}(A) = \sup_{x \in \mathcal{X}} \mu_A(x)$
- The fuzzy set A is said to be **unimodal** if $\text{hgt}(A)$ is the membership degree for only one element of \mathcal{X}
- The fuzzy set A is said to be **normalized** iff $\text{hgt}(A) = 1$
 - The fuzzy sets other than \emptyset with height less than 1 are said to be **subnormal**
 - A is subnormal $\Leftrightarrow 0 < \text{hgt}(A) < 1$
 - A non-empty subnormal fuzzy set A can be normalized into the set A^* by dividing the membership function of A by $\text{hgt}(A)$
 $\forall x \in \mathcal{X} : \mu_{A^*}(x) = \mu_A(x) / \text{hgt}(A)$

Fuzzy sets - Other definitions

- **Cardinality:** For a fuzzy set A with finite support,
$$card(A) = \sum_{x \in \mathcal{A}} \mu_A(x)$$

Fuzzy sets - Other definitions

- **Cardinality:** For a fuzzy set A with finite support,
$$\text{card}(A) = \sum_{x \in \mathcal{X}} \mu_A(x)$$
- A fuzzy set in whose support is single point in \mathcal{X} with membership equal to one is called a **fuzzy singleton**

Fuzzy sets - Other definitions

- **Cardinality:** For a fuzzy set A with finite support,
$$\text{card}(A) = \sum_{x \in \mathcal{A}} \mu_A(x)$$
- A fuzzy set in whose support is single point in \mathcal{X} with membership equal to one is called a **fuzzy singleton**
- Two fuzzy sets A and B are equal ($A = B$) iff
$$\forall x \in \mathcal{X} : \mu_A(x) = \mu_B(x)$$

Fuzzy sets - Other definitions

- **Cardinality:** For a fuzzy set A with finite support,
 $card(A) = \sum_{x \in \mathcal{A}} \mu_A(x)$
- A fuzzy set in whose support is single point in \mathcal{X} with membership equal to one is called a **fuzzy singleton**
- Two fuzzy sets A and B are equal ($A = B$) iff
 $\forall x \in \mathcal{X} : \mu_A(x) = \mu_B(x)$
- A fuzzy set A is said to be included in a fuzzy set B , ($A \subseteq B$) iff
 $\forall x \in \mathcal{X} : \mu_A(x) \leq \mu_B(x)$

Fuzzy sets - Other definitions

- **Cardinality:** For a fuzzy set A with finite support,
 $card(A) = \sum_{x \in \mathcal{A}} \mu_A(x)$
- A fuzzy set in whose support is single point in \mathcal{X} with membership equal to one is called a **fuzzy singleton**
- Two fuzzy sets A and B are equal ($A = B$) iff
 $\forall x \in \mathcal{X} : \mu_A(x) = \mu_B(x)$
- A fuzzy set A is said to be included in a fuzzy set B , ($A \subseteq B$) iff
 $\forall x \in \mathcal{X} : \mu_A(x) \leq \mu_B(x)$
- For a fuzzy set A any element $x \in \mathcal{X}$ that satisfies $\mu_A(x) = 0.5$ is called a **crossover point**

Crisp sets related to fuzzy sets

For a fuzzy set $A = (\mathcal{X}, \mu)$ and $\alpha \in [0, 1]$

- $A^{\geq \alpha} = \{x \in \mathcal{X} : \mu_A(x) \geq \alpha\}$ is called its **α -cut**

Crisp sets related to fuzzy sets

For a fuzzy set $A = (\mathcal{X}, \mu)$ and $\alpha \in [0, 1]$

- $A^{\geq \alpha} = \{x \in \mathcal{X} : \mu_A(x) \geq \alpha\}$ is called its **α -cut**
- $A^{> \alpha} = \{x \in \mathcal{X} : \mu_A(x) > \alpha\}$ is called its **strong α -cut**

Crisp sets related to fuzzy sets

For a fuzzy set $A = (\mathcal{X}, \mu)$ and $\alpha \in [0, 1]$

- $A^{\geq \alpha} = \{x \in \mathcal{X} : \mu_A(x) \geq \alpha\}$ is called its **α -cut**
- $A^{> \alpha} = \{x \in \mathcal{X} : \mu_A(x) > \alpha\}$ is called its **strong α -cut**
- **Support of A** $supp(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$ is also a strong α -cut $A^{>0}$

Crisp sets related to fuzzy sets

For a fuzzy set $A = (\mathcal{X}, \mu)$ and $\alpha \in [0, 1]$


- $A^{\geq \alpha} = \{x \in \mathcal{X} : \mu_A(x) \geq \alpha\}$ is called its **α -cut**
- $A^{> \alpha} = \{x \in \mathcal{X} : \mu_A(x) > \alpha\}$ is called its **strong α -cut**
- **Support of A** $supp(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$ is also a strong α -cut $A^{>0}$
- The 1-cut $A^{\geq 1} = \{x \in \mathcal{X} : \mu_A(x) = 1\}$ is called the **core/kernel** of A

Crisp sets related to fuzzy sets

For a fuzzy set $A = (\mathcal{X}, \mu)$ and $\alpha \in [0, 1]$


- $A^{\geq \alpha} = \{x \in \mathcal{X} : \mu_A(x) \geq \alpha\}$ is called its **α -cut**
- $A^{> \alpha} = \{x \in \mathcal{X} : \mu_A(x) > \alpha\}$ is called its **strong α -cut**
- **Support of A** $supp(A) = \{x \in \mathcal{X} : \mu_A(x) > 0\}$ is also a strong α -cut $A^{>0}$
- The 1-cut $A^{\geq 1} = \{x \in \mathcal{X} : \mu_A(x) = 1\}$ is called the **core/kernel** of A
- The **level** of a fuzzy set A is defined as $A^{\alpha} = \{x \in \mathcal{X} : \mu_A(x) = \alpha\}$

Crisp sets related to fuzzy sets - Example




- This set can be represented as
$$A = 0.0/0 + 0.0/1 + 0.5/2 + 0.8/3 + 1/4 + 0.7/5 + 0.3/6 + 0.0/7 + 0.0/8 + 0.0/9$$
- $supp(A) =$

Crisp sets related to fuzzy sets - Example




- This set can be represented as
$$A = 0.0/0 + 0.0/1 + 0.5/2 + 0.8/3 + 1/4 + 0.7/5 + 0.3/6 + 0.0/7 + 0.0/8 + 0.0/9$$
- $\text{supp}(A) = \{2, 3, 4, 5, 6\}$
- $\text{hgt}(A) =$

Crisp sets related to fuzzy sets - Example




- This set can be represented as
$$\mathbf{A} = 0.0/0 + 0.0/1 + 0.5/2 + 0.8/3 + 1/4 + 0.7/5 + 0.3/6 + 0.0/7 + 0.0/8 + 0.0/9$$
- $\text{supp}(A) = \{2, 3, 4, 5, 6\}$
- $\text{hgt}(A) = 1 \rightarrow A$ is normalized!
- \mathbf{A} is

Crisp sets related to fuzzy sets - Example




- This set can be represented as
$$\mathbf{A} = 0.0/0 + 0.0/1 + 0.5/2 + 0.8/3 + 1/4 + 0.7/5 + 0.3/6 + 0.0/7 + 0.0/8 + 0.0/9$$
- $\text{supp}(A) = \{2, 3, 4, 5, 6\}$
- $\text{hgt}(A) = 1 \rightarrow A$ is normalized!
- \mathbf{A} is unimodal
- $\text{core}(A) =$

Crisp sets related to fuzzy sets - Example




- This set can be represented as
$$\mathbf{A} = 0.0/0 + 0.0/1 + 0.5/2 + 0.8/3 + 1/4 + 0.7/5 + 0.3/6 + 0.0/7 + 0.0/8 + 0.0/9$$
- $\text{supp}(A) = \{2, 3, 4, 5, 6\}$
- $\text{hgt}(A) = 1 \rightarrow A$ is normalized!
- \mathbf{A} is unimodal
- $\text{core}(A) = \{4\}$
- $\text{crossover}(A) =$

Crisp sets related to fuzzy sets - Example




- This set can be represented as
 $A = 0.0/0 + 0.0/1 + 0.5/2 + 0.8/3 + 1/4 + 0.7/5 + 0.3/6 + 0.0/7 + 0.0/8 + 0.0/9$
- $\text{supp}(A) = \{2, 3, 4, 5, 6\}$
- $\text{hgt}(A) = 1 \rightarrow A$ is normalized!
- A is unimodal
- $\text{core}(A) = \{4\}$
- $\text{crossover}(A) = \{2\}$
- $A^{>0.5} =$

Crisp sets related to fuzzy sets - Example



- This set can be represented as
 $A = 0.0/0 + 0.0/1 + 0.5/2 + 0.8/3 + 1/4 + 0.7/5 + 0.3/6 + 0.0/7 + 0.0/8 + 0.0/9$
- $\text{supp}(A) = \{2, 3, 4, 5, 6\}$
- $\text{hgt}(A) = 1 \rightarrow A$ is normalized!
- A is unimodal
- $\text{core}(A) = \{4\}$
- $\text{crossover}(A) = \{2\}$
- $A^{>0.5} = \{3, 4, 5\}$
- The 0.3-level set $A^{=0.3} =$

Crisp sets related to fuzzy sets - Example



- This set can be represented as
 $A = 0.0/0 + 0.0/1 + 0.5/2 + 0.8/3 + 1/4 + 0.7/5 + 0.3/6 + 0.0/7 + 0.0/8 + 0.0/9$
- $\text{supp}(A) = \{2, 3, 4, 5, 6\}$
- $\text{hgt}(A) = 1 \rightarrow A$ is normalized!
- A is unimodal
- $\text{core}(A) = \{4\}$
- $\text{crossover}(A) = \{2\}$
- $A^{>0.5} = \{3, 4, 5\}$
- The 0.3-level set $A^{=0.3} = \{6\}$

Fuzzy set operations

- **Union $A \cup B$:** $\mu_{A \cup B}(x) = \max\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$

Fuzzy set operations

- **Union $A \cup B$:** $\mu_{A \cup B}(x) = \max\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Intersection $A \cap B$:** $\mu_{A \cap B}(x) = \min\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$

Fuzzy set operations

- **Union $A \cup B$:** $\mu_{A \cup B}(x) = \max\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Intersection $A \cap B$:** $\mu_{A \cap B}(x) = \min\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Complement A^c :** $\mu_{A^c}(x) = 1 - \mu_A(x)$

Fuzzy set operations

- **Union $A \cup B$:** $\mu_{A \cup B}(x) = \max\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Intersection $A \cap B$:** $\mu_{A \cap B}(x) = \min\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Complement A^c :** $\mu_{A^c}(x) = 1 - \mu_A(x)$
- **Power of a fuzzy set A^α :** $\mu_{A^\alpha}(x) = \{\mu_A(x)\}^\alpha$ for all $x \in \mathcal{X}$

Fuzzy set operations

- **Union $A \cup B$:** $\mu_{A \cup B}(x) = \max\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Intersection $A \cap B$:** $\mu_{A \cap B}(x) = \min\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Complement A^c :** $\mu_{A^c}(x) = 1 - \mu_A(x)$
- **Power of a fuzzy set A^α :** $\mu_{A^\alpha}(x) = \{\mu_A(x)\}^\alpha$ for all $x \in \mathcal{X}$
- **Cartesian Product $A \times B$:**
 $\mu_{A \times B}(x, y) = \min\{\mu_A(x), \mu_B(y)\}$ for all $x \in \mathcal{X}, y \in \mathcal{Y}$

Example

- $\mathcal{X} = \{x_1, x_2, x_3, x_4\}$
- $A = 0.5/x_1 + 0.4/x_2 + 0.6/x_4$
- $B = 0.1/x_1 + 0.5/x_2 + 1/x_3 + 0.1/x_4$

Fuzzy set operations

- **Union $A \cup B$:** $\mu_{A \cup B}(x) = \max\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Intersection $A \cap B$:** $\mu_{A \cap B}(x) = \min\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Complement A^c :** $\mu_{A^c}(x) = 1 - \mu_A(x)$
- **Power of a fuzzy set A^α :** $\mu_{A^\alpha}(x) = \{\mu_A(x)\}^\alpha$ for all $x \in \mathcal{X}$
- **Cartesian Product $A \times B$:**
 $\mu_{A \times B}(x, y) = \min\{\mu_A(x), \mu_B(y)\}$ for all $x \in \mathcal{X}, y \in \mathcal{Y}$

Example

- $\mathcal{X} = \{x_1, x_2, x_3, x_4\}$
- $A = 0.5/x_1 + 0.4/x_2 + 0.6/x_4$
- $B = 0.1/x_1 + 0.5/x_2 + 1/x_3 + 0.1/x_4$
- $A \cup B = 0.5/x_1 + 0.5/x_2 + 1/x_3 + 0.6/x_4$

Fuzzy set operations

- **Union $A \cup B$:** $\mu_{A \cup B}(x) = \max\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Intersection $A \cap B$:** $\mu_{A \cap B}(x) = \min\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Complement A^c :** $\mu_{A^c}(x) = 1 - \mu_A(x)$
- **Power of a fuzzy set A^α :** $\mu_{A^\alpha}(x) = \{\mu_A(x)\}^\alpha$ for all $x \in \mathcal{X}$
- **Cartesian Product $A \times B$:**
 $\mu_{A \times B}(x, y) = \min\{\mu_A(x), \mu_B(y)\}$ for all $x \in \mathcal{X}, y \in \mathcal{Y}$

Example

- $\mathcal{X} = \{x_1, x_2, x_3, x_4\}$
- $A = 0.5/x_1 + 0.4/x_2 + 0.6/x_4$
- $B = 0.1/x_1 + 0.5/x_2 + 1/x_3 + 0.1/x_4$
- $A \cup B = 0.5/x_1 + 0.5/x_2 + 1/x_3 + 0.6/x_4$
- $A \cap B = 0.1/x_1 + 0.4/x_2 + 0/x_3 + 0.1/x_4$


Fuzzy set operations

- **Union $A \cup B$:** $\mu_{A \cup B}(x) = \max\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Intersection $A \cap B$:** $\mu_{A \cap B}(x) = \min\{\mu_A(x), \mu_B(x)\}$ for all $x \in \mathcal{X}$
- **Complement A^c :** $\mu_{A^c}(x) = 1 - \mu_A(x)$
- **Power of a fuzzy set A^α :** $\mu_{A^\alpha}(x) = \{\mu_A(x)\}^\alpha$ for all $x \in \mathcal{X}$
- **Cartesian Product $A \times B$:**
 $\mu_{A \times B}(x, y) = \min\{\mu_A(x), \mu_B(y)\}$ for all $x \in \mathcal{X}, y \in \mathcal{Y}$


Example

- $\mathcal{X} = \{x_1, x_2, x_3, x_4\}$
- $A = 0.5/x_1 + 0.4/x_2 + 0.6/x_4$
- $B = 0.1/x_1 + 0.5/x_2 + 1/x_3 + 0.1/x_4$
- $A \cup B = 0.5/x_1 + 0.5/x_2 + 1/x_3 + 0.6/x_4$
- $A \cap B = 0.1/x_1 + 0.4/x_2 + 0/x_3 + 0.1/x_4$
- $A^c = 0.5/x_1 + 0.6/x_2 + 1.0/x_3 + 0.4/x_4$




Fuzzy set operations




Fuzzy set operations



Fuzzy set operations



Fuzzy set operations



Properties of fuzzy sets

Commutativity

$$A \cup B = B \cup A$$

$$A \cap B = B \cap A$$

Properties of fuzzy sets

Commutativity

$$A \cup B = B \cup A$$

$$A \cap B = B \cap A$$

Associativity

$$A \cup (B \cup C) = (A \cup B) \cup C$$

$$A \cap (B \cap C) = (A \cap B) \cap C$$

Properties of fuzzy sets

Commutativity

$$A \cup B = B \cup A$$

$$A \cap B = B \cap A$$

Associativity

$$A \cup (B \cup C) = (A \cup B) \cup C$$

$$A \cap (B \cap C) = (A \cap B) \cap C$$

Distributivity

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

Properties of fuzzy sets

Idempotence

$$A \cup \mathcal{X} = \mathcal{X}, A \cap \mathcal{X} = A$$

$$A \cup A = A, A \cap A = A$$

$$A \cup \emptyset = A, A \cap \emptyset = \emptyset$$

$$(A^c)^c = A$$

Properties of fuzzy sets

Idempotence

$$A \cup \mathcal{X} = \mathcal{X}, A \cap \mathcal{X} = A$$

$$A \cup A = A, A \cap A = A$$

$$A \cup \emptyset = A, A \cap \emptyset = \emptyset$$

$$(A^c)^c = A$$

De Morgan's Laws

$$(A \cap B)^c = A^c \cup B^c$$

$$(A \cup B)^c = A^c \cap B^c$$

Properties of fuzzy sets

Idempotence

$$A \cup \mathcal{X} = \mathcal{X}, A \cap \mathcal{X} = A$$

$$A \cup A = A, A \cap A = A$$

$$A \cup \emptyset = A, A \cap \emptyset = \emptyset$$

$$(A^c)^c = A$$

De Morgan's Laws

$$(A \cap B)^c = A^c \cup B^c$$

$$(A \cup B)^c = A^c \cap B^c$$

Transitivity

If $A \subseteq B$, $B \subseteq C$, then $A \subseteq C$

Fuzzy relations

- fuzzy sets defined over the Cartesian product of the universe of discourses $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n$

Fuzzy relations

- fuzzy sets defined over the Cartesian product of the universe of discourses $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n$
- map elements of one universe, say \mathcal{X}_1 , to those of another universe, say \mathcal{X}_2 , and so on

Fuzzy relations

- fuzzy sets defined over the Cartesian product of the universe of discourses $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n$
- map elements of one universe, say \mathcal{X}_1 , to those of another universe, say \mathcal{X}_2 , and so on
- the Cartesian product, denoted as, $\mathcal{X}_1 \times \mathcal{X}_2 \times \dots \times \mathcal{X}_n$ is a collection of ordered pairs, such that
$$\mathcal{X}_1 \times \mathcal{X}_2 \times \dots \times \mathcal{X}_n = \{(x_1, x_2, \dots, x_n) \mid x_1 \in \mathcal{X}_1, x_2 \in \mathcal{X}_2, \dots, x_n \in \mathcal{X}_n\}$$

Fuzzy relations

- fuzzy sets defined over the Cartesian product of the universe of discourses $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n$
- map elements of one universe, say \mathcal{X}_1 , to those of another universe, say \mathcal{X}_2 , and so on
- the Cartesian product, denoted as, $\mathcal{X}_1 \times \mathcal{X}_2 \times \dots \times \mathcal{X}_n$ is a collection of ordered pairs, such that
$$\mathcal{X}_1 \times \mathcal{X}_2 \times \dots \times \mathcal{X}_n = \{(x_1, x_2, \dots, x_n) \mid x_1 \in \mathcal{X}_1, x_2 \in \mathcal{X}_2, \dots, x_n \in \mathcal{X}_n\}$$
- within a relation R , the n -tuples (x_1, x_2, \dots, x_n) have varying degree of memberships given by $\mu_R(x_1, x_2, \dots, x_n)$

Fuzzy relations

- fuzzy sets defined over the Cartesian product of the universe of discourses $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n$
- map elements of one universe, say \mathcal{X}_1 , to those of another universe, say \mathcal{X}_2 , and so on
- the Cartesian product, denoted as, $\mathcal{X}_1 \times \mathcal{X}_2 \times \dots \times \mathcal{X}_n$ is a collection of ordered pairs, such that
$$\mathcal{X}_1 \times \mathcal{X}_2 \times \dots \times \mathcal{X}_n = \{(x_1, x_2, \dots, x_n) \mid x_1 \in \mathcal{X}_1, x_2 \in \mathcal{X}_2, \dots, x_n \in \mathcal{X}_n\}$$
- within a relation R , the n -tuples (x_1, x_2, \dots, x_n) have varying degree of memberships given by $\mu_R(x_1, x_2, \dots, x_n)$
- membership values indicate the strength of the relation between the tuples

Fuzzy relations - Example

- $\mathcal{X} = \{\text{typhoid, viral, cold}\}$, $\mathcal{Y} = \{\text{running nose, high temp, shivering}\}$

Fuzzy relations - Example

- $\mathcal{X} = \{\text{typhoid, viral, cold}\}$, $\mathcal{Y} = \{\text{running nose, high temp, shivering}\}$
- The fuzzy relation \mathbf{R} over $\mathcal{X} \times \mathcal{Y}$ can be represented by a matrix with elements from $[0, 1]$ as follows:

Fuzzy relations - Example

- $\mathcal{X} = \{\text{typhoid, viral, cold}\}$, $\mathcal{Y} = \{\text{running nose, high temp, shivering}\}$
- The fuzzy relation \mathbf{R} over $\mathcal{X} \times \mathcal{Y}$ can be represented by a matrix with elements from $[0, 1]$ as follows:

$$\begin{array}{l} \text{typhoid} \\ \text{viral} \\ \text{cold} \end{array} \begin{pmatrix} \text{runningnose} & \text{hightemp} & \text{shivering} \\ 0.1 & 0.9 & 0.8 \\ 0.2 & 0.9 & 0.7 \\ 0.9 & 0.4 & 0.6 \end{pmatrix}$$

Fuzzy relations - Example

- $\mathcal{X} = \{\text{typhoid}, \text{viral}, \text{cold}\}$, $\mathcal{Y} = \{\text{running nose}, \text{high temp}, \text{shivering}\}$
- The fuzzy relation \mathbf{R} over $\mathcal{X} \times \mathcal{Y}$ can be represented by a matrix with elements from $[0, 1]$ as follows:

	<i>runningnose</i>	<i>hightemp</i>	<i>shivering</i>
<i>typhoid</i>	0.1	0.9	0.8
<i>viral</i>	0.2	0.9	0.7
<i>cold</i>	0.9	0.4	0.6

- $\mathbf{A} = 0.2/a_1 + 0.7/a_2 + 0.4/a_3$, $\mathbf{B} = 0.5/b_1 + 0.6/b_2$
- $\mu_{A \times B}(x, y) = \min\{\mu_A(x), \mu_B(y)\}$ for all $x \in \mathcal{X}$

Fuzzy relations - Example

- $\mathcal{X} = \{\text{typhoid}, \text{viral}, \text{cold}\}$, $\mathcal{Y} = \{\text{running nose}, \text{high temp}, \text{shivering}\}$
- The fuzzy relation \mathbf{R} over $\mathcal{X} \times \mathcal{Y}$ can be represented by a matrix with elements from $[0, 1]$ as follows:

$$\begin{array}{l} \text{typhoid} \\ \text{viral} \\ \text{cold} \end{array} \begin{pmatrix} \text{runningnose} & \text{hightemp} & \text{shivering} \\ 0.1 & 0.9 & 0.8 \\ 0.2 & 0.9 & 0.7 \\ 0.9 & 0.4 & 0.6 \end{pmatrix}$$


- $\mathbf{A} = 0.2/a_1 + 0.7/a_2 + 0.4/a_3$, $\mathbf{B} = 0.5/b_1 + 0.6/b_2$
- $\mu_{A \times B}(x, y) = \min\{\mu_A(x), \mu_B(y)\}$ for all $x \in \mathcal{X}$

$$\mathbf{R} = \mathbf{A} \times \mathbf{B} = \begin{array}{l} a_1 \\ a_2 \\ a_3 \end{array} \begin{pmatrix} b_1 & b_2 \\ 0.2 & 0.2 \\ 0.5 & 0.6 \\ 0.4 & 0.4 \end{pmatrix}$$

Fuzzy relations - Example

Consider the relation R : "x is considerably larger than y", where $x, y \in \mathbb{R}$

$$R(x, y) = \begin{cases} 0 & \text{for } x \leq y \\ \{x - y\} / (10y), & \text{for } y < x \leq 11y \\ 1 & \text{for } x > 11y \end{cases}$$



Operations on fuzzy relations

Since fuzzy relations are only special fuzzy sets:

- propositions that hold true for fuzzy sets also hold true for fuzzy relations
- operations defined for fuzzy sets also applicable to fuzzy relations

Operations on fuzzy relations

Since fuzzy relations are only special fuzzy sets:

- propositions that hold true for fuzzy sets also hold true for fuzzy relations
- operations defined for fuzzy sets also applicable to fuzzy relations

Let R and S be two fuzzy relations on $A \times B$

Union: $\mu_{R \cup S}(a, b) = \max\{\mu_R(a, b), \mu_S(a, b)\}$

Intersection: $\mu_{R \cap S}(a, b) = \min\{\mu_R(a, b), \mu_S(a, b)\}$

Complement: $\mu_{R^c}(a, b) = 1 - \mu_R(a, b)$

Operations on fuzzy relations

Since fuzzy relations are only special fuzzy sets:

- propositions that hold true for fuzzy sets also hold true for fuzzy relations
- operations defined for fuzzy sets also applicable to fuzzy relations

Let R and S be two fuzzy relations on $A \times B$

Union: $\mu_{R \cup S}(a, b) = \max\{\mu_R(a, b), \mu_S(a, b)\}$

Intersection: $\mu_{R \cap S}(a, b) = \min\{\mu_R(a, b), \mu_S(a, b)\}$

Complement: $\mu_{R^c}(a, b) = 1 - \mu_R(a, b)$

Inverse: $\mu_{R^{-1}}(a, b) = \mu_R(b, a)$

Operations on fuzzy relations

Since fuzzy relations are only special fuzzy sets:

- propositions that hold true for fuzzy sets also hold true for fuzzy relations
- operations defined for fuzzy sets also applicable to fuzzy relations

Let R and S be two fuzzy relations on $A \times B$

Union: $\mu_{R \cup S}(a, b) = \max\{\mu_R(a, b), \mu_S(a, b)\}$

Intersection: $\mu_{R \cap S}(a, b) = \min\{\mu_R(a, b), \mu_S(a, b)\}$

Complement: $\mu_{R^c}(a, b) = 1 - \mu_R(a, b)$

Inverse: $\mu_{R^{-1}}(a, b) = \mu_R(b, a)$

Composition: For fuzzy relations $R \in \mathbb{F}(\mathcal{X}_1 \times \mathcal{X}_2)$ and $S \in \mathbb{F}(\mathcal{X}_2 \times \mathcal{X}_3)$, their composition or relational product $R \circ S$ is defined as:

$$\mu_{R \circ S}(x, y) = \max_{z \in \mathcal{X}_2} \min\{\mu_R(x, z), \mu_S(z, y)\} \text{ for all } (x, y) \in \mathcal{X}_1 \times \mathcal{X}_3$$

Composition of fuzzy relations: Example

$$X = (x_1, x_2, x_3), Y = (y_1, y_2), Z = (z_1, z_2, z_3)$$

$$R = \begin{matrix} & y_1 & y_2 \\ \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix} & \begin{pmatrix} 0.5 & 0.1 \\ 0.2 & 0.9 \\ 0.8 & 0.6 \end{pmatrix} \end{matrix}$$

$$S = \begin{matrix} & z_1 & z_2 & z_3 \\ \begin{matrix} y_1 \\ y_2 \end{matrix} & \begin{pmatrix} 0.6 & 0.4 & 0.7 \\ 0.5 & 0.8 & 0.9 \end{pmatrix} \end{matrix}$$

Composition of fuzzy relations: Example

$$X = (x_1, x_2, x_3), \quad Y = (y_1, y_2), \quad Z = (z_1, z_2, z_3)$$

$$R = \begin{matrix} & \begin{matrix} y_1 & y_2 \end{matrix} \\ \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix} & \begin{pmatrix} 0.5 & 0.1 \\ 0.2 & 0.9 \\ 0.8 & 0.6 \end{pmatrix} \end{matrix}$$

$$S = \begin{matrix} & \begin{matrix} z_1 & z_2 & z_3 \end{matrix} \\ \begin{matrix} y_1 \\ y_2 \end{matrix} & \begin{pmatrix} 0.6 & 0.4 & 0.7 \\ 0.5 & 0.8 & 0.9 \end{pmatrix} \end{matrix}$$

$$\mu_{R \circ S}(x_1, z_1) = \max\{\min\{\mu_R(x_1, y_1), \mu_S(y_1, z_1)\}, \min\{\mu_R(x_1, y_2), \mu_S(y_2, z_1)\}\}$$

Composition of fuzzy relations: Example

$$X = (x_1, x_2, x_3), Y = (y_1, y_2), Z = (z_1, z_2, z_3)$$

$$R = \begin{matrix} & \begin{matrix} y_1 & y_2 \end{matrix} \\ \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix} & \begin{pmatrix} 0.5 & 0.1 \\ 0.2 & 0.9 \\ 0.8 & 0.6 \end{pmatrix} \end{matrix}$$

$$S = \begin{matrix} & \begin{matrix} z_1 & z_2 & z_3 \end{matrix} \\ \begin{matrix} y_1 \\ y_2 \end{matrix} & \begin{pmatrix} 0.6 & 0.4 & 0.7 \\ 0.5 & 0.8 & 0.9 \end{pmatrix} \end{matrix}$$

$$\begin{aligned} \mu_{R \circ S}(x_1, z_1) &= \max\{\min\{\mu_R(x_1, y_1), \mu_S(y_1, z_1)\}, \min\{\mu_R(x_1, y_2), \mu_S(y_2, z_1)\}\} \\ &= \max\{\min\{0.5, 0.6\}, \min\{0.1, 0.5\}\} \end{aligned}$$

Composition of fuzzy relations: Example

$$X = (x_1, x_2, x_3), Y = (y_1, y_2), Z = (z_1, z_2, z_3)$$

$$R = \begin{matrix} & \begin{matrix} y_1 & y_2 \end{matrix} \\ \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix} & \begin{pmatrix} 0.5 & 0.1 \\ 0.2 & 0.9 \\ 0.8 & 0.6 \end{pmatrix} \end{matrix}$$

$$S = \begin{matrix} & \begin{matrix} z_1 & z_2 & z_3 \end{matrix} \\ \begin{matrix} y_1 \\ y_2 \end{matrix} & \begin{pmatrix} 0.6 & 0.4 & 0.7 \\ 0.5 & 0.8 & 0.9 \end{pmatrix} \end{matrix}$$

$$\begin{aligned} \mu_{R \circ S}(x_1, z_1) &= \max\{\min\{\mu_R(x_1, y_1), \mu_S(y_1, z_1)\}, \min\{\mu_R(x_1, y_2), \mu_S(y_2, z_1)\}\} \\ &= \max\{\min\{0.5, 0.6\}, \min\{0.1, 0.5\}\} \\ &= \max\{0.5, 0.1\} \end{aligned}$$

Composition of fuzzy relations: Example

$$X = (x_1, x_2, x_3), Y = (y_1, y_2), Z = (z_1, z_2, z_3)$$

$$R = \begin{matrix} & \begin{matrix} y_1 & y_2 \end{matrix} \\ \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix} & \begin{pmatrix} 0.5 & 0.1 \\ 0.2 & 0.9 \\ 0.8 & 0.6 \end{pmatrix} \end{matrix}$$

$$S = \begin{matrix} & \begin{matrix} z_1 & z_2 & z_3 \end{matrix} \\ \begin{matrix} y_1 \\ y_2 \end{matrix} & \begin{pmatrix} 0.6 & 0.4 & 0.7 \\ 0.5 & 0.8 & 0.9 \end{pmatrix} \end{matrix}$$

$$\begin{aligned} \mu_{R \circ S}(x_1, z_1) &= \max\{\min\{\mu_R(x_1, y_1), \mu_S(y_1, z_1)\}, \min\{\mu_R(x_1, y_2), \mu_S(y_2, z_1)\}\} \\ &= \max\{\min\{0.5, 0.6\}, \min\{0.1, 0.5\}\} \\ &= \max\{0.5, 0.1\} \\ &= 0.5 \end{aligned}$$

Composition of fuzzy relations: Example

$$X = (x_1, x_2, x_3), Y = (y_1, y_2), Z = (z_1, z_2, z_3)$$

$$R = \begin{matrix} & y_1 & y_2 \\ \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix} & \begin{pmatrix} 0.5 & 0.1 \\ 0.2 & 0.9 \\ 0.8 & 0.6 \end{pmatrix} \end{matrix}$$

$$S = \begin{matrix} & z_1 & z_2 & z_3 \\ \begin{matrix} y_1 \\ y_2 \end{matrix} & \begin{pmatrix} 0.6 & 0.4 & 0.7 \\ 0.5 & 0.8 & 0.9 \end{pmatrix} \end{matrix}$$

$$R \circ S = \begin{matrix} & z_1 & z_2 & z_3 \\ \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix} & \begin{pmatrix} 0.5 & 0.4 & 0.5 \\ 0.5 & 0.8 & 0.9 \\ 0.6 & 0.6 & 0.7 \end{pmatrix} \end{matrix}$$

References

- 1 <https://cse.iitkgp.ac.in/~dsamanta/courses/sca/resources/slides/FL-01%20Introduction.pdf>
- 2 <https://cse.iitkgp.ac.in/~dsamanta/courses/sca/resources/slides/FL-02%20Fuzzy%20Rules.pdf>
- 3 http://webhome.csc.uvic.ca/~mcheng/460/notes/fuzzy_logic.pdf
- 4 Fuzzy Sets, Fuzzy Logic, Fuzzy Methods with Applications, John Wiley & Sons, ISBN: 0-471-95636-8,
https://www.researchgate.net/publication/260990913_Fuzzy_Sets_Fuzzy_Logic_Fuzzy_Methods_with_Applications
- 5 https://en.wikipedia.org/wiki/Fuzzy_set

Image Sources

- 1 https://www.researchgate.net/figure/Traditional-crisp-set-and-fuzzy-set-membership-functions_fig2_224605202
- 2 <https://www.iitk.ac.in/eeold/archive/courses/2013/intel-info/dlpdf3.pdf>
- 3 <https://cse.iitkgp.ac.in/~dsamanta/courses/sca/resources/slides/FL-01%20Introduction.pdf>
- 4 http://web.cecs.pdx.edu/~mperkows/CLASS_479/LECTURES479/FL001.PDF
- 5 http://webhome.csc.uvic.ca/~mcheng/460/notes/fuzzy_logic.pdf
- 6 <http://osp.mans.edu.eg/elbeltagi/AI%20FuzzyRelations.pdf>