

# 3. Multiple linear regression and Gradient descent

**Shabana K M**

PhD Research Scholar

Computer Science and Engineering

IIT Palakkad

14 August 2021



INDIAN INSTITUTE  
OF TECHNOLOGY  
**PALAKKAD**

# Recap

# Recap

- Regression

# Recap

- Regression
  - dependent, independent variables

# Recap

- Regression
  - dependent, independent variables
  - loss functions

# Recap

- Regression
  - dependent, independent variables
  - loss functions
- Different regression models

# Recap

- Regression
  - dependent, independent variables
  - loss functions
- Different regression models
  - simple, multiple

# Recap

- Regression
  - dependent, independent variables
  - loss functions
- Different regression models
  - simple, multiple
  - linear, polynomial, non-linear, logistic
- Linear regression



# Recap

- Regression
  - dependent, independent variables
  - loss functions
- Different regression models
  - simple, multiple
  - linear, polynomial, non-linear, logistic
- Linear regression
  - intercept, regression coefficients(weights)

# Recap

- Regression
  - dependent, independent variables
  - loss functions
- Different regression models
  - simple, multiple
  - linear, polynomial, non-linear, logistic
- Linear regression
  - intercept, regression coefficients(weights)
- Simple linear regression


# Recap

- Regression
  - dependent, independent variables
  - loss functions
- Different regression models
  - simple, multiple
  - linear, polynomial, non-linear, logistic
- Linear regression
  - intercept, regression coefficients(weights)
- Simple linear regression
  - Ordinary least squares

# Recap


- Regression
  - dependent, independent variables
  - loss functions
- Different regression models
  - simple, multiple
  - linear, polynomial, non-linear, logistic
- Linear regression
  - intercept, regression coefficients(weights)
- Simple linear regression
  - Ordinary least squares
  - Interpreting the regression coefficients

# Revisiting simple linear regression



(a) A simple linear regression model

# Revisiting simple linear regression




(a) A simple linear regression model




(b) Training dataset


# Revisiting simple linear regression



(a) A simple linear regression model




(b) Training dataset




(c) Fitting a linear regression model


# Revisiting simple linear regression




(a) A simple linear regression model



(b) Training dataset



(c) Fitting a linear regression model



(d) Loss function for the model



# Multiple regression

- more than one independent variable:  $x_1, x_2, \dots, x_D$

# Multiple regression

- more than one independent variable:  $x_1, x_2, \dots, x_D$
- the model is of the form

$$\hat{y} = f(w, x) = w_0 + w_1x_1 + \dots + w_Dx_D$$

# Multiple regression

- more than one independent variable:  $x_1, x_2, \dots, x_D$
- the model is of the form

$$\hat{y} = f(w, x) = w_0 + w_1x_1 + \dots + w_Dx_D$$

- when  $D = 2$ , the model is of the form  $\hat{y} = w_0 + w_1x_1 + w_2x_2$ 
  - describes a plane in the three dimensional space of  $\hat{y}, x_1$  and  $x_2$  with  $w_0$  as the intercept of the plane

# Multiple regression

- more than one independent variable:  $x_1, x_2, \dots, x_D$
- the model is of the form

$$\hat{y} = f(w, x) = w_0 + w_1x_1 + \dots + w_Dx_D$$

- when  $D = 2$ , the model is of the form  $\hat{y} = w_0 + w_1x_1 + w_2x_2$ 
  - describes a plane in the three dimensional space of  $\hat{y}, x_1$  and  $x_2$  with  $w_0$  as the intercept of the plane
- regression coefficient  $w_i$  measures the association between the predictor variable  $x_i$  and the outcome  $y$


# Multiple regression

- more than one independent variable:  $x_1, x_2, \dots, x_D$
- the model is of the form

$$\hat{y} = f(w, x) = w_0 + w_1x_1 + \dots + w_Dx_D$$

- when  $D = 2$ , the model is of the form  $\hat{y} = w_0 + w_1x_1 + w_2x_2$ 
  - describes a plane in the three dimensional space of  $\hat{y}, x_1$  and  $x_2$  with  $w_0$  as the intercept of the plane
- regression coefficient  $w_i$  measures the association between the predictor variable  $x_i$  and the outcome  $y$ 
  - $w_i$  represents the mean change in  $y$  corresponding to a unit increase in  $x_i$  when all other predictors are held fixed

# Multiple regression



**Figure:** Visualizing the multiple regression model for predicting the response variable Income ( $y$ ) based on the independent variables Seniority ( $x_1$ ) and Years of Education ( $x_2$ )

# Problem definition

**Given:** Training data set comprising  $N$  observations  $(x_n, y_n)_{n=1}^N$ , where  $x_n = [x_{n1}, x_{n2}, \dots, x_{nD}]$  is the input and  $y_n$  is the corresponding output

# Problem definition

**Given:** Training data set comprising  $N$  observations  $(x_n, y_n)_{n=1}^N$ , where  $x_n = [x_{n1}, x_{n2}, \dots, x_{nD}]$  is the input and  $y_n$  is the corresponding output

**Goal:** Predict the  $y$  value for a new value of  $x$



# Problem definition

**Given:** Training data set comprising  $N$  observations  $(x_n, y_n)_{n=1}^N$ , where  $x_n = [x_{n1}, x_{n2}, \dots, x_{nD}]$  is the input and  $y_n$  is the corresponding output

**Goal:** Predict the  $y$  value for a new value of  $x$

**Estimate:** The weights  $w = [w_0, w_1, \dots, w_D]$

# Problem definition

**Given:** Training data set comprising  $N$  observations  $(x_n, y_n)_{n=1}^N$ , where  $x_n = [x_{n1}, x_{n2}, \dots, x_{nD}]$  is the input and  $y_n$  is the corresponding output

**Goal:** Predict the  $y$  value for a new value of  $x$

**Estimate:** The weights  $w = [w_0, w_1, \dots, w_D]$

**Minimize:** Mean-squared error:  $l(w) = \frac{1}{N} \sum_{i=1}^N (y_i - \hat{y}_i)^2$   
where  $\hat{y}_i = f(w, x_i) = w_0 + w_1 x_{i1} + \dots + w_D x_{iD}$

# A small mathematics refresher

**Scalar:**

24

**Vector:**

[ 2, -6, 9 ]

row

or  
column

$\begin{bmatrix} 2, \\ 6, \\ 9 \end{bmatrix}$

**Matrix:**

$\begin{bmatrix} 2, & -6, & 9 \\ 4, & 5, & -7 \end{bmatrix}$

row(s) x column(s)

**Scalar:** a single number

**Vector:** an ordered array of numbers  
can be in a row or a column  
an index points to a specific value within the vector

**Matrix:** two dimensional array of numbers  
each element identified by two numbers

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

# Multiple regression

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation

- $x_{13} = 75$



# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation

- $x_{13} = 75$
- $x_{42} =$

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation

- $x_{13} = 75$

- $x_{42} = 98$

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation

- $x_{13} = 75$

- $x_{42} = 98$

- $y_i$  -

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation
  - $x_{13} = 75$
  - $x_{42} = 98$
- $y_i$  -  $y$  value of the  $i^{th}$  observation

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation
  - $x_{13} = 75$
  - $x_{42} = 98$
- $y_i$  -  $y$  value of the  $i^{th}$  observation
  - $y_1 =$

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation
  - $x_{13} = 75$
  - $x_{42} = 98$
- $y_i$  -  $y$  value of the  $i^{th}$  observation
  - $y_1 = 152$
  - $y_4 =$

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Consider the problem of predicting the Final exam score ( $y$ ) based on the scores obtained in the first three exams ( $x_1, x_2, x_3$ )

- Using a linear regression model for this problem, we have:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

- $x_{ij}$  -  $j^{th}$  feature of the  $i^{th}$  observation
  - $x_{13} = 75$
  - $x_{42} = 98$
- $y_i$  -  $y$  value of the  $i^{th}$  observation
  - $y_1 = 152$
  - $y_4 = 196$

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.



# Multiple regression

Our model:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

# Multiple regression

Our model:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

We can write:

$$y = w_0 + w_1x_1 + w_2x_2 + w_3x_3 + e$$

where  $e = y - \hat{y}$

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

# Multiple regression

EXAM1	EXAM2	EXAM3	FINAL
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142
53	46	55	101
69	74	77	149
47	56	60	115
87	79	90	175
79	70	88	164
69	70	73	141
70	65	74	141
93	95	91	184
79	80	73	152
70	73	78	148
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Our model:

$$\hat{y} = w_0 + w_1x_1 + w_2x_2 + w_3x_3$$

We can write:

$$y = w_0 + w_1x_1 + w_2x_2 + w_3x_3 + e$$

where  $e = y - \hat{y}$

Now we have:

$$y_1 = w_0 + w_1x_{11} + w_2x_{12} + w_3x_{13} + e_1$$

$$y_2 = w_0 + w_1x_{21} + w_2x_{22} + w_3x_{23} + e_2$$

$$\vdots$$

$$y_N = w_0 + w_1x_{N1} + w_2x_{N2} + w_3x_{N3} + e_N$$

# Matrix Representation

$$y_1 = w_0 + w_1x_{11} + w_2x_{12} + w_3x_{13} + e_1$$

$$y_2 = w_0 + w_1x_{21} + w_2x_{22} + w_3x_{23} + e_2$$

$$\vdots$$

$$y_N = w_0 + w_1x_{N1} + w_2x_{N2} + w_3x_{N3} + e_N$$

# Matrix Representation

$$y_1 = w_0 + w_1x_{11} + w_2x_{12} + w_3x_{13} + e_1$$

$$y_2 = w_0 + w_1x_{21} + w_2x_{22} + w_3x_{23} + e_2$$

$$\vdots$$

$$y_N = w_0 + w_1x_{N1} + w_2x_{N2} + w_3x_{N3} + e_N$$

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} X = \begin{bmatrix} 1 & x_{11} & x_{12} & x_{13} \\ 1 & x_{21} & x_{22} & x_{23} \\ \vdots & & & \\ 1 & x_{N1} & x_{N2} & x_{N3} \end{bmatrix} w = \begin{bmatrix} w_0 \\ w_1 \\ w_2 \\ w_3 \end{bmatrix} e = \begin{bmatrix} e_1 \\ e_2 \\ \vdots \\ e_N \end{bmatrix}$$

# Matrix Representation

$$y_1 = w_0 + w_1x_{11} + w_2x_{12} + w_3x_{13} + e_1$$

$$y_2 = w_0 + w_1x_{21} + w_2x_{22} + w_3x_{23} + e_2$$

$$\vdots$$

$$y_N = w_0 + w_1x_{N1} + w_2x_{N2} + w_3x_{N3} + e_N$$

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} \quad X = \begin{bmatrix} 1 & x_{11} & x_{12} & x_{13} \\ 1 & x_{21} & x_{22} & x_{23} \\ \vdots & & & \\ 1 & x_{N1} & x_{N2} & x_{N3} \end{bmatrix} \quad w = \begin{bmatrix} w_0 \\ w_1 \\ w_2 \\ w_3 \end{bmatrix} \quad e = \begin{bmatrix} e_1 \\ e_2 \\ \vdots \\ e_N \end{bmatrix}$$

The  $N$  equations can be written as:

$$y = Xw + e$$

# Mean Square Error (MSE)

$$y = Xw + e$$

# Mean Square Error (MSE)

$$y = Xw + e$$

Rearranging the terms, we get

$$e = y - Xw$$



# Mean Square Error (MSE)

$$y = Xw + e$$

Rearranging the terms, we get

$$e = y - Xw$$

$$MSE = \frac{\sum_i^N e_i^2}{N}$$

# Mean Square Error (MSE)

$$y = Xw + e$$

Rearranging the terms, we get

$$e = y - Xw$$

$$MSE = \frac{\sum_i^N e_i^2}{N}$$

$$\sum_i^N e^2 = e_1^2 + e_2^2 + \dots + e_N^2$$

$$= \begin{bmatrix} e_1 & e_2 & \dots & e_N \end{bmatrix} \begin{bmatrix} e_1 \\ e_2 \\ \vdots \\ e_N \end{bmatrix} = e^T e = (y - Xw)^T (y - Xw)$$

# Computing $w$ using Ordinary Least Squares (OLS)

$$MSE = \frac{\sum_i^N e_i^2}{N} = \frac{(y - Xw)^T (y - Xw)}{N}$$

# Computing $w$ using Ordinary Least Squares (OLS)

$$\begin{aligned}MSE &= \frac{\sum_i^N e_i^2}{N} = \frac{(y - Xw)^T (y - Xw)}{N} \\&= \frac{1}{N} (y^T y - y^T Xw - w^T X^T y + w^T X^T Xw)\end{aligned}$$

# Computing $w$ using Ordinary Least Squares (OLS)

$$MSE = \frac{\sum_i^N e_i^2}{N} = \frac{(y - Xw)^T (y - Xw)}{N}$$

$$= \frac{1}{N} (y^T y - y^T Xw - w^T X^T y + w^T X^T Xw)$$

Now  $y^T Xw = (w^T X^T y)^T$  is a scalar, so we have

# Computing $w$ using Ordinary Least Squares (OLS)

$$MSE = \frac{\sum_i^N e_i^2}{N} = \frac{(y - Xw)^T (y - Xw)}{N}$$

$$= \frac{1}{N} (y^T y - y^T Xw - w^T X^T y + w^T X^T Xw)$$

Now  $y^T Xw = (w^T X^T y)^T$  is a scalar, so we have

$$MSE = \frac{1}{N} (y^T y - 2w^T X^T y + w^T X^T Xw)$$

# Computing $w$ using Ordinary Least Squares (OLS)

$$\begin{aligned}MSE &= \frac{\sum_i^N e_i^2}{N} = \frac{(y - Xw)^T (y - Xw)}{N} \\&= \frac{1}{N} (y^T y - y^T Xw - w^T X^T y + w^T X^T Xw)\end{aligned}$$

Now  $y^T Xw = (w^T X^T y)^T$  is a scalar, so we have

$$MSE = \frac{1}{N} (y^T y - 2w^T X^T y + w^T X^T Xw)$$

Using OLS, the regression coefficients  $w$  are estimated by solving the following minimization problem:

$$\min_w \frac{1}{N} (y^T y - 2w^T X^T y + w^T X^T Xw)$$

# Computing $w$ using Ordinary Least Squares (OLS)

$$l(w) = \frac{1}{N}(y^T y - 2w^T X^T y + w^T X^T X w)$$



# Computing $w$ using Ordinary Least Squares (OLS)

$$l(w) = \frac{1}{N}(y^T y - 2w^T X^T y + w^T X^T X w)$$

The error is minimized for the value of  $w$  such that  $\frac{\partial l(w)}{\partial w} = 0$

# Computing $w$ using Ordinary Least Squares (OLS)

$$l(w) = \frac{1}{N}(y^T y - 2w^T X^T y + w^T X^T X w)$$

The error is minimized for the value of  $w$  such that  $\frac{\partial l(w)}{\partial w} = 0$

$$-2 * X^T y + 2 * X^T X w = 0$$

# Computing $w$ using Ordinary Least Squares (OLS)

$$l(w) = \frac{1}{N}(y^T y - 2w^T X^T y + w^T X^T X w)$$

The error is minimized for the value of  $w$  such that  $\frac{\partial l(w)}{\partial w} = 0$

$$-2 * X^T y + 2 * X^T X w = 0$$

$$w = (X^T X)^{-1}(X^T y)$$

# Computing $w$ using Ordinary Least Squares (OLS)

$$l(w) = \frac{1}{N}(y^T y - 2w^T X^T y + w^T X^T X w)$$

The error is minimized for the value of  $w$  such that  $\frac{\partial l(w)}{\partial w} = 0$

$$-2 * X^T y + 2 * X^T X w = 0$$

$$w = (X^T X)^{-1}(X^T y)$$

## Normal equation

■  $w = (X^T X)^{-1}(X^T y)$

# Normal equation

$$w = (X^T X)^{-1} (X^T y)$$



- gives the solution in one go

# Normal equation

$$w = (X^T X)^{-1} (X^T y)$$



- gives the solution in one go
- works well when using small feature sets

# Normal equation

$$w = (X^T X)^{-1} (X^T y)$$



- gives the solution in one go
- works well when using small feature sets



- numerical complexity
  - need to compute  $(X^T X)^{-1}$  - can be very slow

# Normal equation

$$w = (X^T X)^{-1} (X^T y)$$



- gives the solution in one go
- works well when using small feature sets



- numerical complexity
  - need to compute  $(X^T X)^{-1}$  - can be very slow
- $X^T X$  may not be always invertible



# Normal equation

$$w = (X^T X)^{-1} (X^T y)$$



- gives the solution in one go
- works well when using small feature sets



- numerical complexity
  - need to compute  $(X^T X)^{-1}$  - can be very slow
- $X^T X$  may not be always invertible
  - too many features (eg:-  $N < D$ )

# Normal equation

$$w = (X^T X)^{-1} (X^T y)$$



- gives the solution in one go
- works well when using small feature sets



- numerical complexity
  - need to compute  $(X^T X)^{-1}$  - can be very slow
- $X^T X$  may not be always invertible
  - too many features (eg:-  $N < D$ )
  - some columns are linearly dependent (redundant features)

# Normal equation

$$w = (X^T X)^{-1} (X^T y)$$



- gives the solution in one go
- works well when using small feature sets




- numerical complexity
  - need to compute  $(X^T X)^{-1}$  - can be very slow
- $X^T X$  may not be always invertible
  - too many features (eg:-  $N < D$ )
  - some columns are linearly dependent (redundant features)
- cannot be easily parallelized

# Loss function for simple linear regression

Consider a simple linear regression problem. If we compute the cost/error function for all possible values of  $w_1$ , the resulting plot will always be convex, or kind of bowl-shaped



# Loss function for simple linear regression

Consider a simple linear regression problem. If we compute the cost/error function for all possible values of  $w_1$ , the resulting plot will always be convex, or kind of bowl-shaped





# Loss function for simple linear regression

Consider a simple linear regression problem. If we compute the cost/error function for all possible values of  $w_1$ , the resulting plot will always be convex, or kind of bowl-shaped



# Loss function for simple linear regression



Consider a simple linear regression problem. If we compute the cost/error function for all possible values of  $w_1$ , the resulting plot will always be convex, or kind of bowl-shaped



- Convex problems have only one minimum; i.e., only one place where the slope is exactly 0

# Loss function for simple linear regression

Consider a simple linear regression problem. If we compute the cost/error function for all possible values of  $w_1$ , the resulting plot will always be convex, or kind of bowl-shaped





- Convex problems have only one minimum; i.e., only one place where the slope is exactly 0 - **convergence point**



## Loss function for simple linear regression



Consider a simple linear regression problem. If we compute the cost/error function for all possible values of  $w_1$ , the resulting plot will always be convex, or kind of bowl-shaped



- Convex problems have only one minimum; i.e., only one place where the slope is exactly 0 - **convergence point**
- Compute the loss function for all possible values of  $w_1$  over the entire data set and then find the minimum

# Loss function for simple linear regression

Consider a simple linear regression problem. If we compute the cost/error function for all possible values of  $w_1$ , the resulting plot will always be convex, or kind of bowl-shaped



- Convex problems have only one minimum; i.e., only one place where the slope is exactly 0 - **convergence point**
- Compute the loss function for all possible values of  $w_1$  over the entire data set and then find the minimum - **inefficient!**

# Gradient Descent

## Gradient descent

- iterative optimization algorithm to find the minimum of a function

# Gradient Descent

## Gradient descent

- iterative optimization algorithm to find the minimum of a function  
loss function in our case!!

# Gradient Descent


## Gradient descent

- iterative optimization algorithm to find the minimum of a function  
loss function in our case!!
- idea: follow the gradients of the loss function

# Gradient Descent

## Gradient descent

- iterative optimization algorithm to find the minimum of a function  
loss function in our case!!
- idea: follow the gradients of the loss function



# Gradient

- used for functions with several inputs and a single output

# Gradient

- used for functions with several inputs and a single output
- vector of partial derivatives with respect to each of its inputs



# Gradient

- used for functions with several inputs and a single output
- vector of partial derivatives with respect to each of its inputs
- points in the direction of greatest increase of a function

# Gradient

- used for functions with several inputs and a single output
- vector of partial derivatives with respect to each of its inputs
- points in the direction of greatest increase of a function
- is zero at a local maximum or local minimum

# Gradient

- used for functions with several inputs and a single output
- vector of partial derivatives with respect to each of its inputs
- points in the direction of greatest increase of a function
- is zero at a local maximum or local minimum
- $\nabla f = \left[ \frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial y} \quad \dots \right]^T$

# Gradient

- used for functions with several inputs and a single output
- vector of partial derivatives with respect to each of its inputs
- points in the direction of greatest increase of a function
- is zero at a local maximum or local minimum
- $\nabla f = \left[ \frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial y} \quad \dots \right]^T$

Let  $f(x, y) = ax^2 + by^2$ ,

then  $\nabla f =$

# Gradient

- used for functions with several inputs and a single output
- vector of partial derivatives with respect to each of its inputs
- points in the direction of greatest increase of a function
- is zero at a local maximum or local minimum
- $\nabla f = \left[ \frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial y} \quad \dots \right]^T$

Let  $f(x, y) = ax^2 + by^2$ ,

then  $\nabla f = \begin{bmatrix} 2ax \\ 2by \end{bmatrix}$


# Gradient

- used for functions with several inputs and a single output
- vector of partial derivatives with respect to each of its inputs
- points in the direction of greatest increase of a function
- is zero at a local maximum or local minimum

$$\nabla f = \left[ \frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial y} \quad \dots \right]^T$$

Let  $f(x, y) = ax^2 + by^2$ ,

$$\text{then } \nabla f = \begin{bmatrix} 2ax \\ 2by \end{bmatrix}$$



# Gradient Descent

**AIM:** Find the value of  $w_1$  corresponding to the minimum value of the loss function

# Gradient Descent

**AIM:** Find the value of  $w_1$  corresponding to the minimum value of the loss function

**BASIC IDEA:** Start with an initial parameter value of  $w_1$  and iteratively move towards a new value of  $w_1$  that minimizes the loss function



# Gradient Descent

**AIM:** Find the value of  $w_1$  corresponding to the minimum value of the loss function

**BASIC IDEA:** Start with an initial parameter value of  $w_1$  and iteratively move towards a new value of  $w_1$  that minimizes the loss function


- can set  $w_1$  to 0 or some random value

# Gradient Descent

**AIM:** Find the value of  $w_1$  corresponding to the minimum value of the loss function

**BASIC IDEA:** Start with an initial parameter value of  $w_1$  and iteratively move towards a new value of  $w_1$  that minimizes the loss function

- can set  $w_1$  to 0 or some random value



# Gradient Descent: Determining the next value of $w_1$

1. Calculate the gradient of the loss curve at the starting point

# Gradient Descent: Determining the next value of $w_1$

## 1. Calculate the gradient of the loss curve at the starting point

- the gradient  $\nabla f$  points in the direction of the greatest rate of increase of the function

# Gradient Descent: Determining the next value of $w_1$

## 1. Calculate the gradient of the loss curve at the starting point

- the gradient  $\nabla f$  points in the direction of the greatest rate of increase of the function
  - its magnitude is the slope of the graph in that direction

# Gradient Descent: Determining the next value of $w_1$

## 1. Calculate the gradient of the loss curve at the starting point


- the gradient  $\nabla f$  points in the direction of the greatest rate of increase of the function
  - its magnitude is the slope of the graph in that direction
- the negative of the gradient ( $-\nabla f$ ) points in the direction of maximum decrease in height

# Gradient Descent: Determining the next value of $w_1$

## 1. Calculate the gradient of the loss curve at the starting point



- the gradient  $\nabla f$  points in the direction of the greatest rate of increase of the function
  - its magnitude is the slope of the graph in that direction
- the negative of the gradient ( $-\nabla f$ ) points in the direction of maximum decrease in height
- gradient descent algorithm takes a step in the direction of the negative gradient in order to reduce loss as quickly as possible

# Gradient Descent





# Gradient Descent



# Gradient Descent: Determining the next value

**2. Add some fraction of the gradient's magnitude to the starting point**

# Gradient Descent: Determining the next value

**2. Add some fraction of the gradient's magnitude to the starting point**

$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

# Gradient Descent: Determining the next value

**2. Add some fraction of the gradient's magnitude to the starting point**

$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

This process is repeated until convergence

# Gradient Descent: Determining the next value

**2. Add some fraction of the gradient's magnitude to the starting point**

$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

This process is repeated until convergence

$\alpha$  (learning rate) - controls the step size

# Gradient Descent: Determining the next value

## 2. Add some fraction of the gradient's magnitude to the starting point

$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

This process is repeated until convergence

$\alpha$  (learning rate) - controls the step size

- high value - might step over the minimum

# Gradient Descent: Determining the next value

## 2. Add some fraction of the gradient's magnitude to the starting point

$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

This process is repeated until convergence

$\alpha$  (learning rate) - controls the step size

- high value - might step over the minimum
- low value - slow convergence

# Gradient Descent: Determining the next value


## 2. Add some fraction of the gradient's magnitude to the starting point

$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

This process is repeated until convergence

$\alpha$  (learning rate) - controls the step size

- high value - might step over the minimum
- low value - slow convergence





# Gradient Descent: Determining the next value


## 2. Add some fraction of the gradient's magnitude to the starting point

$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$


This process is repeated until convergence

$\alpha$  (learning rate) - controls the step size

- high value - might step over the minimum
- low value - slow convergence



# Choosing the learning rate



# Gradient Descent for linear regression

**Loss function:** 
$$l(w) = \frac{1}{N} \sum_{i=1}^N \underbrace{(w_0 + w_1 x_{i1} + \dots + w_D x_{iD})}_{\hat{y}_i} - y_i)^2$$

# Gradient Descent for linear regression

**Loss function:** 
$$l(w) = \frac{1}{N} \sum_{i=1}^N \underbrace{(w_0 + w_1 x_{i1} + \dots + w_D x_{iD})}_{\hat{y}_i} - y_i)^2$$

**Gradient descent update:** 
$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

# Gradient Descent for linear regression

**Loss function:** 
$$l(w) = \frac{1}{N} \sum_{i=1}^N \underbrace{(w_0 + w_1 x_{i1} + \dots + w_D x_{iD})}_{\hat{y}_i} - y_i)^2$$

**Gradient descent update:** 
$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

**Updates:**

# Gradient Descent for linear regression

**Loss function:** 
$$l(w) = \frac{1}{N} \sum_{i=1}^N \underbrace{(w_0 + w_1 x_{i1} + \dots + w_D x_{iD})}_{\hat{y}_i} - y_i)^2$$

**Gradient descent update:** 
$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

**Updates:**

$$w_0 := w_0 - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i)$$

# Gradient Descent for linear regression

**Loss function:** 
$$l(w) = \frac{1}{N} \sum_{i=1}^N \underbrace{((w_0 + w_1 x_{i1} + \dots + w_D x_{iD}) - y_i)}_{\hat{y}_i}^2$$

**Gradient descent update:** 
$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

**Updates:**

$$w_0 := w_0 - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i)$$

$$w_1 := w_1 - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i) * x_{i1}$$

# Gradient Descent for linear regression

**Loss function:** 
$$l(w) = \frac{1}{N} \sum_{i=1}^N \underbrace{((w_0 + w_1 x_{i1} + \dots + w_D x_{iD}) - y_i)^2}_{\hat{y}_i}$$

**Gradient descent update:** 
$$w_i^{next} = w_i^{old} - \alpha * \frac{\partial l(w)}{\partial w_i}$$

**Updates:**

$$w_0 := w_0 - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i)$$

$$w_1 := w_1 - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i) * x_{i1}$$

$$\vdots$$

$$w_D := w_D - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i) * x_{iD}$$



# Gradient Descent for linear regression

repeat until convergence {

$$w_j = w_j - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i) * x_{ij} \quad \text{for } j := 0 \dots D \quad \}$$

# Gradient Descent for linear regression

repeat until convergence {

$$w_j = w_j - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i) * x_{ij} \quad \text{for } j := 0 \dots D \quad \}$$

**Matrix notation:**  $w := w - \alpha \nabla I(w)$

# Gradient Descent for linear regression

repeat until convergence {

$$w_j = w_j - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i) * x_{ij} \quad \text{for } j := 0 \dots D \}$$

**Matrix notation:**  $w := w - \alpha \nabla I(w)$

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \quad X = \begin{bmatrix} 1 & x_{11} & x_{12} & \cdots & x_{1D} \\ 1 & x_{21} & x_{22} & \cdots & x_{2D} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{N1} & x_{N2} & \cdots & x_{ND} \end{bmatrix} \quad w = \begin{bmatrix} w_0 \\ w_1 \\ w_2 \\ \vdots \\ w_D \end{bmatrix}$$

# Gradient Descent for linear regression

repeat until convergence {

$$w_j = w_j - \alpha * \frac{2}{N} \sum_{i=1}^N (\hat{y}_i - y_i) * x_{ij} \quad \text{for } j := 0 \dots D \}$$


**Matrix notation:**  $w := w - \alpha \nabla I(w)$

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \quad X = \begin{bmatrix} 1 & x_{11} & x_{12} & \cdots & x_{1D} \\ 1 & x_{21} & x_{22} & \cdots & x_{2D} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{N1} & x_{N2} & \cdots & x_{ND} \end{bmatrix} \quad w = \begin{bmatrix} w_0 \\ w_1 \\ w_2 \\ \vdots \\ w_D \end{bmatrix}$$

The matrix notation of the Gradient Descent rule is:


$$w := w - \frac{2\alpha}{N} X^T (Xw - y)$$

# Gradient Descent: Demo




(a) Initial  $w$

# Gradient Descent: Demo




(a) Initial  $w$




(b) After one iteration

# Gradient Descent: Demo




(a) After 2 iterations

# Gradient Descent: Demo



(a) After 2 iterations



(b) After 3 iterations



# References

- 1 <https://towardsdatascience.com/multiple-linear-regression-with-math-and-code-c1052f3c7446>
- 2 <https://developers.google.com/machine-learning/crash-course/reducing-loss/gradient-descent>
- 3 <https://www.coursera.org/learn/machine-learning/resources/QQx8l>

Thanks Google for the pictures!