

Propagation of UHECRs in simulated Extra Galactic Magnetic Fields

Stefan Hackstein

University of Hamburg

shackste@physnet.uni-hamburg.de

12.04.2017

Overview

Basics

Cosmic Rays

Extragalactic Magnetic Fields

Simulations

Magnetic Field Models

CRPropa

Sources

Results

Energy Spectrum

Composition

Angular Power

Source Contribution

Conclusion

charged Nuclei

gyro radius $r_g = E/eZB$


low energy $E < 10^{18}$ eV

galactic origin

sources: remnants of SNe

Cosmic Rays

Cosmic Ray Spectra of Various Experiments



Cosmic Rays

charged Nuclei

$$\text{gyro radius } r_g = E/eZB$$

$$\text{low energy } E < 10^{18} \text{ eV}$$

galactic origin


sources: remnants of SNe


$$\text{high energy } E > 10^{18} \text{ eV}$$

extragalactic origin

sources: *unknown*

Cosmic Ray Spectra of Various Experiments





Voids ($\approx 80\%$ of volume)

$$\begin{aligned}B_0 &\leq 0.55 - 5.6 \text{ nG} \\B_{\text{void}} &\geq 10^{-16} \text{ G}\end{aligned}$$


LSS ($\approx 20\%$ of volume)

galaxies $\sim 5 - 15 \mu\text{G}$

clusters $\sim \mu\text{G}$

outskirts and filaments

unknown



Extra-Galactic Magnetic Fields measure EGMFs with UHECRs? constrain seeding processes?

Voids ($\approx 80\%$ of volume)

$$B_0 \leq 0.55 - 5.6 \text{ nG}$$

$$B_{\text{void}} \geq 10^{-16} \text{ G}$$

huge range of uncertainty

LSS ($\approx 20\%$ of volume)

galaxies $\sim 5 - 15 \mu\text{G}$

clusters $\sim \mu\text{G}$

outskirts and filaments

unknown

Combine ENZO

(large MHD simulations on cosmic scale)

with CRPropa

(propagation of UHECRs in 3D models of EGMF)

Magnetic Field Models


ENZO (cosmological grid MHD solver)

unconstrained

$$B_0 = 10^{-14} \text{ G} - 10^{-8} \text{ G}$$

18 observers

Hackstein et al. 2016 MNRAS




constrained (*Sorce 2015*)

$$\text{primordial } B_0 = 10^{-9} \text{ G}$$

astrophysical

magnetic feedback from AGN



Magnetic Field


ENZO (cosmological)

unconstrained

$$B_0 = 10^{-14} \text{ G} - 10^{-8} \text{ G}$$

18 observers


Hackstein et al. 2016 MNRAS



**Energy spectrum
not affected
by EGMFs**

**strong EGMFs
increase
Anisotropy**

**Imprint of
nearby sources
at highest energy**



Magnetic Field

ENZO (cosmological)

unconstrained

$$B_0 = 10^{-14} \text{ G} - 10^{-8} \text{ G}$$

18 observers


Hackstein et al. 2016 MNRAS

What about
- heavy nuclei?
- constrained
simulation?

Energy spectrum
not affected
by EGMFs

strong EGMFs
increase
Anisotropy

Imprint of
nearby sources
at highest energy



Magnetic Field Models

ENZO (cosmological grid MHD solver)

unconstrained

$$B_0 = 10^{-14} \text{ G} - 10^{-8} \text{ G}$$

18 observers



Hackstein et al. 2016 MNRAS

constrained (*Sorce 2015*)


$$\text{primordial } B_0 = 10^{-9} \text{ G}$$

astrophysical

magnetic feedback from AGN



CRPropa



Lorentz deflection

energy loss processes

- pair production
- pion production
(GZK-effect, $E > 4 \cdot 10^{19}$ eV,
 \Rightarrow distance limit ~ 100 Mpc)
- cosmic expansion


disintegration processes

- photo disintegration
- nuclear decay


I. Dutan, L. I. Caramete et al. 2015
<https://crpropa.desy.de/>

Sources

- **homogeneous** distribution




- follow gas **density** 1:1




- **mass halo**: viral haloes

- **vcv_velzen**: AGN & radio galaxies

(Véron-Cetty&Véron 2010, van Velzen 2012)




Energy Spectrum




- softer spectrum for heavy injection (*Allard 2012*)
due to **shorter energy loss length** & remnant protons
- low energy: spectrum universal (*Aloisio 2004*)

Energy Spectrum



- softer spectrum for heavy injection (*Allard 2012*) due to **shorter energy loss length** & remnant protons
- low energy: spectrum universal (*Aloisio 2004*)

Energy Spectrum





- softer spectrum for heavy injection (*Allard 2012*) due to **shorter energy loss length** & remnant protons
- low energy: spectrum universal (*Aloisio 2004*)
- nearby sources** determine spectrum above GZK-limit
- no significant effect of EGMFs

Composition


- not affected by choice of sources

(Globus 2007)





Composition



- not affected by choice of sources (Globus 2007)
- lighter composition in strongest models due to **longer travel distance**
- increased variance, independent of sources
 ⇒ magnetization of observer favours effect



Angular Power $C_l = \frac{1}{2l+1} \sum_{m=-l}^l a_{lm} a_{lm}^*$




$$\text{Angular Power } C_l = \frac{1}{2l+1} \sum_{m=-l}^l a_{lm} a_{lm}^*$$




- source **imprint at highest Energy** in all models (*Harari 2013*)
- constrained: **wash out** anisotropy (*Takami 2012*)
- unconstrained: EGMFs **favour directions**
- stronger deflection** of heavy nuclei

Contribution of Sources




Contribution of Sources



weak magnetic fields:

- geometric diffusion

Contribution of Sources




weak magnetic fields:


- geometric diffusion

strong magnetic fields:

- increased detection probability (*Harari 2013*)
- magnetic horizon (*Batista 2014*)



Contribution of Sources




weak magnetic fields:

- geometric diffusion

strong magnetic fields:

- increased detection probability (*Harari 2013*)
- magnetic horizon (*Batista 2014*)



Conclusion


	Sources	Strong CMFs	Heavy Nuclei
spectrum	determine E_{\max} & slope $> E_{\text{GZK}}$	x	softer spectrum
anisotropy	imprint at highest E	increase wash out	stronger deflection
observed composition	x	x	(obvious)

- UHECR astronomy possible at highest Energies ($\sim 10^{20}$ eV)
- chance to constrain magnetic seeding processes


Artefacts

effect of finite observer



artificial deflection




ideal observer



finite observer



magnetized observer



References

-  Ade P. A. R. et al.,
Planck 2015 results. XIX. Constraints on primordial magnetic fields,
arXiv:1502.01594 [astro-ph.CO] Feb 5 2015
-  Neronov A. & Vovk I.,
Evidence for strong extragalactic magnetic fields from Fermi observations
of TeV blazars, *arXiv:1006.3504 [astro-ph.HE]* Jun 17 2010
-  Ade et al.,
Cosmic magnetic fields - as observed in the Universe, in galactic dynamos,
and in the Milky Way, *2004NewAR..48..763V* 09/2004
-  Feretti L., Govoni F.,
Magnetic Field in Clusters of Galaxies, *arXiv:astro-ph/0410182* Oct 7 2004
-  S. Hackstein et al.,
Propagation of Ultra High Energy Cosmic Rays in Extragalactic Magnetic
Fields: A view from cosmological simulations *arXiv:1607.08872*
[astro-ph.CO] Jul 29 2016

References

-  [Sorce J. et al.](#)
Cosmicflows Constrained Local UniversE Simulations *arXiv:1510.04900 [astro-ph.CO]* Oct 16 2015
-  [Sigl G. et al.](#)
CRPropa <https://crpropa.desy.de/>
-  [Véron-Cetty, M.-P. & Véron, P.,](#)
A catalogue of quasars and active nuclei: 13th edition,
doi:10.1051/0004-6361/201014188 Aug 20 2010
-  [van Velzen, S.,](#)
Radio galaxies of the local universe: all-sky catalog, luminosity functions, and clustering, *arXiv:1206.0031 [astro-ph.CO]* Jul 20 2012
-  [Allard D.,](#)
Extragalactic propagation of ultrahigh energy cosmic-rays, *arXiv:1111.3290 [astro-ph.HE]* Nov 14 2011

References

-  R. Aloisio, V. Berezinsky,
Diffusive propagation of UHECR and the propagation theorem
arXiv:astro-ph/0403095 Mar 03 2004
-  N. Globus, D. Allard, E. Parizot,
Propagation of high-energy cosmic rays in extragalactic turbulent
magnetic fields: resulting energy spectrum and composition
arXiv:0709.1541 [astro-ph] Sep 11 2007
-  G. Sigl, F. Miniati, T. A. Enßlin,
Ultrahigh energy cosmic ray probes of large scale structure and magnetic
fields, *Phys. Rev. D 70, 043007 - Aug 25 2004*
-  D. Harari, S. Mollerach, E. Roulet,
Anisotropies of ultra-high energy cosmic rays diffusing from extragalactic
sources, *arXiv:1312.1366 [astro-ph.HE] Dec 04 2013*
-  H. Takami, S. Inoue, T. Yamamoto,
Propagation of ultra-high-energy cosmic ray nuclei in cosmic magnetic
fields and implications for anisotropy measurements, *Vol. 35, Issue 12, Jul
2012, p. 767780*