

12340306

Mohammad Shadik Ansari

Exercise sheet I

1. Sol"

$$A = \sum_{j=5}^{10} (3j)$$

$$= 3 \sum_{j=5}^{10} j$$

$$= 3 \sum_{j=5}^{10} j$$

$$= 3(5 + 6 + 7 + 8 + 9 + 10)$$

$$= 3 \times 45$$

$$= 135$$

$$\left[\begin{array}{l} \text{: Sum of } n \text{ square numbers} \\ \text{: Sum of } n \text{ natural numbers} \end{array} \right]$$

$$B = \sum_{i=1}^{50} (i^3 - (i-1)^3)$$

$$= \sum_{i=1}^{50} (i^3 - (i^3 - 3i^2 + 3i - 1))$$

$$= \sum_{i=1}^{50} (3i^2 - 3i + 1)$$

$$= 3 \cdot \sum_{i=1}^{50} i^2 - 3 \sum_{i=1}^{50} i + \sum_{i=1}^{50} 1$$

$$= 3 \times \left(\frac{50(50+1)(2 \times 50+1)}{6} \right) - 3 \times \left(\frac{50 \times (50+1)}{2} \right) + 50 \times 1$$

$$= 128775 - 3825 + 50$$

$$= 125,000$$

~~C~~ C is the sum of first 110 numbers in the list 3, 7, 11, 15, 19, 23, ...

$$\text{then, } a_i = 3 + (i-1) \times 4 = 3 + 4i - 4 = 4i - 1$$

$$\therefore \sum_{i=1}^{110} 4i - 1 = 4 \sum_{i=1}^{110} i - \sum_{i=1}^{110} 1$$

$$= 4 \times \left(\frac{110 \cdot (110+1)}{2} \right) - 110$$

$$= 24310$$

12340306

Mohammad Shadik Ansari

2. Solⁿ

$$A = \prod_{i=1}^5 (2i)$$

$$= (2 \times 1) \cdot (2 \times 2) \cdot (2 \times 3) \cdot (2 \times 4) \cdot (2 \times 5)$$

$$= 2^5 \times 5!$$

$$= 3840 //$$

$$B = \prod_{i=3}^5 (i-2)!$$

$$= (3-2)! \cdot (4-2)! \cdot (5-2)!$$

$$= 1! \cdot 2! \cdot 3!$$

$$= 1 \times 2 \times 6$$

$$= 12 //$$

$$C = \prod_{i=1}^3 \prod_{k=2}^4 (2i+k)$$

$$= \prod_{k=2}^4 (2 \times 1 + k) \times \prod_{k=2}^4 (2 \times 2 + k) \times \prod_{k=2}^4 (2 \times 3 + k)$$

$$= (2+2) \cdot (2+3) \cdot (2+4) \times (4+2) \cdot (4+3) \cdot (4+4) \times (6+2) \cdot (6+3) \cdot (6+4)$$

$$= 4 \cdot 5 \cdot 6 \times 6 \cdot 7 \cdot 8 \times 8 \cdot 9 \cdot 10$$

$$= 29030400 //$$

JKU

JOHANNES KEPLER
UNIVERSITÄT LINZ

12340306

Mohammed Shadik Ansari

$$3. A = \sum_{k=1}^n k \cdot k!$$

$$= \sum_{k=1}^n (k+1-1) k!$$

$$= \sum_{k=1}^n ((k+1) \cdot k! - 1 \cdot k!)$$

$$\therefore \boxed{A = \sum_{k=1}^n ((k+1)! - k!)} \quad [\because 0!(n+1)k! = (n+1)!]$$

For, $n=1$,

$$\cancel{\sum_{k=1}^n} ((k+1)! - k!) = (1+1)! - 1! = 1 =$$

For $n=2$,

$$\sum_{k=1}^n ((k+1)! - k!) = (2+1)! - 2! = 4 =$$

$$B = \prod_{i=2}^n \left(1 - \frac{1}{i}\right)$$

$$= \prod_{i=2}^n \left(\frac{i-1}{i}\right)$$

$$= \prod_{i=2}^n (i-1) \times \prod_{i=2}^n \frac{1}{i}$$

$$= (n-1)! \times \frac{1}{n!} \quad [\because \prod_{i=1}^n \frac{1}{i} \text{ or } \prod_{i=2}^n \frac{1}{i} = 1!]$$

$$= \frac{(n-1)!}{n(n-1)!}$$

$$\therefore \boxed{B = \frac{1}{n}} //$$

For $n=2$,

$$\prod_{i=2}^2 \left(1 - \frac{1}{i}\right) = \frac{1}{2} = \frac{1}{2} //$$

12340306

Mohammed Shadik Ansari

3-continue

$$C \cdot \sum_{k=1}^n \frac{1}{k(k+1)}$$

$$= \sum_{k=1}^n \left(\frac{1}{k} - \frac{1}{k+1} \right)$$

$$= \sum_{k=1}^n \frac{1}{k} - \sum_{k=1}^n \frac{1}{k+1}$$

$$= \left(\frac{1}{1} - \frac{1}{2} \right) \cdot$$

$$= \left(\frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{n} \right) - \left(\frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{n+1} \right)$$

$$= 1 - \frac{1}{n+1}$$

$$\therefore \boxed{C = 1 - \frac{1}{n+1}}$$

$$\text{For } n = 1,$$

$$C = 1 - \frac{1}{1+1} = \frac{1}{2}, \quad \square$$

$$\text{For } n = 2,$$

$$C = 1 - \frac{1}{2+1} = \frac{2}{3}, \quad \square$$

12340306

Mohammad Shadik Ansari

$$4. a) \sum_{i=1}^2 \prod_{k=1}^3 \frac{i}{k}$$

$$= \prod_{k=1}^3 \frac{1}{k} + \prod_{k=1}^3 \frac{2}{k}$$

$$= \frac{1}{1} \times \frac{1}{2} \times \frac{1}{3} + \frac{2}{1} \times \frac{2}{2} \times \frac{2}{3}$$

$$= \frac{1}{6} + \frac{8}{6}$$

$$= \frac{9}{6} = \frac{3}{2}$$

$$b) \prod_{k=1}^3 \sum_{i=1}^2 \frac{i}{k}$$

$$= \sum_{i=1}^2 \frac{i}{1} \times \times \sum_{j=1}^2 \frac{j}{2} \times \sum_{l=1}^2 \frac{l}{3}$$

$$= (1+2) \times \left(\frac{1}{2} + \frac{2}{2}\right) \times \left(\frac{1}{3} + \frac{2}{3}\right)$$

$$= 3 \times \frac{3}{2} \times \frac{3}{3}$$

$$= \frac{9}{2}$$

JKU

JOHANNES KEPLER
UNIVERSITÄT LINZ

12340306

Mohammad Shadik Ansari

4. continue

- c) No, these two instances are not enough to establish the general statement about the order of summation and product operation. But, indeed the order of both operations is very important and should not be changed.

5. Sol^h

x_i	-6	-3	-1	1	3	4
y_i	5	4	2	1	1	0

$$\begin{aligned} a) \quad \mu_x &= \frac{1}{6} \sum_{l=1}^6 x_l \\ &= \frac{1}{6} (x_1 + x_2 + x_3 + x_4 + x_5 + x_6) \\ &= \frac{1}{6} (-6 - 3 - 1 + 1 + 3 + 4) \end{aligned}$$

$$\therefore \boxed{\mu_x = -\frac{1}{3}}$$

$$\begin{aligned} \mu_y &= \frac{1}{6} \sum_{l=1}^6 y_l \\ &= \frac{1}{6} (y_1 + y_2 + y_3 + y_4 + y_5 + y_6) \\ &= \frac{1}{6} (5 + 4 + 2 + 1 + 1 + 0) \\ \therefore \boxed{\mu_y = \frac{13}{6}} \end{aligned}$$

$$b) \sigma_x^2 = \frac{1}{6} \sum_{l=1}^6 (x_l - \mu_x)^2$$

$$\begin{aligned} &= \frac{1}{6} ((x_1 - \mu_x)^2 + (x_2 - \mu_x)^2 + (x_3 - \mu_x)^2 + (x_4 - \mu_x)^2 + (x_5 - \mu_x)^2 + (x_6 - \mu_x)^2) \\ &= \frac{1}{6} \left((-6 + \frac{1}{3})^2 + (-3 + \frac{1}{3})^2 + (-1 + \frac{1}{3})^2 + (1 + \frac{1}{3})^2 + (3 + \frac{1}{3})^2 + (4 + \frac{1}{3})^2 \right) \\ &= \frac{1}{6} \left(\frac{289}{9} + \frac{64}{9} + \frac{4}{9} + \frac{16}{9} + \frac{100}{9} + \frac{169}{9} \right) \\ &= \frac{642}{54} = \frac{107}{9} \quad \therefore \boxed{\sigma_x^2 = \frac{107}{9}} \end{aligned}$$

$$\Rightarrow \sigma_{xy} = \frac{1}{6} \sum_{l=1}^6 (x_l - \mu_x)(y_l - \mu_y)$$

$$\begin{aligned} &= \frac{1}{6} \left((x_1 - \mu_x)(y_1 - \mu_y) + (x_2 - \mu_x)(y_2 - \mu_y) + (x_3 - \mu_x)(y_3 - \mu_y) + (x_4 - \mu_x)(y_4 - \mu_y) \right. \\ &\quad \left. + (x_5 - \mu_x)(y_5 - \mu_y) + (x_6 - \mu_x)(y_6 - \mu_y) \right) \\ &= \frac{1}{6} \left((-6 + \frac{1}{3})(5 - \frac{13}{6}) + (-3 + \frac{1}{3})(4 - \frac{13}{6}) + (-1 + \frac{1}{3})(2 - \frac{13}{6}) + (1 + \frac{1}{3})(1 - \frac{13}{6}) + (3 + \frac{1}{3})(4 - \frac{13}{6}) \right. \\ &\quad \left. + (4 + \frac{1}{3})(0 - \frac{13}{6}) \right) \\ &= \boxed{\frac{-107}{18}} \end{aligned}$$

12340306

Mohammed Shadik Ansari

$$c) k = \frac{\sigma_{xy}}{\sigma_x^2}$$

$$= \frac{-1071}{182}$$

$$= \frac{1071}{91}$$

$$\therefore k = -\frac{1}{2}$$

$$d) u_y = k u_x + d$$

$$\text{or, } d = k u_x - u_y$$

$$= -\frac{1}{2} \times \left(-\frac{1}{3}\right) - \frac{13}{6}$$

$$= \frac{1}{6} - \frac{13}{6}$$

$$= -\frac{12}{6}$$

$$\therefore d = -2$$

(-6, 5)

e)

(-3, 4)

(-1, 2)

(0, 1)

(1, 0)

④

x

$$f(x) = kx + d$$

$$(4, -9)$$

$$\text{For: } f(x) = kx + d = -\frac{1}{2}x - 2$$

$$\begin{array}{r|rr|r} x & 4 & 2 & -2 \\ \hline d & -9 & -3 & -1 \end{array}$$

JKU

JOHANNES KEPLER
UNIVERSITÄT LINZ

12340306

s. f) $f(-2) = 4$

Hence,

$$f(x) = kx + d = -\frac{1}{2}x - 2$$

$$\therefore f(2) = -\frac{1}{2} \times 2 - 2 = -3$$

$$\therefore f(-2) = -\frac{1}{2} \times (-2) - 2 = -1$$

Mohammed Shadik Ansari

JKU

JOHANNES KEPLER
UNIVERSITÄT LINZ