

SECTION 8.6 Applications of Inclusion–Exclusion

2. $1000 - 450 - 622 - 30 + 111 + 14 + 18 - 9 = 32$

4. $C(4+17-1, 17) - C(4+13-1, 13) - C(4+12-1, 12) - C(4+11-1, 11) - C(4+8-1, 8) + C(4+8-1, 8) + C(4+7-1, 7) + C(4+4-1, 4) + C(4+6-1, 6) + C(4+3-1, 3) + C(4+2-1, 2) - C(4+2-1, 2) = 20$

6. Square-free numbers are those not divisible by the square of a prime. We count them as follows: $99 - \lfloor 99/2^2 \rfloor - \lfloor 99/3^2 \rfloor - \lfloor 99/5^2 \rfloor - \lfloor 99/7^2 \rfloor + \lfloor 99/(2^2 3^2) \rfloor = 61$.

8. $5^7 - C(5, 1)4^7 + C(5, 2)3^7 - C(5, 3)2^7 + C(5, 4)1^7 = 16,800$

10. This problem is asking for the number of onto functions from a set with 8 elements (the balls) to a set with 3 elements (the urns). Therefore the answer is $3^8 - C(3, 1)2^8 + C(3, 2)1^8 = 5796$.

12. 2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321

14. We use Theorem 2 with $n = 10$, which gives us

$$\frac{D_{10}}{10!} = 1 - \frac{1}{1!} + \frac{1}{2!} - \cdots + \frac{1}{10!} = \frac{1334961}{3628800} = \frac{16481}{44800} \approx 0.3678794643,$$

which is almost exactly $e^{-1} \approx 0.3678794412 \dots$

16. There are $n!$ ways to make the first assignment. We can think of this first seating as assigning student n to a chair we will label n . Then the next seating must be a derangement with respect to this numbering, so there are D_n second seatings possible. Therefore the answer is $n!D_n$.

18. In a derangement of the numbers from 1 to n , the number 1 cannot go first, so let $k \neq 1$ be the number that goes first. There are $n - 1$ choices for k . Now there are two ways to get a derangement with k first. One way is to have 1 in the k^{th} position. If we do this, then there are exactly D_{n-2} ways to derange the rest of the numbers. On the other hand, if 1 does not go into the k^{th} position, then think of the number 1 as being temporarily relabeled k . A derangement is completed in this case by finding a derangement of the numbers 2 through n in positions 2 through n , so there are D_{n-1} of them. Combining all this, by the product rule and the sum rule, we obtain the desired recurrence relation. The initial conditions are $D_0 = 1$ and $D_1 = 0$.

20. We apply iteration to the formula $D_n = nD_{n-1} + (-1)^n$, obtaining

$$\begin{aligned} D_n &= n((n-1)D_{n-2} + (-1)^{n-1}) + (-1)^n \\ &= n(n-1)D_{n-2} + n(-1)^{n-1} + (-1)^n \\ &= n(n-1)((n-2)D_{n-3} + (-1)^{n-2}) + n(-1)^{n-1} + (-1)^n \\ &= n(n-1)(n-2)D_{n-3} + n(n-1)(-1)^{n-2} + n(-1)^{n-1} + (-1)^n \\ &\vdots \\ &= n(n-1) \cdots 2D_1 + n(n-1) \cdots 3 - n(n-1) \cdots 4 + \cdots + n(-1)^{n-1} + (-1)^n \\ &= n(n-1) \cdots 3 - n(n-1) \cdots 4 + \cdots + n(-1)^{n-1} + (-1)^n, \end{aligned}$$

which yields the formula in Theorem 2 after factoring out $n!$.

22. The numbers not relatively prime to pq are the ones that have p and/or q as a factor. Thus we have

$$\phi(pq) = pq - \frac{pq}{p} - \frac{pq}{q} + \frac{pq}{pq} = pq - q - p + 1 = (p-1)(q-1).$$

24. The left-hand side of course counts the number of permutations of the set of integers from 1 to n . The right-hand side counts it, too, by a two-step process: first decide how many and which elements are to be fixed (this can be done in $C(n, k)$ ways, for each of $k = 0, 1, \dots, n$), and in each case derange the remaining elements (which can be done in D_{n-k} ways).
26. This permutation starts with 4, 5, 6 in some order ($3! = 6$ ways to choose this), followed by 1, 2, 3 in some order ($3! = 6$ ways to decide this). Therefore the answer is $6 \cdot 6 = 36$.

SUPPLEMENTARY EXERCISES FOR CHAPTER 8

2. a) Let a_n be the amount that remains after n hours. Then $a_n = 0.99a_{n-1}$.
 b) By iteration we find the solution $a_n = (0.99)^n a_0$, where a_0 is the original amount of the isotope.
4. a) Let B_n be the number of bacteria after n hours. The initial conditions are $B_0 = 100$ and $B_1 = 300$. Thereafter, $B_n = B_{n-1} + 2B_{n-1} - B_{n-2} = 3B_{n-1} - B_{n-2}$.
 b) The characteristic equation is $r^2 - 3r + 1 = 0$, which has roots $(3 \pm \sqrt{5})/2$. Therefore the general solution is $B_n = \alpha_1((3 + \sqrt{5})/2)^n + \alpha_2((3 - \sqrt{5})/2)^n$. Plugging in the initial conditions we determine that $\alpha_1 = 50 + 30\sqrt{5}$ and $\alpha_2 = 50 - 30\sqrt{5}$. Therefore the solution is $B_n = (50 + 30\sqrt{5})((3 + \sqrt{5})/2)^n + (50 - 30\sqrt{5})((3 - \sqrt{5})/2)^n$.
 c) Plugging in small values of n , we find that $B_9 = 676,500$ and $B_{10} = 1,771,100$. Therefore the colony will contain more than one million bacteria after 10 hours.
6. We can put any of the stamps on first, leaving a problem with a smaller number of cents to solve. Thus the recurrence relation is $a_n = a_{n-4} + a_{n-6} + a_{n-10}$. We need 10 initial conditions, and it is easy to see that $a_0 = 1$, $a_1 = a_2 = a_3 = a_5 = a_7 = a_9 = 0$, and $a_4 = a_6 = a_8 = 1$.
8. If we add the equations, we obtain $a_n + b_n = 2a_{n-1}$, which means that $b_n = 2a_{n-1} - a_n$. If we now substitute this back into the first equation, we have $a_n = a_{n-1} + (2a_{n-2} - a_{n-1}) = 2a_{n-2}$. The initial conditions are $a_0 = 1$ (given) and $a_1 = 3$ (follows from the first recurrence relation and the given initial conditions). We can solve this using the characteristic equation $r^2 - 2 = 0$, but a simpler approach, that avoids irrational numbers, is as follows. It is clear that $a_{2n} = 2^n a_0 = 2^n$, and $a_{2n+1} = 2^n a_1 = 3 \cdot 2^n$. This is a nice explicit formula, which is all that “solution” really means. We also need a formula for b_n , of course. From $b_n = 2a_{n-1} - a_n$ (obtained above), we have $b_{2n} = 3 \cdot 2^n - 2^n = 2^{n+1}$, and $b_{2n+1} = 2 \cdot 2^n - 3 \cdot 2^n = -2^n$.
10. Following the hint, we let $b_n = \log a_n$. Then the recurrence relation becomes $b_n = 3b_{n-1} + 2b_{n-2}$, with initial conditions $b_0 = b_1 = 1$. This is solved in the usual manner. The characteristic equation is $r^2 - 3r - 2 = 0$, which gives roots $(3 \pm \sqrt{17})/2$. Plugging the initial conditions into the general solution and doing some messy algebra gives
- $$b_n = \frac{17 - \sqrt{17}}{34} \left(\frac{3 + \sqrt{17}}{2} \right)^n + \frac{17 + \sqrt{17}}{34} \left(\frac{3 - \sqrt{17}}{2} \right)^n.$$
- The solution to the original problem is then $a_n = 2^{b_n}$.
12. The characteristic equation is $r^3 - 3r^2 + 3r - 1 = 0$. This factors as $(r - 1)^3 = 0$, so there is only one root, 1, and its multiplicity is 3. Therefore the general solution is $a_n = \alpha_1 + \alpha_2 n + \alpha_3 n^2$. Plugging in the initial conditions gives us $2 = \alpha_1$, $2 = \alpha_1 + \alpha_2 + \alpha_3$, and $4 = \alpha_1 + 2\alpha_2 + 4\alpha_3$. Solving yields $\alpha_1 = 2$, $\alpha_2 = -1$, and $\alpha_3 = 1$. Therefore the solution is $a_n = 2 - n + n^2$.