

## Unit 2

# Molecular Orbitals and Pericyclic Reactions

### 2.10 Molecular Orbital Theory - 1

9/27:      • See Georgia's notes on Canvas (also included below).

## Lecture 10: MO Theory (1/2)

NO 1  
DATE 27 Sept 2024

### Exam Reflections

- You all did great
- "How did I do?"
  - 90-100 excellent
  - 80-90 good
  - <80 adequate, reach out
- exams hand back in recitations
- these guidelines are exam specific
  - ↳ first exam typically best
- You learned structure determination!


### What's Next?

- how do you go from 2D structure to where are  $e^-$   
where do they react

### Unit 2 Molecular Orbitals & Pericyclic Reactions

- deeper look at  $e^-$  movement
- new types of reactions, new class of mechanism
- use molecular orbitals (MO's) to predict reaction outcomes

↳ regioselectivity, diastereoselectivity, reactivity



- MO's provide insight into structure & reactivity

### Background/Review & Study

- review gen chem (5.11/5.12) & Orgo I (5.12)
- Clayden: ch. 4, 5, 6


27 Sept

## Lecture 10: (cont)


- orbitals are wavefunctions that describe the ability to find an electron in space
- ↳ they interact constructively & destructively

Rules:

- ① # atomic orbitals (AOs) in = # MO's out
- ② interacting orbitals must have similar energy
  - if large  $\Delta E$ ; no mixing
  - if same  $E$ ; best mixing




- ③ interacting orbitals must overlap efficiently and have similar energy symmetry



perpendicular orbitals

- poor overlap
- no mixing

- ④ More electronegative atom  $\Rightarrow$  lower  $E$  AO = more polarized MO

H<sub>2</sub> vs He<sub>2</sub> MO Diagramsbonding;  $e^-$  in middleNOTE:  $|\Delta E_{\text{bond}}| > |\Delta E_{\text{antibond}}|$ 

antibonding is more destabilizing than the bonding is stabilizing

H<sub>2</sub> MO more stable than 2  $\times$  H  $\cdot$  AO

↳ why H<sub>2</sub> bond forms





- antibonding MO filled!
- this is less stable than two individual H atoms
- forming He-He requires filling  $\sigma^*$  (which is more destabilizing) than.

• this is the MO explanation for the full octet rule



### SN<sub>2</sub> MO Picture



Why backside attack?  
• identify HOMO & LUMO

HOMO highest occupied MO = nucleophile, filled orbitals, lone pair  
LUMO lowest unoccupied MO = electrophile = empty orbitals =  $\pi^*$  or  $\sigma^*$  orbital, cations, C of C-Br bond

MO of C-Br



polarized towards Br

27 Sept

## Lecture 10: (cont)



backside attack because  
there is a larger  $\sigma^*$   
lobe there



meanwhile



$\sigma$  bond breaks because  $\sigma^*$  populated

our mechanistic arrows show this

### C=C vs C=O MOs



consequences:

C=C less reactive towards Nuc b/c  $\pi^*$  E too high

C=C generally nucleophilic unless somehow polarized  
ex:  $\text{CH}_2=\text{CH}-\text{C}(=\text{O})\text{CH}_3$

C=O electrophilic on carbon

## 2.11 Molecular Orbital Theory - 2

9/30: • Lecture 10 recap: What MO theory can explain.




Figure 2.1: MO theory explains these phenomena.

### – Regioselectivity.

- Consider a nucleophile adding into a carbonyl (Figure 2.1a).
  - Experimentally, we observe that the nucleophile attacks the carbon atom (magenta arrow) instead of the oxygen atom (grey dashed arrow).
- To understand why, we must consider the carbonyl's molecular orbitals (Figure 2.1d).
  - Specifically, we must consider the carbonyl's LUMO, since this will be the MO that interacts with the nucleophile's HOMO. Here, the LUMO is the carbonyl's  $\pi^*$ -orbital.
  - The carbonyl's LUMO has big lobes on carbon and small lobes on oxygen; in other words, this LUMO is **polarized** toward carbon.
  - The difference in lobe size explains why the nucleophile attacks carbon instead of oxygen.

### – Diastereoselectivity.

- Consider the bromination of an alkene (Figure 2.1b).
  - Experimentally, we observe that the *anti* adduct is formed instead of the *syn* adduct.
- To understand why, we consider the MOs of the bromonium ion intermediate (Figure 2.1e).
  - For the same reason as before, we must consider the bromonium ion's LUMO. Here, the LUMO is the C–Br  $\sigma^*$ -orbital.
  - The bromonium ion's LUMO has its largest lobe behind carbon.
  - Thus, this is the lobe that will be attacked by the  $\text{Br}^-$  nucleophile. Such an attack is called a “backside attack” and induces the *anti* product.

### – Reactivity.

- Consider a Grignard reagent adding into an olefin (Figure 2.1c).
  - Experimentally, we observe no reaction here.
- To understand why, we must consider the relative energies of the reacting MOs (Figure 2.1f).
  - Essentially, the alkene's LUMO (a  $\pi^*$ -orbital) is much higher in energy than the phenyl anion's HOMO. Thus, the  $\Delta E$  gap is too big, i.e., there is a lack of energy symmetry.
  - Therefore, by Rule 3 from Lecture 10, no reaction occurs.

- Today: More MO theory.
- Lecture outline.
  - The Bürgi-Dunitz angle.
  - Hyperconjugation.
  - The anomeric effect.
  - Stereoelectronic effects and the rate of reaction.
- **Bürgi-Dunitz angle:** The angle at which nucleophiles typically add to carbonyls. *Given by  $107^\circ$ .*




Figure 2.2: Bürgi-Dunitz angle.

- This is the angle between the new C–Nuc bond and the carbonyl's  $\sigma$ -plane (Figure 2.2a).
- Nucleophiles attack at this angle because it's the location of the  $\pi^*$ -lobe on carbon (Figure 2.1d).
- Let's elaborate a bit on Figure 2.1d now (Figure 2.2b).
  - Once again, consider the carbonyl  $\pi^*$ -orbital (its LUMO) and its “butterfly” lobes.
  - The nucleophile must approach the  $\pi^*$ -orbital with the right symmetry. This is why we see its HOMO's lobe approach the carbon atom's  $\pi^*$ -lobe dead-on at exactly the right angle.
    - This angle leads to efficient overlap, and hence an effective sharing of electron density.
    - This is an example of Rule 3 from Lecture 10.
  - Are there any other locations at which we can add into the carbonyl?
    - We can also add into the shaded carbon  $\pi^*$ -lobe on the other side of the  $\sigma$ -plane by reversing the shading of the nucleophile's lobe!
    - However, any other angle of attack will *not* work.
    - Note (\*): A backside attack is good for interacting with the  $\sigma^*$ -orbital, but bad for interacting with the  $\pi^*$ -orbital that we need for carbonyl chemistry.
- **Hyperconjugation:** The mixing of filled and empty orbitals to stabilize a system.
- Example (from 5.12): Stabilizing carbocations.




Figure 2.3: Hyperconjugation stabilizes carbocations.



- Consider a primary ( $1^\circ$ ) carbocation (Figure 2.3a).
  - In a carbocation, the positively charged carbon localizes its lack of electron density to an empty  $p$ -orbital.
  - However, adjacent to this empty  $p$ -orbital is a full  $\sigma$ -orbital, namely, the adjacent C–H bond. Moreover, this bond has the right *geometry* to donate into the empty  $p$ -orbital.
  - Thus, the  $\sigma$ -orbital of the C–H bond will donate electron density into the empty  $p$ -orbital, delocalizing both positive and negative charges and thereby stabilizing the system.
- We denote hyperconjugation interactions using a special **notation**; the particular hyperconjugation in Figure 2.3 is denoted  $\sigma_{\text{CH}} \rightarrow p_{\text{C}}$ .<sup>[1]</sup>
- In a tertiary ( $3^\circ$ ) carbocation, we get electron donation from *three* adjacent  $\sigma_{\text{CH}}$  orbitals.
  - These *three* stabilizing interactions explain why  $3^\circ$  carbocations are more stable than  $1^\circ$  ones!
  - Such effects are also why more substituted cations are more stable in general.
- **Hyperconjugation notation:** The concise method for denoting a certain hyperconjugative orbital interaction. *Given by*

$$\text{orbital}_{\text{atoms}} \rightarrow \text{orbital}_{\text{atoms}}$$
  - The arrow means “donates into.”
    - Indeed, we always write the filled orbital first (before the arrow) and the empty orbital second (after the arrow).
  - Possible orbitals:  $\sigma, \sigma^*, \pi, \pi^*, p, n$ .
    - Note that  $n$  denotes a nonbonding lone pair.
- **Anomeric effect:** The tendency of heteroatom substituents adjacent to heteroatoms in cyclohexane derivatives to prefer the axial orientation.
- Let’s break this rather complicated definition down through an example.







Figure 2.4: Anomeric effect.

<sup>1</sup>This is pronounced “sigma C–H to  $p$  C donation” or (very explicitly) “sigma see aech to pee see donation.”



- In methoxycyclohexane, the methoxy group prefers to be equatorial to avoid 1,3-diaxial interactions (Figure 2.4a).
  - This leads to a 70 : 30 distribution in favor of the equatorial conformer.
- However, in 2-methoxytetrahydropyran, the methoxy group prefers to be *axial* due to the anomeric effect (Figure 2.4b).
  - This *also* leads to a 70 : 30 distribution, but this time in favor of the axial conformer.
  - Notice how this empirical observation reflects the definition of the anomeric effect: We have a heteroatom substituent (the methoxy group) adjacent to a heteroatom in cyclohexane (the oxygen in the six-membered ring), and it is preferring the axial orientation!
- What causes the anomeric effect? Let's investigate the stabilization of the axial conformer further using molecular orbitals (Figure 2.4c).
  - In 2-methoxytetrahydropyran's equatorial conformation, we get poor overlap between the oxygen lone pair's orbital and the C–OMe antibonding orbital. This poor overlap is due to the *gauche* orientation of said orbitals.
  - In 2-methoxytetrahydropyran's axial conformation, we get really nice overlap between the oxygen lone pair and the  $\sigma^*$ -orbital of the C–OMe bond. This is because both orbitals have large lobes pointing axial down. Because of this favorable geometry,  $n_{\text{O}} \rightarrow \sigma_{\text{CO}}^*$  hyperconjugation occurs.<sup>[2]</sup>
- Another way of showing how the anomeric effect stabilizes the axial conformer is by using resonance diagrams (Figure 2.4d).
  - Indeed, starting from the typical picture, we can push the lone pair into an O=C  $\pi$ -bond and formally break the C–OMe  $\sigma$ -bond.
  - The result is called a **no-bond resonance form**.
  - Something should feel off to you here, though.
    - When you learned to draw resonance structures, you learned that you can't break  $\sigma$ -bonds.
    - However, we are now telling you that sometimes, you *are* allowed to break  $\sigma$ -bonds. This is “next-level resonance structures.”
  - Note that 2-methoxytetrahydropyran doesn't go all the way to the no-bond resonance form, but said resonance form *is* a major contributor.
    - This also means that the no-bond resonance form affects the reactivity of the molecule.
- Both hyperconjugation and the anomeric effect fall under the broader category of **stereoelectronic effects**.
  - Note that they are not the only examples of such effects, though.
- **Stereoelectronic effect**: An effect on structure or reactivity of a molecule caused by the spatial orientation of its orbitals.
  - We've previously learned that everything in Orgo can be explained by steric and electronic effects, but stereoelectronic effects are like a secret third option!
- Let's now look at some more places where stereoelectronic effects crop up.

---

<sup>2</sup>Note that there is no particular reason why overlap with a  $\sigma^*$ -orbital, in particular, is stabilizing. Rather, the point is that we have a filled orbital (the lone pair) adjacent to an empty orbital (which just happens to be a  $\sigma^*$  orbital), so hyperconjugation can occur to spread out the negative and positive charges. This delocalization — like any — is then inherently stabilizing.

- Example: Hyperconjugation in noncationic species.




Figure 2.5: Hyperconjugation stabilizes staggered ethane.

- We may have learned that ethane prefers the staggered conformer over the eclipsed conformer (Figure 2.5a) due to sterics.
  - This is not true!
  - We know this because H is really tiny.
- In fact, this preference is due to hyperconjugation, a stereoelectronic effect (Figure 2.5b).
  - Staggered ethane is stabilized by electron donation from the  $\sigma$ -bond of one C–H bond into the adjacent, antiperiplanar C–H bond's  $\sigma^*$  orbital:  $\sigma_{\text{CH}} \rightarrow \sigma_{\text{CH}}^*$ .
  - This is a small interaction, but it occurs six times, once for each C–H  $\sigma$ -bond!
- Takeaway: Electron delocalization is stabilizing, and more delocalization is more stabilizing.
- Example: Stereoelectronic stabilization can accelerate reactions.







Figure 2.6: Stereoelectronic effects accelerate reactions.

- Consider the  $S_N1$  substitution of bromocyclohexane to methoxycyclohexane (Figure 2.6a), vs. the  $S_N1$  substitution of 2-bromotetrahydropyran to 2-methoxytetrahydropyran (Figure 2.6b).
- Which of these substitutions occurs faster?
- To answer this question, let's look at the mechanism of each (Figures 2.6c-2.6d).
  - Note that in Figure 2.6c, either bromide or another equivalent of methanol can do the final deprotonation of the **oxonium** ion.<sup>[3]</sup>
  - Note that in Figure 2.6d, the fact that the **oxocarbenium** ion obeys the octet rule implies that it is the more stable resonance structure.
- In fact, the oxocarbenium ion is an example of oxygen stabilizing a carbocation through  $n_O \rightarrow p_C$  hyperconjugation.
- This is one example of hyperconjugation in this reaction scheme, but there is another effect as well.
  - In the original 2-bromotetrahydropyran molecule, the oxygen lone pair will also hyperconjugate into the C–Br  $\sigma^*$ -orbital per the anomeric effect.
  - In other words, O mediates the departure of the leaving group through  $n_O \rightarrow \sigma_{CBr}^*$  hyperconjugation.
- Thus, since both hyperconjugative stabilizing effects can (and do!) happen, it is better to say mechanistically that the arrow pushing in the first step happens simultaneously (Figure 2.6e).
  - Indeed, the rule in arrow pushing is “make a bond, break a bond,” so that's what we do.
- We can now complete the mechanism for the heteroatom-promoted reaction (Figure 2.6e).
  - MeOH adds into the  $\pi^*$ -orbital of the oxocarbenium (at the Bürgi-Dunitz angle!), also kicking electrons up to the oxygen in a concerted step.
  - Then we get deprotonation again.
- Now that we've got both mechanisms, let's consider the energy surface in order to compare the rates of reaction.
  - Both reactions will have two-humped energy surfaces, befitting a mechanism with only one true catinoic intermediate.
  - However, in the energy surface for the heteroatom-promoted reaction,  $n_O \rightarrow p_C$  hyperconjugation will stabilize the intermediate and  $n_O \rightarrow \sigma_{CBr}^*$  will stabilize the transition state of the first step, lowering its activation energy!
  - Thus, the heteroatom-promoted  $S_N1$  is faster!
- Takeaway: The overall reaction specifics depend on geometry and orbital overlap.

## 2.12 Pericyclics

10/2: • Lecture 11 recap.

- Hyperconjugation involves the delocalization of electrons, and hence is stabilizing.
  - This is a very common phenomenon, and it underlies most 5.12 reactions!
- $\sigma_{CH} \rightarrow p_C$  hyperconjugation makes substituted cations more stable (see Figure 2.3).
- $\sigma_{CH} \rightarrow \sigma_{CH}^*$  makes ethane more stable when staggered than eclipsed (see Figure 2.5b).
- $n_O \rightarrow \sigma_{CX}^*$  stabilizes axially positioned heteroatom substituents with  $\alpha$ -heteroatoms in cyclohexane derivatives, per the anomeric effect (see Figure 2.4c).

---

<sup>3</sup>Note that — comparing the  $pK_a$  of protonated methanol to HBr — methanol is actually almost a million times more basic than bromide. As such, for every one time bromide does the final deprotonation, methanol will do it to almost a million other oxonium intermediates. However, it can still be useful to think of bromide as *formally* doing the final deprotonation so as to balance the reaction  $C_6H_{11}Br + CH_3OH \longrightarrow C_6H_{11}OCH_3 + HBr$ .

- Today: Pericyclic reactions.
  - They are cool because they couple MO theory to reactivity.
  - This is a whole new class of reactions, and we will spend the rest of Unit 2 talking about them.
  - Essentially, the rest of this unit looks like: “Here’s a new reaction, and here’s the MO picture underlying it.”
- **Pericyclic** (reaction): A reaction characterized by a **concerted** movement of electrons in a **cyclic** transition state.
- **Concerted** (reaction): A reaction in which all electron movements happen at the same time, i.e., all bonds break and form in the same step.
  - There are no intermediates, and hence only a single “hump” in the energy diagram (Figure 2.7a).
  - Concerted reactions contrast with **stepwise** reactions.
  - Concerted reactions can be **synchronous** or **asynchronous**.
    - Essentially, we ask, “Is the transition state symmetric? Are all bonds breaking and forming to the same extent at the same time, or are some bonds breaking/forming first with others breaking/forming later?”
    - We touch on this concept to illustrate that even *concerted* reactions can have subtle differences between them.
- **Stepwise** (reaction): A reaction that *has* intermediates in its energy landscape.
  - There *are* intermediates, and hence multiple “humps” in the energy diagram (Figure 2.7b).
- To reiterate, concerted and stepwise reactions can be differentiated using their energy diagrams.




Figure 2.7: Concerted vs. stepwise energy diagrams.

- Example:  $S_N2$  reactions are concerted, and  $S_N1$  reactions are stepwise.
- **Synchronous** (concerted reaction): All bond-making and -breaking occurs to an equal extent in the transition state.
- **Asynchronous** (concerted reaction): All bond-making and -breaking does *not* occur to an equal extent in the transition state.
- **Cyclic** (transition state): A transition state in which all bonds that are being broken and made are connected in a ring.
  - Important implication: This is not as complicated as it sounds; rather, it just means that when you draw your electron arrows, you draw them in a ring.
  - See Figure 2.8 for an example.



- History of pericyclic reactions.
  - MO theory was developed *because* of pericyclic reactions.
    - Essentially, pericyclic reactions used to be called “no mechanism” reactions since everything happened in one step.
    - The extent to which organic chemists didn’t understand how pericyclic reactions worked drove them to develop a theory that explained why they did.
  - Today, pericyclic reactions are credited with introducing quantum theory into organic chemistry.
    - Before pericyclic reactions, organic chemists thought that they could explain everything they needed to with Lewis structures and arrow pushing mechanisms.
    - However, after pericyclic reactions, it became clear to organic chemists that there *was* value in keeping track of where all the electrons are actually located in MOs and such.
- We’ll now do an overview of the different classes of pericyclic reactions, i.e., what we have to look forward to over the next couple of weeks.
  - Types we’ll discuss: **Cycloadditions**, **electrocyclizations**, and **sigmatropic rearrangements**.
  - Specifically, we’ve got one lecture on each of these topics coming up (and then a couple others).
  - As such, it’s a good idea to get a general sense of these reactions now, but you don’t need to think too much about them since you will get much more information in the coming days and weeks.
- **Cycloaddition:** A pericyclic reaction in which two separate  $\pi$ -systems react to convert two  $\pi$ -bonds into two  $\sigma$ -bonds.
  - Nomenclature:  $[m + n]$ , where  $m$  and  $n$  are the numbers of atoms in the two separate  $\pi$ -systems.
    - To reiterate: One of the  $\pi$ -systems has  $m$  atoms, and the other has  $n$  atoms.
  - Example: We may speak of a “[4+2] cycloaddition.” This specific pericyclic reaction is also called the **Diels-Alder reaction**.
  - The reverse reaction of a cycloaddition is called a **cycloreversion**.
- **Diels-Alder reaction:** A  $[4 + 2]$  cycloaddition.




Figure 2.8: Diels-Alder reaction.

- This is a really cool reaction — one of the most powerful in organic chemistry, in fact.
- We’ll spend the next two lectures talking about it!
- Note that the magenta electron arrows in Figure 2.8 can go either counterclockwise *or* clockwise.
  - As long as they lead to the right product, you can draw either!
- **Cycloreversion:** A pericyclic reaction in which one system reacts to convert two  $\sigma$ -bonds back into two  $\pi$ -bonds in two separate  $\pi$ -systems.





Figure 2.9: A cycloreversion.

- Sometimes, we give cycloreversions special names.
- Example: We can call the reaction in Figure 2.9 either a “ $4\pi$  cycloreversion” or a “retro-[2 + 2].”

- **Electrocyclization:** A pericyclic reaction in which one system reacts to convert one  $\pi$ -bond into one  $\sigma$ -bond, or vice versa.

Figure 2.10: A forward and reverse  $6\pi$  electrocyclization.

- Nomenclature:  $m\pi$ , where  $m$  is the number of electrons involved.
- Always ring-opening or ring-closing.
- Example: We may speak of a “ $6\pi$  electrocyclization” (Figure 2.10a).
  - Notice how we go from 5  $\sigma$ - and 3  $\pi$ -bonds to 6  $\sigma$ - and 2  $\pi$ -bonds in Figure 2.10a.
- We can also go in reverse (Figure 2.10b).
- **Sigmatropic rearrangement:** A pericyclic reaction in which a  $\sigma$ -bond moves to the end of the  $\pi$ -system.




Figure 2.11: A [1,5] sigmatropic rearrangement.

- Nomenclature:  $[m,n]$ , where  $m$  and  $n$  are the “numbers” of the atoms to which the two ends of the  $\sigma$ -bond moves.
  - We’ll dive into this nomenclature more in the lecture on sigmatropic rearrangements.
  - In particular, we will discuss a very specific way of “numbering” the atoms in our starting material and product!
- Example: We may speak of a “[1,5] sigmatropic rearrangement” (Figure 2.11).
  - You can push the electron arrows either way, but Prof. Elkin likes to start with the  $\sigma$ -bond and move that to the end of the  $\pi$ -system.
  - Observe that one end of the  $\sigma$ -bond (the side at the hydrogen) moved from atom 1 to atom 5, and the other end (the side at the carbon) moved from atom 1 to atom 5. This is related to the aforementioned “numbering.”
- To reiterate from earlier, start familiarizing yourself with these types of reactions, but remember that we will go over these in more detail later in the course.
- Moving on, let’s bring MOs back into the picture.





Figure 2.12: Two views of buta-1,3-diene.

- Observe that the reactants in Figures 2.8, 2.10, and 2.11 are all conjugated systems!
- Thus, they have  $\pi$ -MOs. Let’s consider the diene from Figure 2.8, in particular.
- Looking at it from the side, we see that each  $\pi$ -bond is made up of two  $p$ -orbitals.

- We can mix the four  $p$ -AOs in Figure 2.12b to make MOs, but we have to do so according to the following rules.
  1. The number of MOs is equal to the number of atoms under consideration.
    - Example: A diene will have four MOs.
    - Example: An olefin will have two MOs.
  2. The lowest-energy MO has no **nodes**.
  3. For every increase in  $E$ , we add a node (in such a way that symmetry is maintained).
- **Node:** A change in sign of the orbital, at which there is no electron density.
- Example of nodes: Think of our  $\pi$  and  $\pi^*$  MOs.

Figure 2.13: Nodes in  $\pi$  and  $\pi^*$  molecular orbitals.

- The lower one has no nodes, because the phases are aligned left to right.
- The upper one has 1 node, because the phases invert left to right.
- We are now ready to draw an MO diagram for the diene in Figure 2.12.
- Diene MOs.




Figure 2.14: Reactive molecular orbitals of a diene.

- Let's start with a rule-by-rule analysis.
  1. We have 4 atoms, so we draw lines for 4 MOs.
  2. The lowest energy MO in Figure 2.14 does indeed have no nodes.
    - Observe that Prof. Elkin shades the top lobes first this time instead of the bottom lobes (as in Figure 2.13) because the shading is arbitrary. This comment is review, but the concept is important to remember!!
  3. For each increase in energy, we do indeed add one node.
    - For the second-lowest energy level, we draw our node symmetrically right in the middle.
      - We go along shading the top (or bottom) lobes until we hit our node, and then we switch to shading the other side.
    - For the third-lowest energy level, we draw our nodes symmetrically as well.
    - For the highest energy level, we draw a node between every orbital and alternate shading.
      - The highest energy level has the same alternating structure in the MOs of every conjugated  $\pi$ -system. For another example, see Figure 2.16.
- Note that we also draw (in purple) the **waveform** for every MO.
- Now let's populate our orbitals with electrons.
  - There are four  $\pi$ -electrons in a diene, so per Aufbau, Pauli, and Hund, we fill the bottom two energy levels of our diagram.
  - Filling electrons allows us to identify our HOMO and LUMO, which will be useful for justifying reactivity.
- Takeaway: You probably wouldn't just guess that the LUMO (or any other MO) of a diene looks the way it does, but you can derive it using the three rules and the method of Figure 2.14.
  - Then you can use the result of your derivation to make predictions about a diene's reactivity!
  - We'll cover such predictions next lecture.
- Why do the nodes have to be symmetric?
  - Because quantum mechanics.
  - Very simply, it has something to do with the waveform of each energy level, which you might notice mirrors the waveforms of the particle in a box.
  - See the end of my notes for this lecture for more detail.
    - Note: All extra detail on this topic is beyond the scope of this class, and will never be tested nor appear on problem sets; it is purely to satisfy your curiosity.
- An interesting finding about pericyclic reactions: They can be started by either heat ( $\Delta$ ) or light ( $h\nu$ )!




Figure 2.15: A diene's reactive orbitals in thermal vs. photochemical pericyclic reactions.



- When you think about it, heat and light are just different ways to add energy to our system so that the reaction goes.
- Indeed, pericyclic reactions are cool because you don't have to add a chemical reagent to make one go; rather, you just heat it up or shine light at it, and it reacts away!
- How do photochemical reactions work?
  - When light is absorbed, one electron is excited from the HOMO to the LUMO, and none of the spins of *any* of the electrons change (Figure 2.15).
    - There's a lot more photophysics here that you can go into, but that's beyond the scope of this course.
  - Such excitation is important because it gives us a new HOMO and a new LUMO.
    - These new reactive orbitals have important consequences that we'll discuss later, especially for the stereochemistry of the product.
- Let's now look at the MOs of one more conjugated system.
- Allyl MOs.




Figure 2.16: Reactive molecular orbitals of an allyl group.

- Let's do another rule-by-rule analysis.
  1. Three atoms going in means three MOs.
  2. Shade all the same phases in the bottom MO.
  3. Add nodes for the upper orbitals.
    - Put your node right in the middle for the middle MO.
      - It has to be symmetric!
      - Every time you have an odd number of atoms, some *p*-orbital will get deleted like this.
    - For the top MO, once again do everything alternating.
- Note that we have not yet specified whether this is an allyl cation, allyl radical, or allyl anion!
- We will have a different number of electrons for all three species (even though we have the same MOs), so let's fill electrons for each of these species.
  - Allyl cation: Two electrons, so fill just the bottom MO.
    - Like any carbocation, the allyl cation will react as an electrophile.
    - If it reacts as an electrophile, it must react with its LUMO (which is the middle orbital).
  - Allyl radical: Three electrons, so fill the bottom MO and start filling the middle MO.
    - Like any radical, the allyl radical reacts as a... well... radical.
    - If it reacts as a radical, it must react with its **SOMO** (also the middle orbital).

- Allyl anion: Four electrons, so fill the bottom and middle MOs.
  - Like any carbanion, the allyl anion will react as a nucleophile.
  - If it reacts as a nucleophile, it must react with its HOMO (still the middle orbital!).
- Interesting consequence of this filling: All three allyl species should only react with their middle-energy MO!
  - This would predict that all allyl reactivity occurs at the termini of the allyl group, not the middle carbon, since all of the density of the middle orbital is at the termini and none of it is at the middle carbon.
  - This prediction is experimentally confirmed!
- **Singly occupied molecular orbital:** The molecular orbital in which an unpaired radical electron exists. *Also known as SOMO.*
- An elaboration on why nodes must be placed symmetrically in the MOs of conjugated  $\pi$ -systems (see Figures 2.14 and 2.16 and the associated discussion).
  - Reminder: Everything from here, on, in these notes is beyond the scope of this class!
  - The long-short.
    - The waveforms in Figure 2.14 are *exactly* equal to their corresponding particle-in-a-box wave functions (according to some analyses of quantum mechanics).
      - This relationship can be rationalized intuitively because a conjugated  $\pi$ -system is like an extended, one-dimensional box in which a quantum particle (namely, an electron) lives.
    - The implication is that the molecular orbitals of a conjugated  $\pi$ -system look *exactly* like the particle-in-a-box orbitals, including having nodes in the same places.
    - This actually also means that the individual  $p$ -orbitals making up the MOs are different sizes!
      - For example, in the lowest energy MO in Figure 2.14, the two middle  $p$ -orbitals will be larger than the two terminal  $p$ -orbitals.
      - More relevantly, the HOMO and LUMO in Figure 2.14 will have larger terminal  $p$ -orbitals, which explains why dienes react at their ends and not in the middle; we have already seen an example of dienes reacting at their terminal carbons instead of their middle carbons in Figure 2.8.
  - More detail.
    - The exact sizes of each  $p$ -orbital in a given MO of a conjugated  $\pi$ -system can be calculated — by hand — using only linear algebra. This calculation is part of something called **Hückel theory**.
    - You can learn about Hückel theory by taking a course in quantum mechanics, inorganic chemistry, or graduate physical organic chemistry.
  - If you are interested in reading more about this now, look through the attached PDF. I'd recommend starting with the diagrams and sine/cosine functions on pages 6.6 and 6.7. Enjoy!

## 2.13 Diels-Alder - 1

10/4:

- Lecture 12 recap.
  - Pericyclic reactions have concerted and cyclic transition states.
    - Essentially, what unites all of these reactions is that they have electron arrows moving in a ring!
  - All of these reactions are theoretically reversible.
  - Prof. Elkin redraws the prototypical pericyclic reactions from last class.

- Three main classes.
  1. Cycloaddition.
    - Bond types changed:  $2\pi \rightleftharpoons 2\sigma$ .
    - Nomenclature:  $[m + n]$ .
    - General form: See Figure 2.8.
      - To reiterate: This reaction can also proceed in reverse, i.e., from right to left!
  2. Electrocyclization.
    - Bond types changed:  $1\pi \rightleftharpoons 1\sigma$ .
    - Nomenclature:  $m\pi$ .
    - General form: See Figure 2.10.
      - To reiterate: This reaction can also proceed in reverse, i.e., from right to left!
  3. Sigmatropic rearrangements.
    - Bonds moved:  $1\sigma$ .
    - Nomenclature:  $[m, n]$ .
    - General form: See Figure 2.11.
      - To reiterate: This reaction can also proceed in reverse, i.e., from right to left!
- Announcements.
  - Please fill out the feedback survey in Canvas > Announcements.
  - PSet 3 is due today.
  - PSet 4 will be posted today.
    - It is the last PSet before Exam 2.
    - It only covers Diels-Alder content. However, the rest of this unit's content (cycloadditions, electrocyclizations, and sigmatropic rearrangements) *will* be on Exam 2 as well.
    - So to prepare for the exam, continue doing the Recitation Worksheets even after PSet 4!!
- Today: Diels-Alder (lecture 1 of 2).
- Recall from last lecture that a *Diels-Alder reaction* is a  $[4 + 2]$  cycloaddition.
  - Specifically, a **diene** reacts with an olefin, which we call the **dienophile**.
  - The simplest Diels-Alder (DA) reaction is drawn in Figure 2.8.
    - This is actually a terrible Diels-Alder reaction because there's a poor HOMO-LUMO energy match (we'll talk more about what that means shortly).
  - The Diels-Alder is a powerful tool to make six-membered rings.
    - We see a lot of six-membered rings in organic chemistry, so the Diels-Alder is very useful.
  - This reaction is very predictable: It is **regioselective**, **stereospecific**, and **reliable**.<sup>[4]</sup>
- **Diene**: A compound that contains two conjugated double bonds.
  - The diene is (usually) the HOMO.
  - You can think of it as the nucleophile.
- **Dienophile**: An olefin. *Etymology* from Latin “lover of dienes.”
  - The dienophile is (usually) the LUMO.
  - You can think of it as the electrophile.
- **Reliable** (reaction): A reaction that almost always works if you have the right energy matching.

---

<sup>4</sup>We'll discuss both regioselectivity and stereospecificity later this lecture, and stereospecificity even further on Monday.

- Let's look at the MO picture for a Diels-Alder reaction.




Figure 2.17: Reactive molecular orbitals in a Diels-Alder.

- Recall our diene MOs from last lecture (Figure 2.14).<sup>[5]</sup>
    - Specifically, recall that our HOMO has the orbital picture of the second energy level. Since we have said that the diene reacts with its HOMO, this is the important orbital to watch.
  - Recall also our dienophile MOs from last lecture (Figure 2.13).
    - The olefin has two electrons, so its LUMO is the second energy level. Since we have said that the dienophile reacts with its LUMO, this is the important orbital to watch.
  - Initially, we have a poor energy match between HOMO and LUMO ( $\Delta E$  is large). Therefore, if we want to improve the reaction, we should strive to bring their energies closer together.
  - Two main things to accomplish this goal of bringing HOMO and LUMO energies closer together.
    - Raise the HOMO by adding EDGs to the diene.
    - Lower the LUMO by adding EWGs to the dienophile.
- Example: A Diels-Alder reaction that does work well.




Figure 2.18: Diels-Alder orbitals in 3D space.

<sup>5</sup>Prof. Elkin reviews the three rules for drawing the MOs of conjugated systems; remember these!!



- Let's analyze the new reactants we've drawn (Figure 2.18a).
  - The methoxy-substituted diene is a better nucleophile than 1,3-butadiene because methoxy groups are electron-donating.
  - The ester-substituted dienophile is a better electrophile than ethylene because ester groups are electron-withdrawing.
  - Recall electrophilic aromatic substitution reactions, which tell you what substituents are electron donating vs. withdrawing. Review this 5.12 content!!
- Now let's look at their orbitals (Figure 2.18b).
  - We draw a "perspective picture" of the HOMO and LUMO.
  - Observe that the phases of the HOMO and LUMO match!
    - Specifically, we mean that the lobes connected by the blue lines have the same shading.
  - A note on shading.
    - The *relative* shading between reacting molecules *does* matter because we've got to see overlap between pairs of shaded lobes and pairs of unshaded lobes when we're forming bonds.
    - Thus, while we could invert the shading of every *p*-orbital in Figure 2.18b and be fine, we *could not* invert the shading of just the diene and leave the dienophile unchanged (or vice versa).
- We now redraw the molecules, but after they've formed  $\sigma$ -bonds (Figure 2.18c).
  - As the  $\pi$ -orbitals come together, the middle lobes fuse and become  $\sigma$ -bonds.
  - Implication: You have to have a top-to-bottom approach so that the *p*-orbitals interact and mix. A side-to-side overlap would not form  $\sigma$ -bonds from *p*-orbitals.
- Accelerating Diels-Alder reactions.




Figure 2.19: The relative rate of reaction of different Diels-Alder substrates.

1. Raise the HOMO by putting EDGs on the diene (Figure 2.19a).
    - Adding a methoxy substituent (an EDG) will increase the rate regardless of the position to which you add it.
    - However, interestingly enough, it will increase the rate *more* when added to some positions over other positions.
    - This is because of the difference between **cross-conjugation** and regular conjugation.
  2. Lower the LUMO by putting EWGs on the dienophile (Figure 2.19b).
    - From left to right, the names of these seven compounds are: ethylene, acrylonitrile, propenal, *cis*-1,2-dicyanoethene, *para*-quinone, tetracyanoethene, and 4-phenyl-1,2,4-triazole-3,5-dione (PTAD<sup>[6]</sup>).
  3. Enforce the ***s-cis*** configuration (Figure 2.19c).
    - The leftmost compound is locked in the ***s-trans*** conformation.
    - The next one has a big steric clash between methyl groups, so it's much more stable in the *s-trans* configuration.
    - Buta-1,3-diene likes to be *s-trans* because it still has sterics from the hydrogens.
    - Penta-1,3-diene has the same mild steric preference for *s-trans* as buta-1,3-diene.
      - However, certain stereoelectronic effects (which you'll work out on PSet 4!) promote its reactivity.
      - Essentially, methyl groups are slightly electron-donating.
    - Cyclopentadiene is locked in an *s-cis* conformation.
- **Cross-conjugated** (molecule): A molecule containing multiple olefins that — despite being arranged in a row — do not delocalize efficiently.




Figure 2.20: Cross-conjugation vs. regular conjugation.

- Notice that for the molecule in Figure 2.20a, we cannot draw a resonance structure that engages the bottom  $\pi$ -bond.
  - In contrast, regular conjugation (Figure 2.20b) disperses the oxygen's electron density across the entire  $\pi$ -system.
- ***s-cis*** (conformer): The rotational isomer of a diene in which the alkenes are *cis* relative to the  $\sigma$ -bond.



Figure 2.21: *s-cis* diene.

<sup>6</sup>According to Sauer and Schröder (1967), PTAD is approximately  $10^5$  times faster than tetracyanoethene. This means that it is approximately  $4 \times 10^{12}$  times faster than acrylonitrile!


- **s-trans** (conformer): The rotational isomer of a diene in which the alkenes are *trans* relative to the  $\sigma$ -bond.

Figure 2.22: s-*trans* diene.

- Essentially, olefins can be *cis* or *trans*.

Figure 2.23: Reacting *cis* and *trans* olefins in the Diels-Alder.




- When you react the *cis*-olefin in a Diels-Alder, you get (exclusively) the *cis*-product.
- When you react the *trans*-olefin in a Diels-Alder, you get (exclusively) the *trans*-product.
- This means that the Diels-Alder is **stereospecific**.<sup>[7]</sup>
- On the other hand, dienes can be s-*cis* or s-*trans*.

Figure 2.24: s-*cis* and s-*trans* conformers rapidly interconvert.


- The “s” stands for “sigma bond.”
- These two diene conformers are not discrete species; rather, they interconvert like the gauche, anti, staggered, etc. conformers of ethane.
- Only s-*cis* dienes react in Diels-Alders, so enforcing that geometry accelerates the reaction.
- **Stereospecific** (reaction): A reaction in which the stereochemistry of the reactants translates directly into a single stereochemical product.
- **Stereoselective** (reaction): A reaction in which when a certain stereochemical product is favored, but a mixture is still produced and the stereochemistry of the reactants doesn’t exert excessive influence.
- **Stereoretentive** (reaction): A reaction in which the exact stereocenters present in the starting material are conserved in the product.
- Does the rightmost diene in Figure 2.19a have to be *trans* at the upper alkene?
  - Yes; we’ll talk about that more next lecture.
- Can you lower the LUMO so much that the Diels-Alder reaction no longer proceeds?
  - Yes! This is related to **inverse electron-demand** Diels-Alder reactions, discussed next Monday.
- Can you have a photochemical Diels-Alder?
  - They’re quite rare, but you can do similar things.
  - Note: Lewis acid acceleration of Diels-Alders will also be discussed next Monday.

<sup>7</sup>Next lecture, you will learn how the Diels-Alder is **stereoselective** but still not **stereoretentive**.


- Regioselectivity.

(a) *meta*-product.(b) *para*-product.


(c) Diene (resonance).




(d) Dienophile (resonance).




(e) Diene (hyperconjugation).



(f) Dienophile (hyperconjugation).



(g) Resonance matching.



(h) Hyperconjugation matching.

Figure 2.25: Diels-Alder regioselectivity.

– Observation: Depending on how the diene and dienophile are oriented when they react, we could conceivably form two different products (Figures 2.25a-2.25b).

■ These two products are called the *meta*-product and the *para*-product.

- So which product do we actually get?
    - We will determine this by using either *resonance* or *hyperconjugation/partial negative charges* to identify and match the electron-rich and electron-poor sites on our diene and dienophile.<sup>[8]</sup>
    - Use whichever method you prefer since they give the same result, but you should learn both!!
  - The resonance analysis (Figures 2.25c-2.25d).
    - As in Figure 2.20, push arrows as far as we can to get the reactive resonance structure.
  - The hyperconjugation analysis (Figures 2.25e-2.25f).
    - Begin with an atom that we *know* will have a partial positive ( $\delta^+$ ) or partial negative ( $\delta^-$ ) charge.
      - For example, we know that oxygen will be  $\delta^-$  because it is the most electronegative atom in both structures.
      - Thus, we can label it first in both Figures 2.25e and 2.25f.
    - Then expand out over the rest of the conjugated system, alternating  $\delta^+$  or  $\delta^-$  atom-to-atom.
      - So since oxygen is  $\delta^-$ , the carbon  $\alpha$  to it should be  $\delta^+$ , the carbon(s)  $\beta$  to it should be  $\delta^-$ , the carbon(s)  $\gamma$  to it should be  $\delta^+$ , etc.
    - Keep track of the partial charges on the termini of the diene and dienophile (i.e., the reactive sites). These are circled in Figures 2.25e-2.25f.
  - Notice the agreement/consistency between the two methods!
    - Indeed, the carbanion in Figure 2.25c corresponds to a  $\delta^-$  carbon in Figure 2.25e, and the carbocation in Figure 2.25d corresponds to a  $\delta^+$  carbon in Figure 2.25f.
  - Once we have performed either analysis, matching up the negatives on the diene to the positives on the dienophile and vice versa predicts our product!
    - Thus, by both analyses, the *para*-product is favored!
    - This matching of positive and negative charges is indicative of the maxim that “organic chemistry is just magnets everywhere.”
  - Exercise: Try drawing the meta-product, which will force you to put positive near positive and negative near negative.
    - “That’s not fun, that’s not how magnets work.”
- **meta-product:** The product of a Diels-Alder reaction in which the substituents would be oriented *meta* to each other on the six-membered ring, if the six-membered ring were aromatic.
  - **para-product:** The product of a Diels-Alder reaction in which the substituents would be oriented *para* to each other on the six-membered ring, if the six-membered ring were aromatic.
  - Prof. Elkin has many more examples, but will go through them on Monday.

## 2.14 Diels-Alder - 2

10/7:

- Lecture 13 recap.
  - Corrections to Lecture 13.
    - The word “Diels-Alder” usually has a hyphen; it is *not* written “Diels Alder” with a space.
    - The methyl-activated diene in Figure 2.19c is faster because of *electronics*, not sterics.
      - Indeed, this diene has the same steric preference for *s-trans* as buta-1,3-diene, but it is slightly faster because methyl groups are slightly electron-donating.





---

<sup>8</sup>Note that there is also a secret third method: MO theory! See Clayden et al. (2012, p. 890) for how to do this. As we would expect, the predictions we get by both resonance and hyperconjugation match the predictions of MO theory. (Note also that you are not required to learn the MO theory method because it was not covered in class; this comment is purely to point out an interesting connection :)



- Definitions of **stereoretentive**, **stereospecific**, and **stereoselective** reactions.
  - Recap of regioselectivity.
    - Prof. Elkin redraws Figure 2.25h as a method of predicting the *para*-product (Figure 2.25b).
- **Stereoretentive** (reaction): A reaction in which the stereocenter(s) in the starting material are retained exactly.
  - Example: (*R*)-butan-2-ol reacts and is still an (*R*)-alcohol.
- **Stereospecific** (reaction): A reaction in which the stereocenter(s) of the starting material decide the stereocenter(s) of the product.
  - Example: Figure 2.23.
  - To reiterate: The phenomenon exemplified by Figure 2.23 is an example of stereospecificity, *not* stereoselectivity (as was incorrectly said last lecture).
- **Stereoselective** (reaction): A reaction in which a certain stereoisomer of the product is preferred.
  - Example: Butan-2-one is reduced to (*R*)-butan-2-ol, instead of racemic ( $\pm$ )-butan-2-ol.
  - Stereoselective is an “umbrella” term: Both stereoretentive and stereospecific reactions are stereoselective, but not all stereoselective reactions are stereoretentive or stereospecific.
- Today: More on the Diels-Alder reaction.
- Lecture outline.
  - More on regioselectivity.
  - *exo* vs. *endo* transition states.
  - *exo* vs. *endo* products.
  - Lewis acid catalysts for Diels-Alder reactions.
  - Inverse electron-demand Diels-Alder reactions.
  - Example Diels-Alder reactions; relevant to PSet 4!
- We'll begin today by continuing our discussion of regioselectivity.
  - Recap of Figure 2.25.
    - Note that like we have the *meta*-product and *para*-product, we can have the *ortho*-product.
  - General rules (timesavers).
    1. A single EDG on the diene and EWG on the dienophile (usually) leads to the *ortho*-product or the *para*-product, not (usually) the *meta*-product.
    2. If there isn't a clear preference (e.g., weak EDG or EWG only), you get a mixture of products.
      - Example: The methyl-activated diene, penta-1,3-diene (see Figure 2.19c), reacts with propenal to give both the *ortho*- and *meta*-products in an 8:1 ratio. On the other hand, the methoxy-activated diene, 1-methoxybuta-1,3-diene (see Figure 2.19a), reacts with propenal to give 100% of the *ortho*-product.
- Switching subjects, let's finally investigate what determines the full stereochemistry of the product.
  - We have previously discussed the *stereospecificity* implied by *cis* or *trans* starting materials, but we will now discuss a type of *stereoselectivity*.
  - Specifically, we will need to begin our investigation with a slight detour to define and analyze the *endo* and *exo* transition states.
- **endo** (transition state): The Diels-Alder TS in which the dienophile's EWG points *toward* the diene.
- **exo** (transition state): The Diels-Alder TS in which the dienophile's EWG points *away from* the diene.

- *exo* vs. *endo* transition states.

(a) *exo* transition state.(b) *endo* transition state.(c) *exo* orbitals.(d) *endo* orbitals.Figure 2.26: *exo* vs. *endo* transition states in the Diels-Alder.

- Essentially, observe that when the starting materials in a Diels-Alder reaction encounter each other, the dienophile's substituent can either point “out” (Figure 2.26a) or “in” (Figure 2.26b).
  - Remember that per Figure 2.18, the substituents have to encounter each other top-to-bottom.
  - Remember that per regioselectivity, the reactants will encounter each other as drawn so as to form the *ortho*-product.
- The lower energy transition state will lead to more product, so let's compare their relative energies.
- To do this, we will have to consider the “full LUMO” of the dienophile.
  - While we've only considered the dienophile's alkene functional group so far, observe that methyl acrylate also has an adjacent carbonyl  $\pi$ -bond that can conjugate with the alkene functional group just like in a diene!
  - Thus, four  $p$ -orbitals will form the “full MOs” of the dienophile.
  - The four dienophile MOs resulting from these four  $p$ -orbitals can be drawn using the three rules from Lecture 12, resulting in a picture exactly like Figure 2.14.
  - Since the dienophile will still be reacting with its LUMO, the “full LUMO” of the dienophile will look like the third energy level in Figure 2.14. Indeed, that's what we draw on the dienophile in both Figures 2.26c-2.26d!
- In both the *exo* and *endo* transition states, our “full LUMO” has **primary orbital interactions** with the diene HOMO analogous to Figure 2.18b. This enables the formation of  $\sigma$ -bonds as we'd expect, even (to reiterate) with the “full LUMO.”
- However, there are also some differences between the transition states.
  - *exo* transition state: There is *less steric clash* between the diene/dienophile substituents.
  - *endo* transition state: There is a new **secondary orbital interaction** that is stabilizing.
- This secondary orbital interaction is (usually) more stabilizing than the lack of sterics, so the *endo* transition state is (usually) preferred!
  - This is technically a  $\pi_{\text{HOMO}} \rightarrow \pi_{\text{full LUMO}}^*$  interaction.

- **Primary** (orbital interaction): An orbital interaction that leads to bonding.
- **Secondary** (orbital interaction): An orbital interaction that doesn't lead to bonding, but is stabilizing (or destabilizing).
- Let's now connect *endo* and *exo* transition states back to the stereochemistry of the product in a Diels-Alder reaction.




Figure 2.27: The transition state predicts Diels-Alder product stereochemistry.

- In the course of a Diels-Alder reaction, once we form our (*endo* or *exo*) transition state, we will subsequently form bonds and then unfold the structure — like a book — into our product.
  - Consider drawing in the hydrogens to help see how we get from the second to the third picture.
- For the particular Diels-Alder reaction we've considered in both Figures 2.26 and 2.27, the *exo* transition state yields the *trans*-product and the *endo* transition state yields the *cis*-product.
  - Because the *cis*-product arises from the *endo* transition state (which, to reiterate, is the preferred transition state), the *trans*-product will be preferred for this reaction!
  - I.e.: The *cis*-product is the major product, and the *trans*-product is the minor product.
- Since the Diels-Alder is not **enantioselective**, we can draw either enantiomer of the product.
  - In other words: We could switch all wedges and dashes for dashes and wedges, respectively, in Figure 2.27a-2.27b and still have the right answer.
- Help digesting this material: Look for some online visualizations, get a molecular model kit, etc.!!
- **Enantioselective** (reaction): A reaction that favors one enantiomer of the product over another.
- Example: Predicting the proper stereochemistry in the following Diels-Alder reaction.




Figure 2.28: Diels-Alder reaction of (*Z*)-1-methoxybuta-1,3-diene and methyl acrylate.

- The rate will be slower than if (*E*)-1-methoxybuta-1,3-diene were used because the diene in Figure 2.28 is *s-cis* destabilized.
  - The *s-cis* destabilization comes from steric clashing between the methoxy group and the hydrogen.
- The regioselectivity will pair the bottom carbon on the diene in Figure 2.28 to the bottom carbon on the dienophile in Figure 2.28, yielding the *ortho*-product.
- The stereoselectivity will favor the *endo* transition state because the dienophile's EWG has a  $\pi$ -system that can participate in secondary orbital interactions with the diene.
  - Thus, we will favor the drawn diastereomer (and its enantiomer!).
  - Note that in this reaction, the *endo* transition state yields the *trans*-product. This is the opposite of Figure 2.27b, in which the *endo* transition state yielded the *cis*-product.
  - This illustrates that it's not always the *cis*-product that's major! Rather, whether *cis* or *trans* is major depends on the transition state (*endo* or *exo*) from which each originates.
- Tip for learning this content: Just practice, esp. drawing the product regio- and stereochemistry.
- This wraps up all we need to say about *endo* and *exo* transition states.
- So, switching topics, let's discuss accelerating Diels-Alder reactions with catalysis.
  - So far, every Diels-Alder reaction we've considered has been thermal.
  - We can accelerate these reactions with a Lewis acid catalyst (such as  $\text{SnCl}_4$ ,  $\text{EtAlCl}_2$ , etc.).
  - This allows us to do our Diels-Alder reactions...
    1. At a lower temperature;
    2. With greater stereoselectivity.
    - Lower temperatures make the *endo* transition state even more preferred.
- Example: How does this Lewis acid catalysis work?




Figure 2.29: Lewis acid catalysis of Diels-Alder reactions.

- Three-step process.
  1. The Lewis acid (LA) coordinates to the dienophile's EWG, making it more electron poor.
  2. This lowers the LUMO even further, which gives you better energy overlap.
  3. Better energy overlap stabilizes the transition state.
- Essentially, Lewis acid catalysts work by making our EWG "better."
- Switching topics again, let's discuss something to which we've alluded a few times: What happens when the diene is super stabilized and the dienophile is super destabilized.
- **Inverse electron-demand** (Diels-Alder reaction): A Diels-Alder reaction in which the HOMO of the dienophile interacts with the LUMO of the diene.
  - Still *ortho*/*para*-directing.
  - Still *endo* TS preferred.
  - Often see this when we have a heteroatom in the ring.

- Example inverse electron-demand Diels-Alder reaction.







Figure 2.30: Diels-Alder reaction with inverse electron-demand.

- Because of the methyl ester EWG, the diene in Figure 2.30 is now electron-poor.
  - As such, it will react with its LUMO (the third energy level in Figure 2.14).
- Because of the amine EDG, the dienophile in Figure 2.30 is now electron-rich.
  - As such, it will react with its HOMO (the first/lowest energy level in Figure 2.13).
- However, when we draw the HOMO of the dienophile and LUMO of the diene, we still get good orbital overlap. Thus, this reaction can still proceed, forming the *ortho-cis*-product through the *endo* transition state.
- Why is the *endo* TS still preferred?
  - The exact orbital interactions here deal more with coefficients and differently sized orbital lobes, but that is beyond the scope of this class.
  - Take 5.43 and 5.53 if you want to learn more!
- More example Diels-Alder reactions.



(a) Alkynes can be dienophiles.

(b) *cis/trans* dienophiles and *s-cis/s-trans* dienes.

(c) Multistep one-pot inverse electron-demand Diels-Alder.

Figure 2.31: Exotic Diels-Alder reactions.

1. Figure 2.31a: Dienophiles can be triple-bonded as well!
  - If a given molecule has at least one double bond, it can (usually) react as a dienophile.
2. Figure 2.31b: Don't be fooled by *s-trans* drawing; the diene is still buta-1,3-diene!
  - We will form the *cis*-product because the alkene hydrogens in the dienophile reactant are locked in the *cis*-orientation. (See Figure 2.23 for more context.)

3. Figure 2.31c: A cool example of an inverse electron-demand Diels-Alder reaction.
- The heteroatoms (nitrogens) in the ring of the diene should clue us into the fact that this might be an inverse electron-demand Diels-Alder reaction. (See the definition of inverse electron-demand Diels-Alder reaction.)
    - Indeed, this diene is called a **tetrazine**, and it can do inverse electron-demand reactions.
  - The diene within the tetrazine ring is highlighted in red, and the reactive alkene within the dienophile is highlighted in blue.
  - Drawing the product of the first Diels-Alder reaction can be a bit tricky, but Prof. Elkin has a good method for doing it.
    - Begin by stacking the starting materials in a perspective drawing.
    - Add dashed lines between the bonding atoms, yielding a drawing of your transition state.
    - Fill in the dashed lines and rearrange the double bonds to complete the transformation.
  - What’s cool about this reaction is that there is an immediate follow-up reaction to the first Diels-Alder.
    - In particular, the product of the first step does a retro-Diels-Alder, releasing  $\text{N}_2$  gas.
      - The release of an extremely stable gas molecule is a driving force for this second reaction.
    - Once  $\text{N}_2$  is released, we can redraw the product (now the final product) as a **diazine**.
- **Tetrazine**: A molecule with a central six-membered aromatic ring containing four nitrogen atoms.
  - **Diazine**: A molecule with a central six-membered aromatic ring containing two nitrogen atoms.