


- The Grignard reagent adds twice to carboxylic acid derivatives, yielding a tertiary alcohol.
  - This happens to acid chlorides, acid anhydrides, and esters.
  - Amides turn into the ketone (this is a special case!).
  - Carboxylates do not react.
- Organolithium reagents (more potent than Grignards) react exactly the same as Grignards, except that they will *also* turn carboxylates into ketones!
  - This is a very surprising result, since we've talked about how unreactive carboxylates are.
- Where do ketones and aldehydes fit into the picture?
  - Ketones and aldehydes are between anhydrides and esters, and aldehydes are more reactive than ketones.
  - $\text{NaBH}_4$  will reduce ketones and aldehydes to the primary alcohol.
  - We'll talk about this more later.
- Next time: A mechanistic explanation for Table 4.2.

## 4.26 Acyl Transfer Reactions - 3 / Nitriles


11/6:

- Announcements.
  - Exam 3 is one week from today!
  - Lots of time today and Friday doing practice synthesis problems.
    - Friday's review will *not* involve a summary of the Unit 3-4 material; instead, Prof. Buchwald will send a synopsis in advance.
  - The more problems you work, the easier synthesis will become!!
  - Take advantage of the fact that we don't know *too* many reactions yet!
- Lecture 25 recap.
  - Evidence for a tetrahedral intermediate in acyl transfer reactions: Isotopic labeling studies.
  - Recall Table 4.2.
    - This gets back to what is key for synthesis: **Chemoselectivity**.
      - Example: Consider a molecule with an aldehyde and an amide. We can selectively reduce the aldehyde to the alcohol and not touch the amide if we reduce with  $\text{NaBH}_4$ .
      - This can be important in fancy molecules if we want to play with the **pharmacokinetics**.
    - Acid chlorides and anhydrides are *super* reactive.
    - Aldehydes and ketones get reduced by  $\text{NaBH}_4$ , too!
    - $\text{NaBH}_4$  is mild, while  $\text{LiAlH}_4$  is *violent*. If you throw LAH into water, you'll get a *violent* reaction.
    - Similarly, Grignards are more mild than alkyllithium reagents.
      - Amides and carboxylates can become asymmetric ketones!
- **Chemoselectivity**: Selectivity for certain functional groups in the presence of other functional groups.
- **Pharmacokinetics**: The speed with which a drug moves into, through, and out of the body.
  - We don't want drugs to go straight through our bodies; we want them to hang around for a bit and do their thing (e.g., reduce our headache, soothe our cough, etc.).
  - We don't want to have to take it 5 times per day, so we modify functional groups with chemoselective reactions to slow the pharmacokinetics.

- TTQ: Synthesize the molecule at left in Figure 4.45a — a simplified version of a recently proposed candidate for treating epilepsy — from the provided starting materials.



(a) The desired molecule and starting materials.



(b) Retrosynthetic pathway.

Figure 4.45: TTQ: Synthesis of a drug molecule.

– Like last time (see Figure 4.42), let's start by mapping out the carbons.

- Using color coding, we can identify which carbons in the starting materials become carbons in the products (Figure 4.45a).

- It then becomes clear that what we need to add is a carbon-nitrogen linkage.
  - This could come from a cyano group!
  - Carbon-nitrogen doesn't *always* mean we need a cyano group, but it often does.
- So thinking backwards, the desired molecule could have come from an imine.
  - In the forward direction, we'd use a reducing agent ( $\text{NaBH}_4$  or  $\text{LiAlH}_4$  and a water workup) to reduce the imine to the amine.
  - Next step: Transform the imine to an amine via reductive amination.
  - Next step: Transform the amine to the nitrile via LAH and a water workup.
  - Next step: Transform the cyano group to an aryl diazonium salt via a **Sandmeyer reaction** (i.e., with  $\text{CuCN}$ ).
  - Next step: Transform the aryl diazonium salt to the amine via HONO.
  - Final step: Transform the amine to the nitro group via reduction with  $\text{H}_2 / \text{Pt}$ ,  $\text{H}_2 / \text{Pd/C}$ , or  $\text{H}_2 / \text{Ni}$ . LAH and  $\text{H}_2\text{O}$  are not ideal here.
- **Sandmeyer reaction:** Any method of displacing an aryl diazonium salt with a nucleophile in the presence of catalytic copper (I) salts.
- Takeaway: A general strategy for synthesis problems.
  1. Identify matching fragments (mostly carbon fragments).
  2. Look for functional groups and disconnections.
- This concludes today's synthesis example; we now return to the chemistry of carboxylic acid derivatives.
- Lecture outline.
  5. Reactions with  $\text{NaBH}_4$ ,  $\text{LiAlH}_4$ ,  $\text{RMgBr}$ , and  $\text{RLi}$ .
  6. Chemistry of nitriles.
    - a. Formation.
    - b. Reactions.
- We begin by resuming Topic 5: Reactions with  $\text{NaBH}_4$ ,  $\text{LiAlH}_4$ ,  $\text{RMgBr}$ , and  $\text{RLi}$ .
  - Specifically, we'll give the mechanistic explanation for Table 4.2 promised at the end of last lecture.
- Let's first consider why an acid chloride would react with hydride so quickly.




Figure 4.46: Reduction of acid chlorides mechanism.

- For the same reasons as with Figure 4.23, both addition to and elimination from an acid chloride is fast — it makes no difference that our nucleophile is a hydride!
- The first equivalent of hydride yields the aldehyde.
  - But we can't stop here!
  - Aldehydes are still reactive, so another equivalent of hydride will add in.
  - Then after a workup, we'll get the alcohol.
- Aside: Reagents exist that *can* convert an acid chloride to an aldehyde and stop there.

- Let's now look at the addition of a Grignard to an acid anhydride.




Figure 4.47: Grignard addition to acid anhydrides mechanism.

- The Grignard ( $R'-MgBr$ ) adds fast because acid anhydrides are not very resonance-stabilized either.
  - Then a good leaving group leaves to give a ketone.
  - Then the ketone reacts again to give us the tertiary alcohol.
- But the carboxylate is still hanging around.
  - It will *not* react with a Grignard.
  - Thus, we get 50% of  $3^\circ$  alcohol and 50% carboxylate, so this is *not* an elegant reaction.
- If we use  $R'Li$  instead of  $R'MgBr$ , this gives us 100% of the  $3^\circ$  alcohol, so this *is* a good reaction.




Figure 4.48: Alkyllithium addition to acid anhydrides.

- $R'Li$  is necessary because alkyllithium reagents are strong enough to reduce carboxylates, too (see Table 4.2).
- Note that this reaction only proceeds with heating.
- This reaction will *not* be tested!!
- With alkyllithium reagents, we can stop the reaction at the dianion and then quench.




Figure 4.49: Alkyllithium addition to carboxylates.

- To quench, use either water or  $H_3O^+$ .
  - You should write one of these two reagents above the arrow on a test, not "quench."
- This gives us the ketone hydrate.
  - But ketone hydrates are not stable, so under workup, we'll lose  $H_2O$  and obtain the ketone.
- This is the money reaction, and very much could be tested!!

- TTQ: How would you make a ketone from RLi and R'Li?




Figure 4.50: TTQ: Applying the addition of alkyllithium reagents to carboxylates.

- First step: Transform the ketone into the carboxylate and R'Li via the reaction in Figure 4.49.
- Second step: Transform the carboxylate into RLi via CO<sub>2</sub> carboxylation.
- Both LAH and R'M (M = MgBr, Li) can do add twice to esters.




Figure 4.51: Organometallic addition to esters mechanism.

- As we've seen before, the first step is addition to an electrophilic carbon center.
  - The resultant alkoxide anion is so powerful it can even push out a methoxide.
  - Then you get another addition to form the tertiary alcohol, after workup.
- Takeaway: If you see a tertiary alcohol with two like substituents, get used to thinking that it might come from the reaction of two equivalents of a Grignard (or alkyllithium reagent) with an ester!
- We now discuss two other reactions to make ketones.
  - These take acyl derivatives — “acyl X” — to ketones.
- Reaction #1: Beginning with an acid chloride.




Figure 4.52: Monoaddition to acid chlorides with dimethylcopper lithium.

- If we introduce a Grignard or alkyllithium reagent, the reaction will proceed all the way to the tertiary alcohol.
  - Thus, we need a gentler, more selective version of a Grignard or alkyllithium.
  - An example of such a reagent is **dimethylcopper lithium**.
- TTQ: Given the reaction above (except for the starting material, reagent, or product), fill in the missing compound.

- **Dimethylcopper lithium:** A reagent composed of an anionic copper atom covalently bonded to two methyl groups and ionically bonded to a lithium cation. *Also known as Gilman reagent, organocuprate. Structure*




Figure 4.53: Dimethylcopper lithium.

- History: Invented by Henry Gilman, an organic chemist at Iowa State University.
- Aside: This compound is really good at 1,4-addition, also known as conjugate addition. We'll cover such this class of reactions in Unit 5.
- Synthesis (not testable material):  $2 \text{ MeLi} + \text{CuX} \longrightarrow \text{Me}_2\text{CuLi}$
- Reaction #2: Beginning with a **Weinreb amide**.




Figure 4.54: Weinreb ketone synthesis.

- This reaction works with either Grignards or alkyllithium reagents.
- After addition to the carbonyl, the metal coordinates to both the *N*-oxygen's lone pair and the alkoxy anion.
  - This is a quasi-stable species.
- Water-workup then gets you the ketone.
- **Weinreb amide:** An amide with an *N*-methyl and *N*-methoxy group. *Structure*






Figure 4.55: Weinreb amide.

- We now move onto Topic 6: The chemistry of nitriles.
- We'll begin with Subtopic 6.a: Formation of nitriles.
  - We'll start with three reactions you already know:  $\text{S}_{\text{N}}2$  displacement, cyanohydrin formation, and the Sandmeyer reaction.
  - Then we'll cover one new method.
- $\text{S}_{\text{N}}2$  displacement (see Figure 3.21).
  - The X group can be Br, I, or OTs.

- Cyanohydrin formation (see Figure 3.22).
  - This reaction should be familiar from 5.12.
  - Note that the base catalyst usually has  $pK_a \approx 9.5$ .
- Sandmeyer reaction (see Figure 4.45b).
- One new method: Dehydration of amides.



(a) The reaction.



(b) A partial mechanism.

Figure 4.56: Nitrile synthesis: Dehydration of amides.

- We add  $\text{POCl}_3$  (the triacid chloride of phosphorous acid) to our amide (Figure 4.56).
  - $\text{POCl}_3$  is a very strong Lewis acid.
  - It rips out an equivalent of  $\text{H}_2\text{O}$  from our amide in a process known as dehydration.
- Approximate mechanism (Figure 4.56b).
  - $\text{POCl}_3$  is a strong Lewis acid, so it will head straight for one of the carbonyl lone pairs. The amide lone pair can then kick up to allow proper  $\text{O}-\text{P}$  bond formation, and kick out a  $\text{Cl}^-$ .
  - Following deprotonation of the amide, we obtain an intermediate with a great leaving group. The new nitrogen lone pair can then kick out this leaving group, which will also lose another  $\text{Cl}^-$  to enable  $\text{O}=\text{P}$  bond formation.
  - A final deprotonation gives us our nitrile.
- We now move onto Subtopic 6.b: Reactions of nitriles.
- Nitrile hydrolysis (see Figure 4.8).
  - Adding a harsh acid or base gets you all the way to the carboxylic acid.
  - Adding a mild acid or base gets you the amide.
  - We will not ask you either set of conditions on an exam!!
- Converting nitriles to ketones.






Figure 4.57: Organometallic addition to nitriles mechanism.

- Use an alkyllithium reagent or Grignard followed by an acidic workup.
- Implication: When you see a ketone in a molecule you're trying to synthesize, you can now think about whether it would be helpful if this retrosynthetically came from a nitrile and organometallic reagent, too!
- TTQ: Synthesize the molecule at left in Figure 4.58a from the provided starting materials.



(a) The desired molecule and starting materials.



(b) Solution.

Figure 4.58: TTQ: Applying nitrile addition chemistry.

- The cyclohexane to cyclohexyl bromide to Grignard reaction sequence in Figure 4.58b should be familiar from 5.12.
- Amide goes to nitrile with dehydration conditions ( $\text{POCl}_3$ ).
- Then the nitrile plus the Grignard makes the product.