

Homogeneous Catalysis

DOI: 10.1002/anie.200503479


A Versatile Method for Suzuki Cross-Coupling Reactions of Nitrogen Heterocycles**

Noriaki Kudo, Mauro Perseghini, and Gregory C. Fu*

The Suzuki cross-coupling reaction is a powerful method for carbon-carbon bond formation that has been applied in a variety of settings, ranging from natural-products synthesis to materials chemistry, including large-scale production.^[1,2] Among the attractive features of the Suzuki reaction are the wide availability, stability to air and moisture, and low toxicity of boronic acids, as well as the facile removal of the boron-containing side products of the coupling process.


In recent years, tremendous progress has been described in the discovery of more active catalysts for the Suzuki reaction. Arguably the most important remaining challenge is the development of a universal method for cross-coupling substrates that include nitrogen heterocycles.^[3] The presence of such groups, which are particularly pervasive in medicinal chemistry,^[4] can lead to low reactivity in coupling reactions. Herein, we describe an unusually versatile catalyst that

achieves Suzuki cross-couplings of an array of nitrogen-containing boronic acids and aryl halides [Eq. (1); dba = dibenzylideneacetone, Cy = cyclohexyl].



Since 1998, we and others have established that trialkylphosphines, including $\text{P}(t\text{Bu})_3$ and PCy_3 , serve as very effective ligands for a variety of palladium-catalyzed coupling processes.^[1,5] In view of the low cost of PCy_3 relative to many of the other ligands that provide highly active cross-coupling catalysts (e.g., $\text{P}(t\text{Bu})_3$, arylalkylphosphines, and carbenes), we decided to examine the possibility of developing a versatile PCy_3 -based catalyst for Suzuki reactions of nitrogen heterocycles.

In an earlier study, we employed $\text{Pd/PCy}_3/\text{KF}/\text{THF}$ as a catalyst system for Suzuki cross-couplings of aryl halides.^[6,7] Unfortunately, when we applied these conditions to the coupling of 3-pyridineboronic acid with 4-*n*-butylchlorobenzene, we obtained essentially none of the desired biaryl [Eq. (2)]. After considerable exploration of the reaction parameters, we determined that $\text{Pd/PCy}_3/\text{K}_3\text{PO}_4/\text{dioxane}/\text{H}_2\text{O}$ achieves the desired cross-coupling in excellent yield [Eq. (2)].^[8,9]



We were pleased to find that this standard protocol can be applied directly to Suzuki reactions of an array of nitrogen heterocycles. As illustrated in Table 1, a variety of heteroarylboronic acids can even be coupled with challenging aryl halides. Thus, 3-pyridineboronic acid reacts with a range of chlorobenzenes, including *ortho*-substituted and electronically deactivated substrates, in good yield (Table 1, entries 1–4). As might be anticipated, the corresponding aryl bromides and iodides also couple under these conditions (Table 1, entries 5 and 6). In addition, substituted 3-pyridineboronic acids are suitable cross-coupling partners (Table 1, entries 7–9). The method is effective not only for 3-pyridineboronic acids, but also for an array of other heteroarylboronic acids (e.g., 4-pyridine, 5-pyrimidine, and indole; Table 1, entries 10–13).

Although substrates that bear an indole NH group undergo Suzuki cross-coupling smoothly with this procedure (Table 1, entry 13), the catalyst appears to be sensitive to pyrazole NH groups (Table 1, entries 14 and 16). However, simple protection of the nitrogen atom leads to coupling with useful efficiency (Table 1, entries 15 and 17).^[10]

[*] Dr. N. Kudo, Dr. M. Perseghini, Prof. Dr. G. C. Fu
Department of Chemistry
Massachusetts Institute of Technology
Cambridge, MA 02139 (USA)
Fax: (+1) 617-324-3611
E-mail: gcf@mit.edu


[**] We thank Frontier Scientific for boronic acids and Johnson Matthey for $[\text{Pd}_2(\text{dba})_3]$. Financial support was provided by the National Institutes of Health (National Institute of General Medical Sciences, R01-GM62871), Merck Research Laboratories, Novartis, Sankyo-Agro Co. Ltd. (N.K.), and Novartis Stiftung (M.P.). Funding for the MIT Department of Chemistry Instrumentation Facility was furnished in part by the National Science Foundation (CHE-9808061 and DBI-9729592).



Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

(entries 13 and 14, respectively).^[11] Protection of the pyrazole NH group is necessary for the coupling of a 4-bromopyrazole with 3-pyridineboronic acid (entry 15 vs. 16; see also Table 1, entries 14–17).

In preliminary experiments, we determined that this Pd/PCy₃/K₃PO₄/dioxane/H₂O-based method is effective not only for Suzuki cross-couplings of heteroarylboronic acids, but also for boronate esters and trifluoroborates [Eq. (3)]. Finally, we



established that the procedure can be conducted on a multigram scale (18 mmol of 4-*n*-butylchlorobenzenechloride + 20 mmol of 3-pyridineboronic acid, yield = 3.4 g (88%) of product).

To the best of our knowledge, this is the most wide-ranging study that has been described for Suzuki reactions of nitrogen-containing cross-coupling partners. Attractive features of this method include its versatility (a single procedure was employed for all of the examples, including boronate esters and trifluoroborates), its compatibility with a variety of unprotected functionalities (e.g., NH₂- and OH-substituted pyridines and unprotected indoles), its convenience (commercially available components), and its efficiency even with inexpensive, unactivated aryl chlorides. We anticipate that this catalyst system will find application in academia and, in particular, in industry.

Experimental Section

General procedure: The heteroarylboronic acid (1.10 mmol), [Pd₂(dba)₃] (9.2 mg, 0.010 mmol), and PCy₃ (6.7 mg, 0.024 mmol) were added to a 25-mL Schlenk flask equipped with a stir bar in air. The flask was evacuated and refilled with argon five times. Dioxane (2.67 mL), the (hetero)aryl halide (1.00 mmol; if the halide was a solid, it was added prior to the evacuation/refill cycle), and aqueous K₃PO₄ (1.27 M, 1.33 mL, 1.70 mmol) were added by syringe. The Schlenk flask was sealed and heated in an oil bath at 100 °C for 18 h with vigorous stirring. The mixture was then filtered through a pad of silica gel (washing with EtOAc), the filtrate concentrated under reduced pressure, and the aqueous residue extracted three times with EtOAc. The combined extracts were dried over anhydrous MgSO₄, filtered, and concentrated. The residue was then purified by column chromatography on silica gel.

Received: October 1, 2005

Revised: December 12, 2005

Published online: January 19, 2006

Keywords: cross-coupling · heterocycles · homogeneous catalysis · nitrogen · palladium

- [1] For leading references, see: a) N. Miyaura in *Metal-Catalyzed Cross-Coupling Reactions* (Eds.: A. de Meijere, F. Diederich), Wiley-VCH, New York, **2004**, chap. 2; b) A. Suzuki in *Handbook of Organopalladium Chemistry for Organic Synthesis* (Ed.: E.-i. Negishi), Wiley Interscience, New York, **2002**, chap. III.2.2.
- [2] For an overview, see: A. M. Rouhi, *Chem. Eng. News* **2004**, 82, 49–58.
- [3] For recent reports that include Suzuki reactions of nitrogen heterocycles, see: a) T. E. Barder, S. D. Walker, J. R. Martinelli, S. L. Buchwald, *J. Am. Chem. Soc.* **2005**, 127, 4685–4696; b) T. E. Barder, S. L. Buchwald, *Org. Lett.* **2004**, 6, 2649–2652; c) I. Kondolff, H. Doucet, M. Santelli, *Synlett* **2005**, 2057–2061; d) A. E. Thompson, G. Hughes, A. S. Batsanov, M. R. Bryce, P. R. Parry, B. Tarbit, *J. Org. Chem.* **2005**, 70, 388–390, and references therein.
- [4] a) For a discussion of the widespread occurrence of nitrogen heterocycles, particularly pyridines, in pharmaceutically active compounds, see: V. Bonnet, F. Mongin, F. Trecourt, G. Breton, F. Marsais, P. Knochel, G. Queguiner, *Synlett* **2002**, 1008–1010 (footnote 1); b) A. F. Pozharskii, A. T. Soldartenko, A. Katritzky, *Heterocycles in Life and Society*, Wiley, New York, **1997**.
- [5] For an overview, see: J. Tsuji, *Palladium Reagents and Catalysts*, Wiley, New York, **2004**.
- [6] A. F. Littke, C. Dai, G. C. Fu, *J. Am. Chem. Soc.* **2000**, 122, 4020–4028; see, also: A. F. Littke, G. C. Fu, *Angew. Chem.* **1998**, 110, 3586–3587; *Angew. Chem. Int. Ed.* **1998**, 37, 3387–3388.
- [7] For an early report of the use of Pd/PCy₃ for palladium-catalyzed Suzuki reactions, see: W. Shen, *Tetrahedron Lett.* **1997**, 38, 5575–5578.
- [8] For an overview of the use of water as a (co)solvent for Suzuki reactions, see: N. E. Leadbeater, *Chem. Commun.* **2005**, 2881–2902.
- [9] For the application of a related system to the cross-coupling of cyclopropylboronic acid with (hetero)aryl bromides, see: D. J. Wallace, C.-y. Chen, *Tetrahedron Lett.* **2002**, 43, 6987–6990.
- [10] For the Suzuki cross-coupling illustrated in entry 15 of Table 1, deboronation is a significant side reaction.
- [11] For a previous example, see: C. N. Johnson, G. Stemp, N. Anand, S. C. Stephen, T. Gallagher, *Synlett* **1998**, 1025–1027.