

# Topic 5


## Exam 1

### 5.1 Exam 1 Review Sheet


- 3/4:      • Important heterocycles and their key properties.
- Pyridine.
    - $\pi$ -deficient ring:  $\delta^+$  on  $\alpha$ - and  $\gamma$  carbons,  $\delta^-$  at  $\beta$ -carbon.
    - Reactivity.
      - > EAS: Bad, except as *N*-oxide.
      - >  $S_NAr$ : Good, especially at the  $\alpha$ -carbons. Electrophile coordination can induce a 10 order of magnitude rate increase (see Figure 2.4), and can even drive dearomatization (as in  $NAD^+/NADH$ ).
    - Acidity:  $\gamma > \beta > \alpha$  ( $\delta^+$  without  $\alpha$ -effect, no  $\alpha$ -effect,  $\alpha$ -effect).
    - Basicity:  $pK_a \approx 5.5$  (modulated by substituents).
    - Nucleophilicity: Modulated by substituents (e.g., pyridine vs. DMAP).
    - Pyridine-containing drug: Nicotine.
  - Pyridone.
  - Quinoline.
    - Quinoline-containing drug: Quinine.
  - Isoquinoline.
    - Reactivity.
      - > More reactive on non-heterocyclic portion.
      - > EAS at 5- and 8-positions.
      - >  $S_NAr$  *always* at 1-position.
  - Quinolone.
    - Quinolone-containing drug: Ciprofloxacin.
  - Pyridazine.
    - Reactivity: Easier to protonate because of unfavorable  $\alpha$ -effect (in neutral form).
  - Pyrimidine.
    - Reactivity.
      - > Relative to pyridine: Better at  $S_NAr$ , worse at EAS.
      - > More reactive at 4- than 2-position (double  $\alpha$ -effect is bad).
    - Pyrimidine-containing drug: Anti-asthma agents.
  - Pyrrole.

- Protonation at  $\alpha$ -carbons ( $\text{pK}_a = -3.8$ ).
- $\pi$ -excessive ring: Slightly more reactive toward EAS at  $\alpha$ - than  $\beta$ -carbons, though can vary depending on the type of carbocation formed??
- Pyrrole-containing drug: Lipitor.
- Imidazole.
  - Hydrogen-bonds well.
  - Undergoes tautomerization.
  - $\text{pK}_{a1} = 14.5$ ,  $\text{pK}_{a2} = 7.1$ .
    - Oxazole's  $\text{pK}_a = 0.8$ , thiazole's  $\text{pK}_a = 2.5$  (no equal-energy resonance form).
  - Reactivity: Good at EAS (but not as good as pyrrole).
  - Imidazole-containing natural product: Histidine.
- Pyrazole.
  - Dimeric in solution (due to hydrogen bonding).
  - Tautomerization: Hydrogen prefers to be farther away from bulky substituents.
  - Reactivity.
    - EAS at 4-position (halogenation, formylation, etc.)
  - Pyrazole-containing drug: DGAT-2 inhibitors.
- Indole.
  - Reactivity.
    - 5-membered ring is most reactive.
    - EAS at 3-position.
    - Basic conditions can make N more nucleophilic than C3 (e.g., for acylation).
    - Alkylation: C3, C3  $\rightarrow$  C2, C3  $\rightarrow$  deprotonation, N.
    - Deprotonation at C2 (esp. with Boc DMG).
  - $\text{pK}_a = 16.2$  for the nitrogen proton.
  - Indole-containing natural product: Strychnine, tryptophan.
- Indazole.
  - Indazole-containing drug: EGFR inhibitor.
- Thiophene.
  - Thiophene-containing natural product: Echinothiophene.
- Furan.
  - Acidity:  $\text{pK}_a \approx 35.6$  ( $\alpha$ -carbons).
  - Furan-containing drug: Zantac.
- Pyridine reactivity.
  - Directed metallation.
    - Lithiation is reversible, hence why it is observed as occurring *thermodynamically* at the  $\gamma$ -position over *kinetically* at the  $\alpha$ -position.
    - DMGs: All the usual suspects ( $3^\circ$  amides, methoxy, carbamates, etc.).
    - DMGs (for  $\pi$ -deficient heterocycles): Halogens and pseudo-halogens (F, Cl,  $\text{CF}_3$ ,  $\text{CO}_2^-$ ).
      - Br sometimes included, but may prefer to do lithium/halogen exchange.
      - MeO is stronger than Cl as a DMG.
    - $\beta$ -DMGs direct  $\gamma$  (almost all) or  $\alpha$  ( $-\text{OEt}$ ).
    - $\gamma$ -DMGs direct  $\beta$ .
    - $\alpha$ -DMGs direct  $\beta$ .

- Remember to do these reactions cold, in ethereal solvent (Et<sub>2</sub>O or THF), and maybe with an additive (e.g., TMEDA).
- Lithium/halogen exchange.
- Lateral deprotonation.
  - pK<sub>a</sub>'s:
    - > γ: 26.
    - > α: 29.5.
    - > β: 33.5.
  - Thermodynamic conditions: γ > α > β-positions.
  - Kinetic conditions (e.g., with <sup>n</sup>BuLi): α > γ > β-positions.
  - Connection to decarboxylation at lateral positions.
- S<sub>N</sub>Ar is more probable than ketone addition.
  - For example, aqueous ammonia will do S<sub>N</sub>Ar on pyridine before it adds to a bonded ketone.
- Hydrogenolysis of aryl chlorides.




- Aryl chloride to methyl group.




- See Figure 1.5.


- Chichibabin (reaction).




- 2-addition.




- The Grignard must be aryl, vinyl, or alkynyl.
- It's not clear what the final step oxidant would be, but perhaps DDQ??
- N-oxide formation and removal.




- Oxidants include *m*CPBA and H<sub>2</sub>O<sub>2</sub>.
- N-oxide nitration.




- *N*-oxide bromination.




- *N*-oxide chlorination.




- **Zincke** (reaction).



- **Zincke** (aldehyde formation).




- Milder, Zincke-inspired *meta*-halogenation.



■ Also works with NIS.


■ Works with substituted pyridines, too.

- *meta*-halogenation via dearomatization.




• Pyridine synthesis.


- Industrial pyridine synthesis.




- 1,5-dicarbonyl pyridine synthesis.




- **Hantzsch** (pyridine synthesis).




- Mechanism is classic condensation reactions.
- Asymmetric variant: Condense aldehyde and 1,3-dicarbonyl first, then condense with a **vinyligous urethane**. No last-step oxidation needed.
- **Kröhnke** (pyridine synthesis).




- If R is enolizable (and not methyl), you will get regioisomers.
- Can also start directly with an  $\alpha$ -bromocarbonyl compound.
- [2 + 2 + 2] pyridine syntheses: Cool, but limited synthetic utility.
- Some more important  $\text{pK}_a$ 's.
  - $^n\text{BuLi}$ : 50.
  - LDA: 36.
  - $\text{LiNEt}_2$ : 31.7.
- Pyridone.
  - Chlorination (see Figure 1.4).



- Cross-coupling.
  - Know generic mechanism.
    - Transmetalation typically occurs through  $\sigma$ -bond metathesis.
  - **Kumada** (coupling).




- **Negishi** (coupling).




- Common solvent: THF.

- Ideal for coupling something to the pyridine  $\alpha$ -position; 2-pyridylzincs are great.

- **Stille** (coupling).



- **Suzuki-Miyaura** (coupling).



- Common base:  $\text{K}_2\text{CO}_3$ .

- Common ligands.


- SPhos (see Figure 5.1b).

- dppf ( $sp^3$ -hybridized boronates).


- Heteroaryl couplings:  $\text{PCy}_3$  or MIDA boronates.

- Common solvent: ACN/ $\text{H}_2\text{O}$ .


- **Sonogashira** (coupling).




- **Hiyama** (coupling).




- **Carbonyl enolate** (coupling).



- **Heck** (coupling).



- **Buchwald-Hartwig** (amination).




- **Carbonylation**.




- Via an acyl palladide ( $\text{ArCOPd}$ ) intermediate.


- **Ullmann** (coupling).




- **Goldberg** (coupling).




- **Miyaura** (borylation).



– **Cyanation.**



– **C–O (coupling).**



■ R can be alkyl or aryl.

– **Ligands.**




Figure 5.1: Dialkylbiaryl phosphine ligands.


■  $\text{PPh}_3$ .

■ Xantphos (Buchwald-Hartwig amination).

■ SPhos (Miyaura borylation and Suzuki-Miyaura borylation).


■ RuPhos (Negishi coupling).

– Direct cross-coupling of two (possibly hetero)aryl halides.




• **Quinoline reactivity.**

– *meta*-bromination.



■ Proceeds through alternate mechanism.

– 2-addition.



– **Hydrogenations.**




Figure 5.2: Quinoline hydrogenations.

- *cis*-decalin mostly formed in complete hydrogenation; some *trans*-though.
- Quinoline synthesis.
  - **Meth-Cohn** (quinoline synthesis): 3-substituted and 2-substitutable quinolines.



- The starting material could come from the (possibly substituted) aniline and acid chloride (plus  $\text{NEt}_3$ ).
- Mechanism: Amide  $\rightarrow$  chloroimine  $\rightarrow$  enamine  $\rightarrow$  attack on a Vilsmeier reagent  $\rightarrow$  Friedel-Crafts.
- **Skraup** (quinoline synthesis): No substitution, mix of (di-)2- and 4-substitution.






Figure 5.3: Skraup quinoline synthesis.


- Figure 5.3a.
  - If at least one of  $\text{R}^2, \text{R}^3 = \text{H}$ , then acid =  $\text{H}_2\text{SO}_4$  and the system is oxidized to a quinoline.




- If both  $R^2, R^3 \neq H$ , then acid = pTsOH and the system is *not* oxidized.
- Figure 5.3b.
  - Acrolein generated *in situ* from glycerol.
  - Mechanism: Michael addition, Friedel-Crafts, dehydration, oxidation.
- **Friedlander** (quinoline synthesis): 2-, 3-, and 4-substitution (or mix and match).



- Mechanism: Imine condensation, followed by enamine attack on the aldehyde/ketone.
- Regioisomer problems: Just reject the unwanted side product.
- Quinolone synthesis.
  - **Conrad-Limpach-Knorr** (quinolone synthesis).




(a) 2-quinolones.




(b) 4-quinolones.

Figure 5.4: Conrad-Limpach-Knorr quinolone synthesis.


- Presumably also works with substituted variants.
- Acid protonates the more electron-rich ester; heat provides energy for attack at the more electrophilic ketone.
- Isoquinoline reactivity.
  - 5-bromination.




- 5- and 8-nitration.




- Chichibabin reaction.



- 1-addition.




- 1-selective  $S_NAr$ .



- Isoquinoline synthesis.


- **Pomeranz-Fritsch** (isoquinoline synthesis): Anything can be substituted.



- Can use methyl or ethyl acetal.

- **Bischler-Napieralski** (isoquinoline synthesis): Enables formation of same derivatives.
- **Pictet-Gams** (isoquinoline synthesis): Enables formation of same derivatives.


- **Pictet-Spengler** (reaction): A  $\beta$ -arylethylamine undergoes condensation with an aldehyde or ketone followed by ring closure.



- Mechanism can be Friedel-Crafts or involve shifts (depending on the most nucleophilic position).


- Pyrimidine synthesis.

- **Grimmaux** (pyrimidine synthesis): 3 carbonyls.




- Can also use NaOR/ROH and di-R malonate esters.


- **Ziegenbein-Franke** (pyrimidine synthesis): 5- and 6-substituted pyrimidines.




- Will have regioselectivity issues if  $\text{R}^1 \neq \text{R}^2$ .
- [3 + 3] (pyrimidine syntheses).
  - Bis-nucleophile: Pinner product, or other group in the middle besides alkyl.
  - Bis-electrophile.
    - 4(5)6-substitution:  $\beta$ -diketone.
    - (5)6-substitution, 4-one:  $\beta$ -ketoester.
    - 4(5)-substitution:  $\alpha, \beta$ -unsaturated ketone with  $\beta$ -leaving group.
    - 6-substitution, 4-one: Propynyl ester.
    - (5)-substitution: Vinamidium salt.
    - (5)6-substitution, 4-amine:  $\beta$ -ketonitrile.
- **Pinner** (reaction).



- Forms **Pinner salts**, which are readily derivatized.
- **Turbogrignard**: The compound  $^i\text{PrMgCl}$ , which is useful for converting  $\text{R}-\text{X}$  to Grignards.
- $\text{KOH} + \text{H}_2\text{O}/\text{THF}$  can sometimes be used to convert chloro-heterocycles to carbonyl groups.
- Boc protection/deprotection.




- Pyrrole reactivity.
  - 2- (and partial 3-) nitration.




- $\text{AcONO}_2$  is a source of  $\text{NO}_2^+$ ; made from  $\text{HNO}_3 + \text{Ac}_2\text{O}$ .
- Perbromination.

- 2,5-bromination.




■ Boc-protection.

- 3-bromination.




■ TIPS-protection.

- 3,4-bromination.




- 2-bromination.




■ Dibromodimethylhydantoin is an alternative  $\text{Br}^+$  equivalent.

- Vilsmeier formylation.




- Can get reactivity at N by deprotonating with  $\text{NaH}$ .
- Diels-Alder reactivity: Possible with Boc (EWG) protection and *very* activated dienophiles.
- Decarboxylation.




■ Carboxylic acids can be used as removable C2-blocking groups.

- Cross-coupling.




■ Big, bulky protecting group needed (TIPS best).

– 2,5-dimethylpyrrole protection/deprotection.




– 2-nitrilation (with CSI).




■ DMF-induced pericyclic reactions can help in workup.

– 2-nitrilation (with Vilsmeier-type chemistry).




• Pyrrole synthesis.

– Industrial pyrrole synthesis.




– **Paal-Knorr** (pyrrole synthesis): (1)(2)(5)-substitution.




■ To avoid  $R^3$ , use  $NH_3$ .

■ To avoid  $R^1$ ,  $R^2$ , or both, use the corresponding acetal(s). 2,5-dimethoxyTHF may be useful.


– **Knorr** (pyrrole synthesis): (2)3-substitution; can keep an ester or carboxylic acid at the 4- and/or 5-position.



- **Hantzsch** (pyrrole synthesis): 2,3,5-substitution.




- **van Leusen** (pyrrole synthesis): 2,3,4-substitution.




■ α,β-unsaturated SM can come from HWE!


- POBr<sub>3</sub> does the same thing as POCl<sub>3</sub> (e.g., can brominate something).
- Imidazole reactivity.
  - Alkylation under neutral conditions: MeI adds to one, both, or neither nitrogen.
  - Alkylation under basic conditions (LDA, NaH, NaHMDS): Deprotonation and alkylation.
  - Selective N<sup>2</sup> alkylation.



- 4-nitration.




- Perbromination.



■ Can also do 2-bromination with just Br<sub>2</sub>??


- Directed metallation.




■ Protect with SEM (deprotonation, SEM-Cl).

■ Direct to C2, which can also be protected/deprotected to direct C4.


- Lithium/halogen exchange: Consider adding a strong base before <sup>n</sup>BuLi to ensure ordering.
- **Minisci** (reaction).



- Radical addition to electrophilic sites.




- Quaternary imidazolium salts to *N*-heterocyclic carbenes.




- Imidazole synthesis.


- **Debus-Radziszewski** (imidazole synthesis): (2)4(5)-substitution.




- Pinner-type (imidazole synthesis): (2)4(5)-substitution.




- **van Leusen** (imidazole synthesis): (3)4-substitution.



- Synthesis 4: 2(3)-substitution.




- Paal-Knorr-type (imidazole synthesis): 24(5)-substitution.




- Pyrazole reactivity.

- Acylation.



- Mechanism probably proceeds through reversible acylation at the other nitrogen first.


- 4-halogenation.



- N-alkylation varies in neutral vs. basic conditions as in imidazole.
- 5-directed metallation upon N-H protection.


- Pyrazole synthesis.

- **Knorr** (pyrazole synthesis): (2)35-substitution.



- Regioisomer issues if asymmetric, unless extreme mismatch in electrophilicity/nucleophilicity is induced.


- Dipolar cycloaddition method: 34-substitution.



- May have regioisomer issues. Can be partially overcome by introducing electronic biases.

- Indole reactivity.

- **Mannich** (reaction): Formaldehyde and dialkylamines add at C3.
- C2 lithiation.




- Gramine (from Mannich reaction) can be methylated and leave to allow other nucleophiles to attach to the offshot position.
- 4-lithiation of gramine with TIPS protection.


- Indole synthesis.

- **Zincke** (indole synthesis): 3-substitution.






- **Bartoli** (indole synthesis): (2)(3)7-substitution, and other on the benzene ring.




- Requires bulky 7-group.

- **Leimgruber-Batcho** (indole synthesis): (3)-substitution, and other on the benzene ring.




- Does not need a bulky 7-group.

- **Bischler** (indole synthesis): Aniline starting material.




- **Fischer** (indole synthesis).




- Regioisomer problems: Enolization both ways, substituents on the ring.
- *meta*-EDG selective for 6-substitution.
- Weak acid selective for thermodynamic enolization; strong acid selective for kinetic enolization.

- **Reissert** (indole synthesis): 2-ester-(3)-substitution, and other on the benzene ring.




- **Madelung** (indole synthesis): 2(3)-substitution, and other on the benzene ring.



- Could prepare starting material from Fridel-Crafts, Clemmensen, bromination, Goldberg (or nitration, reduction, acylation).


– **Hemetsberger** (indole synthesis): 2-ester-substitution.

- *para*-sulfonyl protecting group installation and removal.




- Indazole reactivity.

– N<sup>1</sup>- and N<sup>2</sup>-THP protection.



- Deprotect with pTsOH in MeOH.


– 3-halogenation.




- Feeds into cross-coupling.

- Indazole synthesis.

– Route 1: (3)-substitution, and other on the benzene ring.




– Route 3: (3)-substitution, and other on the benzene ring.




- Can put nothing (aldehyde), amine (nitrile), or hydroxyl (ester) on the 3-position.

- Thiophene reactivity.


– Perbromination.




– 2-bromination/chlorination.




- 2-iodination.




- 2,3,5-tribromination.



- 3-bromination.




- 2,3-dibromination.




- Thiophene synthesis.


- Industrial thiophene synthesis.



- **Paal-Knorr** (thiophene synthesis): 2,5-substitution.




- **Fiesselmann** (thiophene synthesis): 2-ester-4,5-substitution.




■ Can also use esters or nitriles as in indazole route 1.

■ Can saponify ester to 2,3-substituted derivative.

- **Hinsberg** (thiophene synthesis): 3,4-substitution.




- **Gewald** (thiophene synthesis): 2-amino-3-EWG-4,5-substitution.




- Knoevenagel-type mechanism.
- Remember that  $S_1$  is more active than  $S_n$ .

• Furan reactivity.


- 2-bromination.



- 2-addition.




- Diels-Alder with highly activated dienophiles.
- Mannich reaction: 2-substitution.
- 3-addition.




• Furan synthesis.

- **Paal-Knorr** (furan synthesis): 2(3)(4)5-substitution.



- **Feist-Benary** (furan synthesis - aldehydes): 23-substitution.



- **Feist-Benary** (furan synthesis - ketones): 235-substitution.

