

A General, Multimetallic Cross-Ullmann Biheteroaryl Synthesis from Heteroaryl Halides and Heteroaryl Triflates

Kai Kang, Nathan L. Loud, Tarah A. DiBenedetto, and Daniel J. Weix*

Cite This: *J. Am. Chem. Soc.* 2021, 143, 21484–21491

Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: Despite their importance to medicine and materials science, the synthesis of biheteroaryls by cross-coupling remains challenging. We describe here a new, general approach to biheteroaryls: the Ni- and Pd-catalyzed multimetallic cross-Ullmann coupling of heteroaryl halides with triflates. An array of 5-membered, 6-membered, and fused heteroaryl bromides and chlorides, as well as aryl triflates derived from heterocyclic phenols, proved to be viable substrates in this reaction (62 examples, $63 \pm 17\%$ average yield). The generality of this approach to biheteroaryls was further demonstrated in 96-well plate format at 10 μmol scale. An array of 96 possible products provided $>90\%$ hit rate under a single set of conditions. Further, low-yielding combinations could be rapidly optimized with a single “Toolbox Plate” of ligands, additives, and reductants.

Biheteroaryls are commonly found in drugs, electronic materials, ligands, and natural products,¹ but their synthesis remains a challenge.² As all-carbon aromatics are more widely available and have a more discrete range of reactivity than their heteroaryl analogues, strategies toward biheteroaryl synthesis have had to balance substrate availability and stability with reaction generality (Scheme 1). For example, cross-coupling approaches³ are the most general, but can be limited by the poor availability^{4,5} and instability⁶ of heteroaryl nucleophile equivalents. C–H arylation⁷ and cross-dehydrogenative coupling⁸ approaches overcome substrate availability problems, but C–H bond selectivity and generality among the broad reactivity of heteroaryls can be challenging. Recently, sulfur^{9,10} and phosphorus¹¹ chemistry has been developed that allows access to challenging biheteroaryls, but these new approaches share some of the availability and regiocontrol limitations of cross-coupling and C–H functionalization. Given the wide availability⁴ of heteroaromatic bromides, chlorides, and phenols (10–1000 times more than HetAr-[M]), a cross-Ullmann approach would be a powerful addition to biheteroaryl synthesis.

Cross-Ullmann biaryl syntheses can be divided into two categories: 1-pot, 2-step reactions (e.g., borylation of one HetAr-X followed by coupling with a second HetAr-X) and reductive single-step reactions. While both approaches have had success with biaryl^{12,13} and heteroaryl–aryl^{14,15} cross-coupling reactions, neither approach has been widely tested against biheteroaryls.^{16,17} Indeed, the wide range of reactivities (that present a challenge for selectivity) and the potential for biheteroaryls to coordinate strongly to the catalyst¹⁸ (inhibiting turnover and lowering selectivity) could limit cross-Ullmann biheteroaryl synthesis to a narrow range of coupling partners.¹⁹ Our approach was to use our increasing understanding of Ni and Pd multimetallic cross-Ullmann chemistry²⁰ to design a comprehensive solution to the synthesis of biheteroaryls from heteroaryl halides and heteroaryl triflates (Scheme 1). Our results, described here provide a general set of conditions effective for a broad array of

combinations and the tools necessary to optimize low-yielding reactions.

The initial reaction optimization^{20a} using 6-methylpyridin-3-yl triflate (**1a**) and 3-bromopyridine (**2a**) as starting materials (Table 1 and Table S6 in Supporting Information) revealed that nonsymmetric biheteroaryl **3a** could be formed in 50% yield in the presence of 5 mol % $\text{NiBr}_2(\text{dme})/\text{N1}$ and 5 mol % $\text{PdCl}_2/\text{P1}$ as catalysts, 2 equiv of Zn as reductant, and 2 equiv of KF as additive in DMF at 60 °C (Table 1, entry 1). A series of control reactions confirmed the necessity of metal catalysts, reductant, and salt additive.^{20,21} Replacement of the highly hygroscopic²² KF additive with KBr did not lead to any decrease in yield of **3a** (entry 2). Further screening of the amine ligands on nickel showed that, while dtbbpy (**N1**) remained the most promising for the model substrate pair, 1,10-phenanthroline (**N2**) and substituted terpyridine ligands (**N3** and **N4**) were also relatively effective (entries 3–5). An evaluation of phosphine ligands indicated that product **3a** could be obtained in satisfactory yields when **P2** and **P3** were used (entries 6 and 7), although **P1** was still preferred when the amount of **2a** was increased to 1.5 equiv (73 vs 70%, entries 2 and 6). Finally, the use of rigorously anhydrous KBr (stored in a N_2 filled glovebox) further improved the yield of **3a** to 79% (Table 1, entry 8).


An array of 6-membered, 5-membered, and fused heteroaryl bromides were next evaluated as substrates under these optimal conditions (Scheme 2).²³ In addition to 3-pyridyl bromides, a 2-pyridyl bromide (**3k**) and a substituted 4-pyridyl bromide (**3l**) could both be coupled with 3-pyridyl triflates

Received: October 15, 2021

Published: December 17, 2021




Scheme 1. Approaches to Access Nonsymmetric Biheteroaryls by Merging Heteroarene Feedstocks



successfully, affording the biheteroaryls in 58% and 83% yield. These conditions were also effective for a variety of 5-membered heteroaryl bromides (**3m–3w**), including a number that are much more electron-rich than pyridines (**3m–3o**, **3r–3w**), and a diverse array of fused ring heterocycles (**3x–3aj**). In all except three cases,²⁴ lower yields were due to formation of side products (homodimers and HetAr–H) or challenges in isolation.²⁵ In three of the lower-yielding cases, higher yields could be obtained with further optimization (see Figure 1).

Although heteroaryl chlorides are the most widely available heteroaryl electrophile,⁴ their low reactivity can present challenges to selectivity in cross-electrophile coupling.^{20c} Our standard conditions (A) allow the coupling of a variety of heteroaryl chlorides, ranging from those activated toward oxidative addition (**3al–3ao**) to deactivated (**3ap–3ar**). The wider availability of aryl chlorides provides unique coupling partners, such as precursors to drug molecules imiquimod²⁶ and pazopanib²⁷ (**3as** and **3at**).^{28,29} Finally, 2,3'-bipyridine **3ao** was scaled 10-fold, from 0.5 to 5 mmol, with comparable yields (71% to 63%).

These general conditions tolerate a variety of electrophilic functional groups that could be challenging for other methods. For example, ketones, esters, and aldehydes could react with Grignard reagents, but are unreactive under these conditions (**3g**, **3h**, **3p**, **3q**, **3an**, **3ao**). A wide variety of protecting groups are tolerated, including acid-sensitive protecting groups (**3m**, **3s**, **3t**, **3x**, **3ah**) and even an allyl group is tolerated (**3ag**).

Table 1. Optimization of Model Reaction^a

entry	variations from above conditions	3a (%)
1	KF (2.0 equiv) as additive	50
2	none	50 (73 ^b)
3	N2 instead of N1	40
4	N3 instead of N1	43
5	N4 instead of N1	42
6	P2 instead of P1	52 (70 ^b)
7	P3 instead of P1	45
8	rigorously anhydrous KBr	79 ^b (81 ^c)

^aReactions on a 0.25 mmol scale in 0.75 mL of DMF. Calibrated GC yield. KBr was stored in air unless otherwise noted. See Supporting Information, Table S6 for additional selectivity data. ^b1.5 equiv of 2a was used. ^c¹H NMR yield.


Finally, some steric hindrance is tolerated (**3e**, **3o**, **3u**, **3w**, **3x-aa**, **3ac**, **3ae**).

While electron-neutral and electron-rich pyridyl and pyrimidyl bromides proved to be viable substrates (**3a–3e**), we found that couplings with electron-poor heteroaryl bromides (**3f–3j**) led to low yields due to rapid homocoupling. Re-examination of the other effective ligands in Table 1 led to a new set of conditions (B) utilizing ttbtpy (**N3**) instead of dtbbpy (**N1**) that suppressed competitive homocoupling and delivered the products **3f–3j** in 50–75% yields.³⁰

The scope of heteroaryl triflates was also found to be broad (Scheme 3). In addition to the pyridyl triflates substituted with electron donating groups used in Scheme 2, pyridyl triflates with electron-neutral (–H, **3au**) or -deficient (–CO₂Me, **3aw**) substituents could be coupled, as could more electron-rich benzothiazole (**3ax**).

More importantly, these conditions can be used to synthesize challenging chelating biheteroaryls with varying electronic and steric properties (**3aac–3aai**) that could be useful in medicinal chemistry or as nonsymmetric bidentate ligands.³¹ While these products should be competitive ligands for both Ni and Pd, reactions were complete by 24 h.³² Notably, products **3aae** and **3aai** feature biheteroaryl cores that have not been previously reported.³³ Finally, the conditions could be extended to a nonaromatic vinyl triflate (**3aaj**).

To gain additional information on the applicability of these conditions to medicinal chemistry, we adapted our chemistry to a standard high-throughput experimentation (HTE) setup. Modern medicinal chemistry often utilizes HTE to facilitate the synthesis of collections of molecules for biological

Scheme 2. Scope of Heteroaryl Bromides and Chlorides Coupled with Heteroaryl Triflates^a

^aIsolated yields of reactions on a 0.5 mmol scale. Reactions were conducted under conditions A unless otherwise noted. ^bNMR yield of a mixture of product and side products, see [Supporting Information](#) for details. ^cReactions were conducted under conditions B. ^dReaction temperature was 80 °C. ^eIsolated yield of a 5.0 mmol scale reaction.





Figure 1. (A) General conditions used for HTE scope investigation (E). (B) Heteroaryl triflates used in HTE scope investigation (E). (C) Heteroaryl halides used in HTE scope investigation (E). (D) Ligands, salt additives, and reductants used in optimization Toolbox Plates (F–H). (E) Heatmap of results of HTE scope (using conditions from A and substrates from B and C). (F) Heatmap of results of product B3 reoptimization (using toolbox from D). (G) Heatmap of results of product C6 reoptimization (using toolbox from D). (H) Heatmap of results of product H2 reoptimization (using toolbox from D).

Scheme 3. Scope of Heteroaryl Triflates^a

^aIsolated yields from reactions on a 0.5 mmol scale under conditions A. Aryls from triflates are blue and aryls from halides are red. ^bHetAr-X was HetAr-Br. ^cNMR yield of a mixture of product and side products, see Supporting Information for details. ^dHetAr-X was HetAr-Cl.

testing.³⁴ Generally, μmol scale strikes the right balance between material cost and providing enough material for isolation and characterization (although the actual testing often needs only nanomoles³⁵ of material). Here, the solid reagent coated glass ChemBeads³⁶ in combination with our multi-variable catalytic system can quickly access arrays of biheteroaryls (Figure 1E) and efficiently improve the yields of target molecules via fast reoptimizations with a “Toolbox Plate” (Figure 1F–H).

Starting with standard conditions on 10 μmol scale (Figure 1A),³⁷ we evaluated all combinations of eight heteroaryl triflates (Figure 1B, A–H) and 12 heteroaryl halides (Figure 1C, 1–12) in a 96-well plate. The results are shown in a heatmap based on the product/internal standard ratio (Pdt/IS) (Figure 1E). These ratios are useful for comparing the same coupling under different conditions (as in the optimization plates Figure 1F–H), but the ratios between different pairs of coupling partners are not easily compared with each other. The standard conditions were generally effective, with 94% (90/96) of cross-coupled products observed by UV and MS analysis.

Inspired by the complementary reactivity of ttbtpy (N3) in couplings involving electron-poor heteroaryl halides (Scheme 2, Conditions B, 3f–3j), we developed a 96-well “Toolbox Plate”, which consists of known-useful variations of ligands, additives, and reductants. Such a Toolbox Plate could be useful

for improving the yields of target molecules from the initial HTE substrate scope screen. To maximize coverage, we included suboptimal, but promising ligands and additives from our initial optimization (Table 1), including amine ligands for Ni (Figure 1-D, N1–N4), phosphine ligands for Pd (P1 to P3), salt additives (KBr, KF, KCl, and KI), and reductants (Zn and Mn). Products detected under standard conditions (Figure 1-E) in well B3 (Pdt/IS = 0.01), C6 (Pdt/IS = 1.03), and H2 (Pdt/IS = 3.12) were chosen for reoptimization to represent three scenarios: low, medium, and relatively high initial yields.

When optimized using the Toolbox Plate, a significant increase in Pdt/IS ratios was observed for all these three products (Figure 1F–H). For challenging 2,2'-bipyridine B3, replacement of P1 and KBr in the standard conditions with P2 and KF was found to boost the Pdt/IS ratio of product B3 from 0.01 to 0.73. For benzoxazolyl quinoline C6, a combination of N4, P2, KCl, and Zn increased the yield of product C6 by more than 6-fold. Ligand-like pyrimidinyl imidazoquinoline H2 benefitted from a combination of N3, P1, KF, and Zn, which improved the Pdt/IS from 3 to 11. Finally, the syntheses of products B3, C6, and H2 under these reoptimized conditions³⁸ were conducted on a 0.5 mmol scale with 14%³⁹ (B3), 29% (C6), and 44% (H2) isolated yields, respectively. While further optimization could be conducted, these yields would be sufficient for testing³⁵ and can be obtained quickly with minimal material (2–4 mg of substrate per reaction).

In conclusion, we have developed a new cross-Ullmann approach that offers a general, reliable, and complementary solution to biheteroaryls. This strategy leverages abundant starting materials (heteroaryl halides and phenols), exhibits good functional group tolerance due to mild reductive conditions and demonstrates outstanding compatibility with a wide array of heterocycles. Indeed, the 62 examples in this manuscript include 38 different biheteroaryl cores. In addition, although the products in this manuscript are low molecular weight and quickly accessed from commercial compounds, 82% are new compounds (51/62). Finally, because these conditions are amenable to HTE format, the rapid synthesis of biheteroaryl collections and optimization of individual couplings can be accomplished quickly and with minimal material cost.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.1c10907>.

Additional tables of optimization data, detailed experimental procedures, characterization of products, and copies of product NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Daniel J. Weix – University of Wisconsin-Madison, Madison, Wisconsin 53706, United States; orcid.org/0000-0002-9552-3378; Email: dweix@wisc.edu

Authors

Kai Kang – University of Wisconsin-Madison, Madison, Wisconsin 53706, United States; orcid.org/0000-0002-7625-1661

Nathan L. Loud – University of Wisconsin-Madison, Madison, Wisconsin 53706, United States

Tarah A. DiBenedetto – University of Rochester, Rochester, New York 14627, United States; orcid.org/0000-0001-8861-0740

Complete contact information is available at:
<https://pubs.acs.org/10.1021/jacs.1c10907>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by the NIH (R01GM097243). The following instrumentation in the PBCIC was supported by Thermo Q Exactive Plus by NIH 1S10 OD020022-1, Bruker Avance III 400 by NSF CHE-1048642, Bruker Avance III 500 by a generous gift from Paul J. and Margaret M. Bender. We are grateful to Joe Barendt and Chiral Technologies for the donation of the SFC-A column used in this work.

REFERENCES

- (1) (a) Cepanec, I. *Synthesis of Biaryls*; Elsevier: New York, 2004; (b) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among US FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257–10274. (c) Wu, J. S.; Cheng, S. W.; Cheng, Y. J.; Hsu, C. S. Donor-Acceptor Conjugated Polymers Based on Multifused Ladder-Type Arenes for Organic Solar Cells. *Chem. Soc. Rev.* **2015**, *44*, 1113–1154. (d) Jin, Z. Muscarine, Imidazole, Oxazole and Thiazole Alkaloids. *Nat. Prod. Rep.* **2009**, *26*, 382–445. (e) Hansen, E. C.; Pedro, D. J.; Wotal, A. C.; Gower, N. J.; Nelson, J. D.; Caron, S.; Weix, D. J. New Ligands for Nickel Catalysis from Diverse Pharmaceutical Heterocycle Libraries. *Nat. Chem.* **2016**, *8*, 1126–1130.
- (2) (a) Badenock, J. C.; Gribble, G. W. Metal-Catalyzed Coupling with Heterocycles. In *Heterocyclic Chemistry in the 21st Century*; Advances in Heterocyclic Chemistry; Scriven, E. F. V., Ramsden, C. A., Eds.; Academic Press, 2016; Vol. 120, Chapter 4, pp 99–136. (b) Slagt, V. F.; de Vries, A. H. M.; de Vries, J. G.; Kellogg, R. M. Practical Aspects of Carbon–Carbon Cross-Coupling Reactions Using Heteroarenes. *Org. Process Res. Dev.* **2010**, *14*, 30–47. (c) Zhao, D.; You, J.; Hu, C. Recent Progress in Coupling of Two Heteroarenes. *Chem. - Eur. J.* **2011**, *17*, 5466–5492. (d) Revuelta, J.; Machetti, F.; Cicchi, S. In *Modern Heterocyclic Chemistry*; Wiley-VCH: Weinheim, Germany, 2011; (e) Rubtsov, A. E.; Malkov, A. V. Recent Advances in the Synthesis of 2,2'-Bipyridines and Their Derivatives. *Synthesis* **2021**, *53*, 2559–2569.
- (3) For selected examples, see ref 2 and (a) Almond-Thynne, J.; Blakemore, D. C.; Pryde, D. C.; Spivey, A. C. Site-Selective Suzuki-Miyaura Coupling of Heteroaryl Halides - Understanding the Trends for Pharmaceutically Important Classes. *Chem. Sci.* **2017**, *8*, 40–62. (b) Kim, S. H.; Rieke, R. D. Recent Advance in Heterocyclic Organozinc and Organomanganese Compounds; Direct Synthetic Routes and Application in Organic Synthesis. *Molecules* **2010**, *15*, 8006–8038. (c) Knochel, P.; Schade, M. A.; Bernhardt, S.; Manolikakes, G.; Metzger, A.; Piller, F. M.; Rohbogner, C. J.; Mosrin, M. Functionalization of Heterocyclic Compounds Using Polyfunctional Magnesium and Zinc Reagents. *Beilstein J. Org. Chem.* **2011**, *7*, 1261–1277. (d) Cordovilla, C.; Bartolome, C.; Martinez-Illarduya, J. M.; Espinet, P. The Stille Reaction, 38 Years Later. *ACS Catal.* **2015**, *5*, 3040–3053.
- (4) Commercial availability of heteroarene sources: HetAr-[M] (M = Zn, Mg, Li, Cu) (325), HetAr-[Sn] (292), HetAr-B(OH)₂ (2842); HetAr-[Si] (546), HetAr-OH (85739), HetAr-I (28720), HetAr-Br (225789), HetAr-Cl (299113). Source: eMolecules database, accessed 2019-04-08 using Elsevier REAXYS.
- (5) For synthesis of heteroaryl nucleophiles, see (a) Ila, H.; Baron, O.; Wagner, A. J.; Knochel, P. Preparation and Reactions of Heteroaryl Organomagnesium Compounds. *Chem. Lett.* **2006**, *35*, 2–7. (b) Knochel, P.; Schade, M. A.; Bernhardt, S.; Manolikakes, G.; Metzger, A.; Piller, F. M.; Rohbogner, C. J.; Mosrin, M. Functionalization of Heterocyclic Compounds Using Polyfunctional Magnesium and Zinc Reagents. *Beilstein J. Org. Chem.* **2011**, *7*, 1261–1277. (c) Snieckus, V. Directed Ortho Metalation. Tertiary Amide and O-Carbamate Directors in Synthetic Strategies for Polysubstituted Aromatics. *Chem. Rev.* **1990**, *90*, 879–933. (d) Lipshutz, B. H.; Hagen, W. Single-Flask Polyfunctionalization of the Imidazole Ring; a Streamlined Route to the Antitumor Agent Carmethizole. *Tetrahedron Lett.* **1992**, *33*, 5865–5868. (e) Cook, X. A. F.; de Gombert, A.; McKnight, J.; Pantaine, L. R. E.; Willis, M. C. The 2-Pyridyl Problem: Challenging Nucleophiles in Cross-Coupling Arylations. *Angew. Chem., Int. Ed.* **2021**, *60*, 11068–11091.
- (6) (a) Cox, P. A.; Reid, M.; Leach, A. G.; Campbell, A. D.; King, E. J.; Lloyd-Jones, G. C. Base-Catalyzed Aryl-B(OH)₂ Protodeboronation Revisited: From Concerted Proton Transfer to Liberation of a Transient Aryl Anion. *J. Am. Chem. Soc.* **2017**, *139*, 13156–13165. (b) Cox, P. A.; Leach, A. G.; Campbell, A. D.; Lloyd-Jones, G. C. Protodeboronation of Heteroaromatic, Vinyl, and Cyclopropyl Boronic Acids: Ph-Rate Profiles, Autocatalysis, and Disproportionation. *J. Am. Chem. Soc.* **2016**, *138*, 9145–9157. (c) Dick, G. R.; Woerly, E. M.; Burke, M. D. A General Solution for the 2-Pyridyl Problem. *Angew. Chem., Int. Ed.* **2012**, *51*, 2667–2672. (d) Billingsley, K. L.; Buchwald, S. L. A General and Efficient Method for the Suzuki-Miyaura Coupling of 2-Pyridyl Nucleophiles. *Angew. Chem., Int. Ed.* **2008**, *47*, 4695–4698. (e) Kinzel, T.; Zhang, Y.; Buchwald, S. L. A New Palladium Precatalyst Allows for the Fast Suzuki-Miyaura Coupling Reactions of Unstable Polyfluorophenyl and 2-Heteroaryl Boronic Acids. *J. Am. Chem. Soc.* **2010**, *132*, 14073–14075.
- (7) (a) Rossi, R.; Bellina, F.; Lessi, M.; Manzini, C. Cross-Coupling of Heteroarenes by C-H Functionalization: Recent Progress Towards Direct Arylation and Heteroarylation Reactions Involving Heteroarenes Containing One Heteroatom. *Adv. Synth. Catal.* **2014**, *356*, 17–117. (b) Alberico, D.; Scott, M. E.; Lautens, M. Aryl-Aryl Bond Formation by Transition-Metal-Catalyzed Direct Arylation. *Chem. Rev.* **2007**, *107*, 174–238. (c) Rossi, R.; Lessi, M.; Manzini, C.; Marianetti, G.; Bellina, F. Direct (Hetero) Arylation Reactions of (Hetero) Arenes as Tools for the Step-and Atom-Economical Synthesis of Biologically Active Unnatural Compounds Including Pharmaceutical Targets. *Synthesis* **2016**, *48*, 3821–3862. (d) Besson, T.; Fruit, C. Recent Developments in Microwave-Assisted Metal-Catalyzed C–H Functionalization of Heteroarenes for Medicinal Chemistry and Material Applications. *Synthesis* **2016**, *48*, 3879–3889. (e) Huang, H.-Y.; Benzai, A.; Shi, X.; Doucet, H. Effective Tools for the Metal-Catalyzed Regiodivergent Direct Arylations of (Hetero)-Arenes. *Chem. Rec.* **2021**, *21*, 343–356. (f) Bellina, F. Recent Developments in Pd-Catalyzed Direct Arylations of Heteroarenes with Aryl Halides. In *C-H Bond Activation and Catalytic Functionalization I*; Topics in Organometallic Chemistry; Dixneuf, P. H., Doucet, H., Eds.; Springer International: Cham, 2016; pp 77–102. (g) Hagui, W.; Periasamy, K.; Soule, J.-F. Synthesis of 2,2'-Bipyridines through Catalytic C–C Bond Formations from C–H Bonds. *Eur. J. Org. Chem.* **2021**, *2021*, 5388–5402. (h) Murakami, K.; Yamada, S.; Kaneda, T.; Itami, K. C–H Functionalization of Azines. *Chem. Rev.* **2017**, *117*, 9302–9332. (i) Verbitskiy, E. V.; Rusinov, G. L.; Chupakhin, O. N.; Charushin, V. N. Recent Advances in Direct C–H Functionalization of Pyrimidines. *Synthesis* **2018**, *50*, 193–210. (j) Song, A.-X.; Zeng, X.-X.; Ma, B.-B.; Xu, C.; Liu, F.-S. Direct (Hetero)Arylation of Heteroarenes Catalyzed by Unsymmetrical Pd-PEPPSI-NHC Complexes under Mild Conditions. *Organometallics* **2020**, *39*, 3524–3534.
- (8) (a) Yang, Y. D.; Lan, J. B.; You, J. S. Oxidative C-H/C-H Coupling Reactions between Two (Hetero)Arenes. *Chem. Rev.* **2017**, *117*, 8787–8863. (b) Choy, P. Y.; Wong, S. M.; Kapdi, A.; Kwong, F. Y. Recent Developments in Palladium-Catalysed Non-Directed Coupling of (Hetero) Arene C-H Bonds with C-Z (Z = B, Si, Sn,

- S, N, C, H) Bonds in Bi(Hetero)Aryl Synthesis. *Org. Chem. Front.* **2018**, *5*, 288–321. (c) Liu, C.; Zhang, H.; Shi, W.; Lei, A. W. Bond Formations between Two Nucleophiles: Transition Metal Catalyzed Oxidative Cross-Coupling Reactions. *Chem. Rev.* **2011**, *111*, 1780–1824. (d) Li, C. J. Cross-Dehydrogenative Coupling (CDC): Exploring C–C Bond Formations Beyond Functional Group Transformations. *Acc. Chem. Res.* **2009**, *42*, 335–344. (e) Yeung, C. S.; Dong, V. M. Catalytic Dehydrogenative Cross-Coupling: Forming Carbon–Carbon Bonds by Oxidizing Two Carbon–Hydrogen Bonds. *Chem. Rev.* **2011**, *111*, 1215–1292. (f) Han, W.; Ofial, A. R. No Detours: Palladium-Catalyzed Oxidative C–H/C–H Cross-Couplings of Heteroarenes. *Synlett* **2011**, *2011*, 1951–1955. (g) Bugaut, X.; Glorius, F. Palladium-Catalyzed Selective Dehydrogenative Cross-Couplings of Heteroarenes. *Angew. Chem., Int. Ed.* **2011**, *50*, 7479–7481. (h) Liu, C.; Yuan, J.; Gao, M.; Tang, S.; Li, W.; Shi, R.; Lei, A. Oxidative Coupling between Two Hydrocarbons: An Update of Recent C–H Functionalizations. *Chem. Rev.* **2015**, *115*, 12138–12204. (i) Bansal, S.; Shabade, A. B.; Punji, B. Advances in C(sp²)–H/C(sp²)–H Oxidative Coupling of (Hetero)Arenes Using 3d Transition Metal Catalysts. *Adv. Synth. Catal.* **2021**, *363*, 1998–2022.
- (9) (a) Markovic, T.; Roche, B. N.; Blakemore, D. C.; Mascitti, V.; Willis, M. C. Pyridine Sulfonates as General Nucleophilic Coupling Partners in Palladium-Catalyzed Cross-Coupling Reactions with Aryl Halides. *Chem. Sci.* **2017**, *8*, 4437–4442. (b) Markovic, T.; Murray, P. R. D.; Roche, B. N.; Shavnya, A.; Blakemore, D. C.; Willis, M. C. Heterocyclic Allylsulfonates as Latent Heteroaryl Nucleophiles in Palladium-Catalyzed Cross-Coupling Reactions. *J. Am. Chem. Soc.* **2018**, *140*, 15916–15923. (c) Cook, X. A. F.; Pantaine, L. R. E.; Blakemore, D. C.; Moses, I. B.; Sach, N. W.; Shavnya, A.; Willis, M. C. Base-Activated Latent Heteroaromatic Sulfonates as Nucleophilic Coupling Partners in Palladium-Catalyzed Cross-Coupling Reactions. *Angew. Chem., Int. Ed.* **2021**, *60*, 22461–22468.
- (10) Zhou, M.; Tsien, J.; Qin, T. Sulfur(IV)-Mediated Unsymmetrical Heterocycle Cross-Couplings. *Angew. Chem., Int. Ed.* **2020**, *59*, 7372–7376.
- (11) (a) Hilton, M. C.; Zhang, X.; Boyle, B. T.; Alegre-Requena, J. V.; Paton, R. S.; McNally, A. Heterobiaryl Synthesis by Contractive C–C Coupling via P(V) Intermediates. *Science* **2018**, *362*, 799–804. (b) Boyle, B. T.; Hilton, M. C.; McNally, A. Nonsymmetrical Bis-Azine Biaryls from Chloroazines: A Strategy Using Phosphorus Ligand-Coupling. *J. Am. Chem. Soc.* **2019**, *141*, 15441–15449. (c) Koniarczyk, J. L.; Greenwood, J. W.; Alegre-Requena, J. V.; Paton, R. S.; McNally, A. A Pyridine–Pyridine Cross-Coupling Reaction via Dearomatized Radical Intermediates. *Angew. Chem., Int. Ed.* **2019**, *58*, 14882–14886. (d) Che, Y.-Y.; Yue, Y.; Lin, L.-Z.; Pei, B.; Deng, X.; Feng, C. Palladium-Catalyzed Electrophilic Functionalization of Pyridine Derivatives through Phosphonium Salts. *Angew. Chem., Int. Ed.* **2020**, *59*, 16414–16419.
- (12) Examples of one-pot metalation + arylation for normal biaryls: (a) Ishiyama, T.; Itoh, Y.; Kitano, T.; Miyaura, N. Synthesis of Arylborationates via the Palladium(0)-Catalyzed Cross-Coupling Reaction of Tetra(Alkoxo)Diborons with Aryl Triflates. *Tetrahedron Lett.* **1997**, *38*, 3447–3450. (b) Baudoin, O.; Guénard, D.; Guéritte, F. Palladium-Catalyzed Borylation of *Ortho*-Substituted Phenyl Halides and Application to the One-Pot Synthesis of 2,2'-Disubstituted Biphenyls. *J. Org. Chem.* **2000**, *65*, 9268–9271. (c) Chow, W. K.; So, C. M.; Lau, C. P.; Kwong, F. Y. Palladium-Catalyzed Borylation of Aryl Mesylates and Tosylates and Their Applications in One-Pot Sequential Suzuki–Miyaura Biaryl Synthesis. *Chem. - Eur. J.* **2011**, *17*, 6913–6917.
- (13) Examples for cross-electrophile couplings for normal biaryl synthesis: (a) Amatore, M.; Gosmini, C. Efficient Cobalt-Catalyzed Formation of Unsymmetrical Biaryl Compounds and Its Application in the Synthesis of a Sartan Intermediate. *Angew. Chem., Int. Ed.* **2008**, *47*, 2089–2092. (b) Marina, N.; Lanterna, A. E.; Scaiano, J. C. Expanding the Color Space in the Two-Color Heterogeneous Photocatalysis of Ullmann C–C Coupling Reactions. *ACS Catal.* **2018**, *8*, 7593–7597. (c) Lv, L.; Qiu, Z.; Li, J.; Liu, M.; Li, C.-J. N₂H₄ as Traceless Mediator for Homo- and Cross- Aryl Coupling. *Nat. Commun.* **2018**, *9*, 4739. (d) Tang, J.; Liu, L. L.; Yang, S.; Cong, X.; Luo, M.; Zeng, X. Chemoselective Cross-Coupling between Two Different and Unactivated C(Aryl)–O Bonds Enabled by Chromium Catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 7715–7720. (e) Xiong, B.; Li, Y.; Wei, Y.; Kramer, S.; Lian, Z. Dual Nickel-/Palladium-Catalyzed Reductive Cross-Coupling Reactions between Two Phenol Derivatives. *Org. Lett.* **2020**, *22*, 6334–6338. (f) Nohira, I.; Chatani, N. Nickel-Catalyzed Cross-Electrophile Coupling between C(sp²)–F and C(sp²)–Cl Bonds by the Reaction of *Ortho*-Fluoro-Aromatic Amides with Aryl Chlorides. *ACS Catal.* **2021**, *11*, 4644–4649.
- (14) Examples for aryl-heteroaryl cross-coupling by one-pot, two-step metalation/cross-coupling: (a) Baudoin, O.; Cesario, M.; Guénard, D.; Guéritte, F. Application of the Palladium-Catalyzed Borylation/Suzuki Coupling (BSC) Reaction to the Synthesis of Biologically Active Biaryl Lactams. *J. Org. Chem.* **2002**, *67*, 1199–1207. (b) Molander, G. A.; Trice, S. L. J.; Dreher, S. D. Palladium-Catalyzed, Direct Boronic Acid Synthesis from Aryl Chlorides: A Simplified Route to Diverse Boronate Ester Derivatives. *J. Am. Chem. Soc.* **2010**, *132*, 17701–17703. (c) Molander, G. A.; Trice, S. L. J.; Kennedy, S. M. Scope of the Two-Step, One-Pot Palladium-Catalyzed Borylation/Suzuki Cross-Coupling Reaction Utilizing Bis-Boronic Acid. *J. Org. Chem.* **2012**, *77*, 8678–8688. (d) Molander, G. A.; Trice, S. L. J.; Kennedy, S. M.; Dreher, S. D.; Tudge, M. T. Scope of the Palladium-Catalyzed Aryl Borylation Utilizing Bis-Boronic Acid. *J. Am. Chem. Soc.* **2012**, *134*, 11667–11673.
- (15) Examples for cross-electrophile aryl-heteroaryl coupling: (a) Gosmini, C.; Lasry, S.; Nedelec, J. Y.; Perichon, J. Electrochemical Cross-Coupling between 2-Halopyridines and Aryl or Heteroaryl Halides Catalysed by Nickel-2,2'-Bipyridine Complexes. *Tetrahedron* **1998**, *54*, 1289–1298. (b) Gosmini, C.; Nedelec, J. Y.; Perichon, J. Electrosynthesis of Functionalized 2-Arylpyridines from Functionalized Aryl and Pyridine Halides Catalyzed by Nickel Bromide 2,2'-Bipyridine Complex. *Tetrahedron Lett.* **2000**, *41*, 5039–5042. (c) Sengmany, S.; Léonel, E.; Polissaint, F.; Nédélec, J.-Y.; Pipelier, M.; Thobie-Gautier, C.; Dubreuil, D. Preparation of Functionalized Aryl- and Heteroarylpyridazines by Nickel-Catalyzed Electrochemical Cross-Coupling Reactions. *J. Org. Chem.* **2007**, *72*, 5631–5636. (d) Miao, W.; Ni, C.; Xiao, P.; Jia, R.; Zhang, W.; Hu, J. Nickel-Catalyzed Reductive 2-Pyridination of Aryl Iodides with Difluoromethyl 2-Pyridyl Sulfone. *Org. Lett.* **2021**, *23*, 711–715. (e) Sengmany, S.; Vitu-Thiebaud, A.; Le Gall, E.; Condon, S.; Leonel, E.; Thobie-Gautier, C.; Pipelier, M.; Lebreton, J.; Dubreuil, D. An Electrochemical Nickel-Catalyzed Arylation of 3-Amino-6-Chloropyridazines. *J. Org. Chem.* **2013**, *78*, 370–379. (f) Sengmany, S.; Vasseur, S.; Lajnef, A.; Le Gall, E.; Leonel, E. Beneficial Effects of Electrochemistry in Cross-Coupling Reactions: Electroreductive Synthesis of 4-Aryl- or 4-Heteroaryl-6-Pyrrolylpyrimidines. *Eur. J. Org. Chem.* **2016**, *2016*, 4865–4871. (g) Dewanji, A.; Bülow, R. F.; Rueping, M. Photoredox/Nickel Dual-Catalyzed Reductive Cross Coupling of Aryl Halides Using an Organic Reducing Agent. *Org. Lett.* **2020**, *22*, 1611–1617. (h) Qin, J.; Zhu, S.; Chu, L. Dual Photoredox-/Palladium-Catalyzed Cross-Electrophile Couplings of Polyfluoroarenes with Aryl Halides and Triflates. *Organometallics* **2021**, *40*, 2246–2252. (i) Mirabi, B.; Marchese, A. D.; Lautens, M. Nickel-Catalyzed Reductive Cross-Coupling of Heteroaryl Chlorides and Aryl Chlorides. *ACS Catal.* **2021**, *11*, 12785–12793.
- (16) Examples for heteroaryl-heteroaryl cross-coupling by one-pot, two-step metalation/cross-coupling: (a) Billingsley, K. L.; Barder, T. E.; Buchwald, S. L. Palladium-Catalyzed Borylation of Aryl Chlorides: Scope, Applications, and Computational Studies. *Angew. Chem., Int. Ed.* **2007**, *46*, 5359–5363. (b) Shu, W.; Pellegatti, L.; Oberli, M. A.; Buchwald, S. L. Continuous-Flow Synthesis of Biaryls Enabled by Multistep Solid-Handling in a Lithiation/Borylation/Suzuki–Miyaura Cross-Coupling Sequence. *Angew. Chem., Int. Ed.* **2011**, *50*, 10665–10669. (c) Wang, L.; Cui, X.; Li, J.; Wu, Y.; Zhu, Z.; Wu, Y. Synthesis of Biaryls through a One-Pot Tandem Borylation/Suzuki–Miyaura Cross-Coupling Reaction Catalyzed by a Palladacycle. *Eur. J. Org. Chem.* **2012**, *2012*, 595–603. (d) Ji, H.; Wu, L.-Y.; Cai, J.-H.; Li, G.-R.; Gan, N.-N.; Wang, Z.-H. Room-Temperature Borylation and One-

Pot Two-Step Borylation/Suzuki–Miyaura Cross-Coupling Reaction of Aryl Chlorides. *RSC Adv.* **2018**, *8*, 13643–13648. (e) Duan, X.; Li, P.; Zhu, G.; Fu, C.; Chen, Q.; Huang, X.; Ma, S. Pd/Gorlos-Phos-Catalyzed Cross-Coupling between Two Different Aryl Chlorides in the Presence of B₂Pin₂ and Cytotoxicity Studies of the Products. *Org. Chem. Front.* **2018**, *5*, 3319–3323. (f) Xu, S. D.; Sun, F. Z.; Deng, W. H.; Hao, H.; Duan, X. H. One-Step Highly Selective Borylation/Suzuki Cross-Coupling of Two Distinct Aryl Bromides in Pure Water. *New J. Chem.* **2018**, *42*, 16464–16468.

(17) Reference 15 contains sporadic examples. For papers with a large number of examples, see: (a) Liao, L.-Y.; Kong, X.-R.; Duan, X.-F. Reductive Couplings of 2-Halopyridines without External Ligand: Phosphine-Free Nickel-Catalyzed Synthesis of Symmetrical and Unsymmetrical 2,2'-Bipyridines. *J. Org. Chem.* **2014**, *79*, 777–782. (b) Wu, T.-F.; Zhang, Y.-J.; Fu, Y.; Liu, F.-J.; Tang, J.-T.; Liu, P.; Toste, F. D.; Ye, B. Zirconium-Redox-Shuttled Cross-Electrophile Coupling of Aromatic and Heteroaromatic Halides. *Chem.* **2021**, *7*, 1963–1974.

(18) (a) Ge, S. Z.; Hartwig, J. F. Highly Reactive, Single-Component Nickel Catalyst Precursor for Suzuki–Miyaura Cross-Coupling of Heteroaryl Boronic Acids with Heteroaryl Halides. *Angew. Chem., Int. Ed.* **2012**, *51*, 12837–12841. (b) Wang, H.; Lorian, M. M.; Ackermann, L. Overcoming the Limitations of C–H Activation with Strongly Coordinating N-Heterocycles by Cobalt Catalysis. *Angew. Chem., Int. Ed.* **2016**, *55*, 10386–10390.

(19) There are several reports on the coupling of a single class of heteroaryl halide with a wide variety of aryl halides. See refs 15c–h.

(20) (a) Ackerman, L. K. G.; Lovell, M. M.; Weix, D. J. Multimetallic Catalyzed Cross-Coupling of Aryl Bromides with Aryl Triflates. *Nature* **2015**, *524*, 454–457. (b) Olivares, A. M.; Weix, D. J. Multimetallic Ni- and Pd-Catalyzed Cross-Electrophile Coupling to Form Highly Substituted 1,3-Dienes. *J. Am. Chem. Soc.* **2018**, *140*, 2446–2449. (c) Huang, L. B.; Ackerman, L. K. G.; Kang, K.; Parsons, A. M.; Weix, D. J. LiCl-Accelerated Multimetallic Cross-Coupling of Aryl Chlorides with Aryl Triflates. *J. Am. Chem. Soc.* **2019**, *141*, 10978–10983. (d) Kang, K.; Huang, L.; Weix, D. J. Sulfonate Versus Sulfonate: Nickel and Palladium Multimetallic Cross-Electrophile Coupling of Aryl Triflates with Aryl Tosylates. *J. Am. Chem. Soc.* **2020**, *142*, 10634–10640.

(21) Fagnou, K.; Lautens, M. Halide Effects in Transition Metal Catalysis. *Angew. Chem., Int. Ed.* **2002**, *41*, 26–47.

(22) Rodinkov, O. V.; Zhuravlyova, G. A.; Vaskova, E. A.; Platonov, I. A. Potassium Fluoride as a Selective Moisture Trapping Agent for SPE-TD-GC-FID Determination of Volatile Organic Compounds in Air. *Anal. Methods* **2015**, *7*, 458–465.

(23) 6-Methyl- and 6-methoxy-3-pyridyl triflates were used for the evaluation of heteroaryl halide scope because the methyl and methoxy group allows reliable determination of yield by ¹H NMR. Simple 3-pyridyl triflate showed similar reactivity to 6-methyl-3-pyridyl triflate in its reactions with both electron-rich and electron-poor heteroaryl bromides. This was illustrated by the syntheses of compounds 3aak and 3aal in Supporting Information, pp S51–S52.

(24) Reactions run with 4-bromo-3,5-dimethylisoxazole, 2-chloro-1-(4-fluorobenzyl)-1H-benzo[d]imidazole, and 6-bromo-2-methylthiazolo[4,5-b]pyridine resulted in no conversion of either starting material. This appears to be a catalyst poisoning effect because adding 4-bromo-3,5-dimethylisoxazole to a productive coupling pair shut down reactivity as well.

(25) Additional substrate pairs that resulted in low yields are listed in the Supporting Information, Scheme S10.

(26) Baumann, M.; Baxendale, I. R. An Overview of the Synthetic Routes to the Best Selling Drugs Containing 6-Membered Heterocycles. *Beilstein J. Org. Chem.* **2013**, *9*, 2265–2319.

(27) Qi, H.; Chen, L.; Liu, B.; Wang, X.; Long, L.; Liu, D. Synthesis and Biological Evaluation of Novel Pazopanib Derivatives as Antitumor Agents. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 1108–1110.

(28) The coupling of 6-methylpyridin-3-yl triflate (1a) with 3-pyridyl tosylate under the optimal conditions was also investigated. Because of the slow conversion of 3-pyridyl tosylate, the cross-product

3a was delivered in only 39% yield, accompanied by almost equal amount (32%) of dimerization product from 1a. We expect that further optimization could lead to a general cross-coupling reaction. See ref 29.

(29) For examples of cross-electrophile couplings between two different phenol derivatives, see ref 20 and (a) Tang, J.; Liu, L. L.; Yang, S.; Cong, X.; Luo, M.; Zeng, X. Chemoselective Cross-Coupling between Two Different and Unactivated C(Aryl)–O Bonds Enabled by Chromium Catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 7715–7720. (b) Xiong, B.; Li, Y.; Wei, Y.; Kramer, S.; Lian, Z. Dual Nickel-/Palladium-Catalyzed Reductive Cross-Coupling Reactions between Two Phenol Derivatives. *Org. Lett.* **2020**, *22*, 6334–6338.

(30) Our studies show that, compared with conditions A, conditions B convert electron-deficient heteroaryl bromides to the heteroaryl zinc reagent faster and are more selective for heteroaryl zinc reagent formation over homocoupling. See Supporting Information for more details.

(31) See ref 1e and (a) Constable, E. C.; Steel, P. J. N,N'-Chelating Biheteroaromatic Ligands; a Survey. *Coord. Chem. Rev.* **1989**, *93*, 205–223. (b) Kaes, C.; Katz, A.; Hosseini, M. W. Bipyridine: The Most Widely Used Ligand. A Review of Molecules Comprising at Least Two 2, 2'-Bipyridine Units. *Chem. Rev.* **2000**, *100*, 3553–3590.

(32) Note that reactions to form 3aad (35%) and 3aag (29%) were selective for cross-product and isolation from dimeric products was not challenging. The cause of the lower yields remains unclear.

(33) No examples were found in a substructure search using the CAS SciFinder database.

(34) (a) Buitrago Santanilla, A.; Regalado, E. L.; Pereira, T.; Shevlin, M.; Bateman, K.; Campeau, L.-C.; Schneeweis, J.; Berritt, S.; Shi, Z.-C.; Nantermet, P.; Liu, Y.; Helmy, R.; Welch, C. J.; Vachal, P.; Davies, I. W.; Cernak, T.; Dreher, S. D. Nanomole-Scale High-Throughput Chemistry for the Synthesis of Complex Molecules. *Science* **2015**, *347*, 49–53. (b) Krska, S. W.; DiRocco, D. A.; Dreher, S. D.; Shevlin, M. The Evolution of Chemical High-Throughput Experimentation To Address Challenging Problems in Pharmaceutical Synthesis. *Acc. Chem. Res.* **2017**, *50*, 2976–2985. (c) Perera, D.; Tucker, J. W.; Brahmabhatt, S.; Helal, C. J.; Chong, A.; Farrell, W.; Richardson, P.; Sach, N. W. A Platform for Automated Nanomole-Scale Reaction Screening and Micromole-Scale Synthesis in Flow. *Science* **2018**, *359*, 429–434. (d) Mennen, S. M.; Alhambra, C.; Allen, C. L.; Barberis, M.; Berritt, S.; Brandt, T. A.; Campbell, A. D.; Castañón, J.; Cherney, A. H.; Christensen, M.; Damon, D. B.; Eugenio de Diego, J.; García-Cerrada, S.; García-Losada, P.; Haro, R.; Janey, J.; Leitch, D. C.; Li, L.; Liu, F.; Lobben, P. C.; MacMillan, D. W. C.; Magano, J.; McInturff, E.; Monfette, S.; Post, R. J.; Schultz, D.; Sitter, B. J.; Stevens, J. M.; Strambeanu, I. I.; Twilton, J.; Wang, K.; Zajac, M. A. The Evolution of High-Throughput Experimentation in Pharmaceutical Development and Perspectives on the Future. *Org. Process Res. Dev.* **2019**, *23*, 1213–1242.

(35) Ref 34 and Biyani, S. A.; Moriuchi, Y. W.; Thompson, D. H. Advancement in Organic Synthesis Through High Throughput Experimentation. *Chem. Methods* **2021**, *1*, 323–339.

(36) (a) Tu, N. P.; Dombrowski, A. W.; Goshu, G. M.; Vasudevan, A.; Djuric, S. W.; Wang, Y. High-Throughput Reaction Screening with Nanomoles of Solid Reagents Coated on Glass Beads. *Angew. Chem., Int. Ed.* **2019**, *58*, 7987–7991. (b) Aguirre, A. L.; Loud, N. L.; Johnson, K. A.; Weix, D. J.; Wang, Y. ChemBead Enabled High-Throughput Cross-Electrophile Coupling Reveals a New Complementary Ligand. *Chem. - Eur. J.* **2021**, *27*, 12981–12986.

(37) Solid reagent coated ChemBeads are dispensed with a suite of calibrated scoops (5–200 mg) by volume, see ref 36 for details. The ChemBeads used in this study (10% w/w for salt additives and 5% w/w for reductants) were added by a 15 mg-scoop. See Supporting Information for details.

(38) In these 0.5 mmol scale reactions, 2.0 equiv of reductant and 1.0 or 2.0 equiv of salt additives were used. See Supporting Information for details.

(39) NMR yield from a mixture of product and side products.