





1/21: I) Do the following (VSEPR) problems from your text (Miessler et al. (2014)): Chapter 3: #8, 9f-i, 20, 29.

3.8 Give Lewis dot structures and sketch the shapes of the following:



a. SeCl_4





b. I_3^-





c. PSCl_3





d. IF_4^-





e. PH_2^-





f. TeF_4^{2-}





g. N_3^-



h. SeOCl_4





i. PH_4^+





3.9 Give Lewis dot structures and sketch the shapes of the following.



f. $\text{IO}(\text{OH})_5$





g. SOCl_2





h. ClOF_4^-




i. XeO_2F_2



3.20 Predict and sketch the structure of the (as yet) hypothetical ion IF_3^{2-} .



3.29 Sketch the most likely structure of PCl_3Br_2 and explain your reasoning.



Answer. Bromine is more electropositive than chlorine. Thus, by Bent's rule, the bromines will bond to the hybrid orbitals with greater s -character (the equatorial sp^2 ones) first. \square

- II) Assign the symmetry point group to the 13 ions and molecules in problems #8, 9f-i in Chapter 3 of your text.

3.8**a. SeCl₄**

Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes.

Therefore, SeCl₄ is of the C_{2v} point group. ☐

b. I₃⁻

Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes.

Therefore, I₃⁻ is of the C_{2v} point group. ☐

c. PSCl₃

Answer. Not low or high symmetry. Has a C_3 axis. No perpendicular C_2 axes. No σ_h . Has three σ_v planes all offset by 60°.

Therefore, PSCl₃ is of the C_{3v} point group. ☐

d. IF₄⁻

Answer. Not low or high symmetry. Has a C_4 axis. Has 4 perpendicular C_2 axes. Has σ_h .

Therefore, IF₄⁻ is of the D_{4h} point group. ☐

e. PH₂⁻

Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes.

Therefore, PH₂⁻ is of the C_{2v} point group. ☐

f. TeF₄²⁻

Answer. Not low or high symmetry. Has a C_4 axis. Has 4 perpendicular C_2 axes. Has σ_h .

Therefore, TeF₄²⁻ is of the D_{4h} point group. ☐

g. N₃⁻

Answer. N₃⁻ is of the $D_{\infty h}$ point group. ☐

h. SeOCl₄

Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes.

Therefore, SeOCl₄ is of the C_{2v} point group. ☐

i. PH₄⁺

Answer. PH₄⁺ is of the T_d point group. ☐

3.9**f. IO(OH)₅**

Answer. Not low or high symmetry. Has a C_4 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes and two perpendicular σ_d planes.

Therefore, IO(OH)₅ is of the C_{4v} point group. ☐

g. SOCl₂

Answer. SOCl₂ is of the C_s point group. ☐

h. ClOF₄⁻

Answer. Not low or high symmetry. Has a C_4 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes and two perpendicular σ_d planes.

Therefore, ClOF₄⁻ is of the C_{4v} point group. ☐


i. XeO₂F₂

Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes.

Therefore, XeO₂F₂ is of the C_{2v} point group. ☐


III) Assign the symmetry point group of the following molecules and objects. Ignore the H atoms in (a), (e), and (g). Note that (e) has pseudooctahedral geometry and (g) is square-planar.

a) The molecule pictured below.



Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. Has a σ_h . Therefore, the above molecule is of the C_{2h} point group. ☐

b) The molecule pictured below.



Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. Has a σ_h . Therefore, the above molecule is of the C_{2h} point group. ☐

c) POCl_3

Answer. Not low or high symmetry. Has a C_3 axis. No perpendicular C_2 axes. No σ_h . Has three σ_v planes all offset by 60° . Therefore, POCl_3 is of the C_{3v} point group. ☐

d) Tennis ball (including the seam)

Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes. Therefore, a tennis ball is of the C_{2v} point group. ☐

e) $\text{trans}[\text{CrCl}_2(\text{H}_2\text{O})_4]^+$

Answer. Not low or high symmetry. Has a C_4 axis. Has 4 perpendicular C_2 axes. Has σ_h . Therefore, $\text{trans}[\text{CrCl}_2(\text{H}_2\text{O})_4]^+$ is of the D_{4h} point group. ☐

f) 1,3,5-trichlorobenzene.

Answer. Not low or high symmetry. Has a C_3 axis. Has 3 perpendicular C_2 axes. Has σ_h . Therefore, 1,3,5-trichlorobenzene is of the D_{3h} point group. ☐

g) $\text{trans-Pt}(\text{NH}_3)_2\text{Cl}_2$

Answer. Not low or high symmetry. Has a C_2 axis. Has 2 perpendicular C_2 axes. Has σ_h . Therefore, TeF_4^{2-} is of the D_{2h} point group. ☐

h) SF_5Cl

Answer. Not low or high symmetry. Has a C_4 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes and two perpendicular σ_d planes.

Therefore, SF_5Cl is of the C_{4v} point group. □

i) BFClBr

Answer. BFClBr is of the C_s point group. □

j) PF_2^+

Answer. Not low or high symmetry. Has a C_2 axis. No perpendicular C_2 axes. No σ_h . Has two perpendicular σ_v planes.

Therefore, PF_2^+ is of the C_{2v} point group. □

- IV) In the octahedral ion FeF_6^{3-} , what symmetry elements are destroyed if two *trans* F ions are moved away from the Fe^{3+} center in an equidistant fashion?

Answer. If the described change is made, the point group changes from O_h to D_{4h} . In this change, every C_3 and S_6 axis, two of the three C_4 axes, and every σ_d that does not contain the axis along which the F ions are stretched are destroyed. \square